

# Formalisation of CW-complexes

Hannah Scholz

Born 21<sup>st</sup> July, 2003 in Bonn, Germany

22<sup>nd</sup> August, 2024

Bachelor's Thesis Mathematics

Advisor: Prof. Dr. Floris van Doorn

Second Advisor: Prof. Dr. X Y

MATHEMATISCHES INSTITUT

MATHEMATISCH-NATURWISSENSCHAFTLICHE FAKULTÄT DER  
RHEINISCHEN FRIEDRICH-WILHELMS-UNIVERSITÄT BONN



# Contents

<b>1. Introduction</b>	<b>5</b>
1.1. First section . . . . .	5
1.1.1. Subsections . . . . .	5
1.2. How to cite . . . . .	5
<b>2. One</b>	<b>7</b>
2.1. How to cross reference . . . . .	7
2.2. How to create Index . . . . .	7
2.3. How to create symbol index . . . . .	7
<b>3. Definition</b>	<b>9</b>
3.1. Definition of a CW-complexes . . . . .	9
<b>4. Product</b>	<b>15</b>
4.1. K-spaces and the k-ification . . . . .	15
4.2. The product of CW-complexes . . . . .	20
<b>Appendix</b>	<b>21</b>
A. a . . . . .	21
B. b . . . . .	21
<b>Symbol Index</b>	<b>23</b>
<b>Index</b>	<b>25</b>
<b>Bibliography</b>	<b>27</b>



# 1. Introduction

Here you have your introduction. This template is mainly based on Felix Boes Master thesis you can find it here [https://github.com/felixboes/masters\\_thesis/tree/master](https://github.com/felixboes/masters_thesis/tree/master)

## 1.1. First section

Congratulations you have created your first section

### 1.1.1. Subsections

If you need to divide your thesis use subsections.

You can iterate this with subsub-sections if you want. If you want to do this you have to change the depth of the subsub-sections in the main.

If you want to create a subsection, which does not appear in the contents table use "subsection\*"

## 1.2. How to cite

In this section we will discuss how to cite.

Just add your reference in masterthesis\_\_your\_\_name\_\_bibliography.bib

Then use this line [Abh05] or [Dis06]



## 2. One

In this chapter we will learn some useful tools.

### 2.1. How to cross reference

In this section we will learn to cross reference.

For this just use command 1.1. You previously need to create a label.

You can also reference equations in this way

$$\langle v, \operatorname{Re} Av \rangle = \langle v, U^* DU v \rangle = \langle Uv, DUv \rangle \geq \lambda_1 \|Uv\|^2 = \lambda_1 \|v\|^2 \quad (2.1)$$

**Lemma 2.1.1.** *Let  $A \in \mathbb{C}^{M \times M}$  be diagonally dominant then  $A$  is invertible.*

This equations can be accessed by 2.1 and 2.1.1

### 2.2. How to create Index

In this section we will learn to add elements to the index.

Just use the command as in the example.

**Definition 2.2.1.** A vectorspace is...

### 2.3. How to create symbol index

In this section we will learn to add elements to the symbol index.

Just use the command as in the example.





### 3. Definition

In this chapter we will introduce CW-complexes and prove basic facts about them.

#### 3.1. Definition of a CW-complexes

The modern definition of a CW-complex is the following:

**Definition 3.1.1.** Let  $X$  be a topological space. A *CW-complex* on  $X$  is a filtration  $X_0 \subseteq X_1 \subseteq X_2 \subseteq \dots$  such that

- (i) For every  $n \geq 0$  there is a pushout of topological spaces

$$\begin{array}{ccc} \coprod_{i \in I_n} S_i^{n-1} & \xrightarrow{\coprod_{i \in I_n} q_i^n} & X_{n-1} \\ \downarrow \coprod_{i \in I_n} j_i & & \downarrow \\ \coprod_{i \in I_n} D_i^n & \xrightarrow{\coprod_{i \in I_n} Q_i^n} & X_n \end{array}$$

where  $I_n$  is any indexing set and  $j_i: S_i^{n-1} \rightarrow D_i^n$  is the usual inclusion for every  $i \in I_n$ .

- (ii) We have  $X = \bigcup_{n \geq 0} X_n$ .
- (iii)  $X$  has weak topology, i.e.  $A \subseteq X$  is open  $\iff A \cap X_n$  is open in  $X_n$  for every  $n$ .

$X_n$  is called the *n-skeleton*. An element  $e^n \in \pi_0(X_n \setminus X_{n-1})$  is called an (*open*) *n-cell*.  $Q_i^n$  is called a *characteristic map*.

In this thesis we will however focus on the historical definition of CW-complexes first presented by Whitehead which can be found in [Whi18].

**Definition 3.1.2.** Let  $X$  be a Hausdorff space. A *CW-complex* on  $X$  consists of a family of indexing sets  $(I_n)_{n \in \mathbb{N}}$  and a family of maps  $(Q_i^n: D_i^n \rightarrow X)_{n \geq 0, i \in I_n}$  s.t.

- (i)  $Q_i^n|_{\text{int}(D_i^n)}: \text{int}(D_i^n) \rightarrow Q_i^n(\text{int}(D_i^n))$  is a homeomorphism. We call  $e_i^n := Q_i^n(\text{int}(D_i^n))$  an (*open*) *n-cell* (or a cell of dimension  $n$ ) and  $\bar{e}_i^n := Q_i^n(D_i^n)$  a *closed n-cell*.
- (ii) For all  $n, m \in \mathbb{N}$ ,  $i \in I_n$  and  $j \in I_m$  where  $(n, i) \neq (m, j)$  the cells  $e_i^n$  and  $e_j^m$  are disjoint.
- (iii) For each  $n \in \mathbb{N}$ ,  $i \in I_n$ ,  $Q_i^n(\partial D_i^n)$  is contained in the union of a finite number of closed cells of dimension less than  $n$ .

### 3.1. Definition of a CW-complexes

---

(iv)  $A \subseteq X$  is closed iff  $Q_i^n(D_i^n) \cap A$  is closed for all  $n \in \mathbb{N}$  and  $i \in I_n$ .

(v)  $\bigcup_{n \geq 0} \bigcup_{i \in I_n} Q_i^n(D_i^n) = X$ .

We call  $Q_i^n$  a *characteristic map* and  $\partial e_i^n := Q_i^n(\partial D_i^n)$  the *edge of the  $n$ -cell* for any  $i$  and  $n$ . Additionally we define  $X_n := \bigcup_{m < n+1} \bigcup_{i \in I_m} \bar{e}_i^m$  and call it the  *$n$ -skeleton* of  $X$  for  $-1 \leq n \leq \infty$ .

For the rest of the chapter let  $X$  be a CW-complex.

**Remark 3.1.3.** Property (iii) in the above definition is called *closure finiteness*. Property (iv) is called *weak topology*. Whitehead named CW-complexes or long *closure finite complexes with weak topology* after these two properties [Whi18].

Let us first answer the obvious question about the two definitions:

**Proposition 3.1.4.** *Definition 3.1.1 and 3.1.2 are equivalent.*

The proof to this proposition is long, tedious and not relevant to this thesis so we will skip it here. It can be found as the proof of Proposition A.2. in [Hat01]. From here on the term CW-Complex will always refer to the older definition 3.1.2. As such keep in mind that throughout this thesis any CW-complex will by definition be assumed to be Hausdorff.

**Remark 3.1.5.** The name *open  $n$ -cell* and the notation  $\partial e_i^n$  can be confusing as an open  $n$ -cell is not necessarily open and  $\partial e_i^n$  is not necessarily the boundary of  $\bar{e}_i^n$ .

But at least the notion of a closed  $n$ -cell makes sense:

**Lemma 3.1.6.**  *$\bar{e}_i^n$  is compact and closed for every  $n \in \mathbb{N}$  and  $i \in I_n$ . Similarly  $\partial e_i^n$  is compact and closed for every  $n \in \mathbb{N}$  and  $i \in I_n$ .*

*Proof.*  $D_i^n$  is compact. Therefore its image  $Q_i^n(D_i^n) = \bar{e}_i^n$  is compact as well. In a Hausdorff space any compact set is closed. Thus  $\bar{e}_i^n$  is closed. The proof for  $\partial e_i^n$  works in the same way.  $\square$

And luckily the following is also true:

**Lemma 3.1.7.**  *$\overline{e_i^n} = \bar{e}_i^n$  for every  $n \in \mathbb{N}$  and  $i \in I_n$ .*

*Proof.* Since  $e_i^n \subseteq \bar{e}_i^n$  and  $\bar{e}_i^n$  is closed by the lemma above, the left inclusion is trivial. So let us show now that  $\bar{e}_i^n \subseteq \overline{e_i^n}$ . This statement can be rewritten as  $Q_i^n(\bar{B}_i^n) \subseteq \overline{Q_i^n(B_i^n)}$ . It is generally true for any continuous map that the closure of the image is contained in the image of the closure. Thus we are done.  $\square$

Now let us define what it means for a CW-complex to be finite:

**Definition 3.1.8.** Let  $X$  be a CW-complex. We call  $X$  *of finite type* if there are only finitely many cells in each dimension, i.e. if  $I_n$  is finite for all  $n \in \mathbb{N}$ .  $X$  is said to be *finite dimensional* if there is an  $n \in \mathbb{N}$  such that  $X = X_n$ . Finally,  $X$  is called *finite* if it is of finite type and finite dimensional.

If we already know that the CW-complex we want to construct will be finite or of finite type we can relax some of the conditions:

**Remark 3.1.9.**

- (i) For a CW-complex of finite type condition (iii) in definition 3.1.2 follows from the following: For each  $n \in \mathbb{N}$ ,  $i \in I_n$   $Q_i^n(\partial D_i^n)$  is contained in  $\bigcup_{m \leq n-1} \bigcup_{i \in I_m} e_i^m$ .
- (ii) Additionally for a finite CW-complex condition (iv) in definition 3.1.2 is follows from the other conditions.

*Proof.* Let us begin with statement (i). Take  $n \in \mathbb{N}$  and  $i \in I_n$ . We need to show that  $Q_i^n(\partial D_i^n)$  is contained in a finite number of cells of a lower dimension. But by assumption we have  $Q_i^n(\partial D_i^n) \subseteq \bigcup_{m \leq n-1} \bigcup_{i \in I_m} e_i^m$  which in this case is made up of finitely many cells. Now we can move on to statement (ii). We need to prove condition (iv) of definition 3.1.2, i.e.

$$A \subseteq X \text{ is closed} \iff \bar{e}_i^n \cap A \text{ is closed for all } n \in \mathbb{N} \text{ and } i \in I_n.$$

For the forward direction notice that  $\bar{e}_i^n \cap A$  is just the intersection of two closed sets by assumption and lemma 3.1.6. As such it is closed. For the backward direction take an  $A \subseteq X$  such that  $\bar{e}_i^n$  is closed for all  $n \in \mathbb{N}$  and  $i \in I_n$ . We need to show that  $A$  is closed. But using condition (v) of definition 3.1.2 we get

$$A = A \cap \bigcup_{n \geq 0} \bigcup_{i \in I_n} \bar{e}_i^n = \bigcup_{n \geq 0} \bigcup_{i \in I_n} (A \cap \bar{e}_i^n)$$

which by assumption is a finite union of closed sets, making  $A$  closed.  $\square$

We can also think about the  $n$ -skeletons as being made up of open cells:

**Lemma 3.1.10.**  $X_n = \bigcup_{m < n+1} \bigcup_{i \in I_m} e_i^m$  for every  $-1 \leq n \leq \infty$ .

*Proof.* We show this by induction over  $-1 \leq n \leq \infty$ . For the base case assume that  $n = -1$ . Then we get  $X_n = \bigcup_{m < 0} \bigcup_{i \in I_m} \bar{e}_i^m = \emptyset = \bigcup_{m < 0} \bigcup_{i \in I_m} e_i^m$ .

For the induction step assume that the statement is true for  $n$ . We now show that it also holds for  $n+1$ .

$$\begin{aligned} X_{n+1} &= \bigcup_{m < n+2} \bigcup_{i \in I_m} \bar{e}_i^m \\ &= \bigcup_{i \in I_{n+1}} \bar{e}_i^{n+1} \cup \bigcup_{m < n+1} \bigcup_{i \in I_m} \bar{e}_i^m \\ &= \bigcup_{i \in I_{n+1}} \bar{e}_i^{n+1} \cup X_n \\ &\stackrel{(1)}{=} \bigcup_{i \in I_{n+1}} \bar{e}_i^{n+1} \cup \bigcup_{m < n+1} \bigcup_{i \in I_m} e_i^m \\ &= \bigcup_{i \in I_{n+1}} e_i^{n+1} \cup \bigcup_{i \in I_{n+1}} \partial e_i^{n+1} \cup \bigcup_{m < n+1} \bigcup_{i \in I_m} e_i^m \\ &\stackrel{(2)}{=} \bigcup_{i \in I_{n+1}} e_i^{n+1} \cup \bigcup_{m < n+1} \bigcup_{i \in I_m} e_i^m \\ &= \bigcup_{m < n+2} \bigcup_{i \in I_m} e_i^m \end{aligned}$$

### 3.1. Definition of a CW-complexes

---

Where (1) holds by induction and (2) holds by closure finiteness (property (iii) in definition 3.1.2).

Now we can move on to the case  $n = \infty$ .

$$\begin{aligned}
 X_\infty &= \bigcup_{m < \infty+1} \bigcup_{i \in I_m} \bar{e}_i^m \\
 &= \bigcup_{m < \infty+1} \bigcup_{l < m+1} \bigcup_{i \in I_l} \bar{e}_i^l \\
 &= \bigcup_{m < \infty+1} X_m \\
 &\stackrel{(1)}{=} \bigcup_{m < \infty+1} \bigcup_{l < m+1} \bigcup_{i \in I_l} e_i^l \\
 &= \bigcup_{m < \infty+1} \bigcup_{i \in I_m} \bar{e}_i^m
 \end{aligned}$$

Where (1) holds by induction. □

This also enables us to write  $X$  as a union of open cells:

**Corollary 3.1.11.**  $\bigcup_{n \geq 0} \bigcup_{i \in I_n} e_i^n = X$ .

When we want to show that a set  $A \subseteq X$  is closed the weak topology (property (iv) in 3.1.2) lets us reduce that question to an individual cell. It is then often convenient to do strong induction over the dimension of the cell. We now want to prove a lemma that makes this repeated process easier. We first need the following:

**Lemma 3.1.12.** *Let  $A \subseteq X$  be a set and  $n$  a natural number. Assume that for every  $m \leq n$  and  $j \in I_m$  the intersection  $A \cap \bar{e}_j^m$  is closed. Then  $A \cap \partial e_j^{n+1}$  is closed for every  $j \in I_{n+1}$ .*

*Proof.* By closure finiteness of  $X$  (property (iii) in 3.1.2) there is a set  $E$  of cells of dimension lower than  $n+1$  such that  $\partial e_j^{n+1} \subseteq \bigcup_{e \in E} \bar{e}$ . This gives us

$$A \cap \partial e_j^{n+1} = A \cap \bigcup_{e \in E} \bar{e} \cap \partial e_j^{n+1} = \bigcup_{e \in E} (A \cap \bar{e}) \cap \partial e_j^{n+1}.$$

$\bigcup_{e \in E} (A \cap \bar{e})$  is closed as a finite union of sets that are by assumption closed and  $\partial e_j^{n+1}$  is closed by lemma 3.1.6. Therefore the intersection is also closed. □

Now we can move on to the lemma that we actually want. We can think of this lemma as being a weaker condition than the weak topology i.e. property (iv) in 3.1.2.

**Lemma 3.1.13.** *Let  $A \subseteq X$  be a set such that for every  $n > 0$  and  $j \in I_n$  either  $A \cap e_j^n$  or  $A \cap \bar{e}_j^n$  is closed. Then  $A$  is closed.*

*Proof.* Since  $X$  has weak topology, it is enough to show that  $A \cap \bar{e}_j^n$  is closed for every  $n \in \mathbb{N}$  and  $i \in I_n$ . We show this by doing strong induction over  $n$ . For the base case  $n = 0$  notice that  $\bar{e}_j^0$  is a singleton and the intersection with a singleton is either that singleton or empty. As such the intersection is closed in both cases.

Now let us move on to the induction step. Assume that for every  $m \leq n$  the statement already holds. We now need to show it for  $n+1$ . By assumption either  $A \cap e_j^{n+1}$  or  $A \cap \bar{e}_j^{n+1}$  is closed. The second case is just immediately what we wanted to show.

In the first case we can use that  $A \cap \bar{e}_j^{n+1} = (A \cap \partial e_j^{n+1}) \cup (A \cap e_j^{n+1})$ . The left part of the union is closed by lemma 3.1.12 applied to the induction hypothesis. The right part of the union is closed by the assumption of our case. The union is therefore also closed.  $\square$

Another fact that can be quite helpful is a version of closure finiteness using open cells:

**Lemma 3.1.14.** *For each  $n \in \mathbb{N}$  and  $i \in I_n$   $\partial e_i^n$  is contained in the union of a finite number of open cells of dimension less than  $n$ .*

*Proof.* We show this by doing strong induction on  $n$ . For the base case  $n = 0$  notice that  $\partial e_i^0$  is empty.

Moving on to the induction step assume that the statement holds for all  $m \leq n$ . We need to show that it also holds for  $n+1$ . By closure finiteness there is a finite set  $E$  of cells of dimension less than  $n+1$  such that  $\partial e_i^{n+1} \subseteq \bigcup_{e \in E} \bar{e}$ . If we can show that for every  $e \in E$  there is a finite set  $E_e$  of cells of dimension less than  $n+1$  such that  $\bar{e} \subseteq \bigcup_{e' \in E_e} e'$ , we would then be done since  $\partial e_i^{n+1} \subseteq \bigcup_{e \in E} \bar{e} \subseteq \bigcup_{e \in E} \bigcup_{e' \in E_e} e'$ .

So take  $e \in E$ . By the induction hypothesis there is a finite set  $E'_e$  of cells of a lesser dimension than that of  $e$  such that  $\partial e \subseteq \bigcup_{e' \in E'_e} e'$ . This gives us  $\bar{e} = \partial e \cup e \subseteq (\bigcup_{e' \in E'_e} e') \cup e$  which finishes the proof.  $\square$



## 4. Product

In this chapter we will talk about the product.

### 4.1. K-spaces and the k-ification

Before we can move on to discuss the product of CW-complexes we need to discuss its topology. Therefore we will study k-spaces and the k-ification in this section.

A k-space or also called a compactly generated space is defined for our purposes as follows. Note that we mean quasi-compactness when talking about compactness.

**Definition 4.1.1.** Let  $X$  be a topological space. We call  $X$  a *k-space* if

$A \subseteq X$  is open  $\iff$  for all compact sets  $C \subseteq X$  the intersection  $A \cap C$  is open in  $C$ .

There are a lot of different definitions in the literature. The most popular ones all agree on Hausdorff spaces. An overview of these different notions can be found on Wikipedia [Wik24].

It will also be helpful to characterise k-spaces via closed sets:

**Lemma 4.1.2.** Let  $X$  be a topological space.  $X$  is a k-space iff

$A \subseteq X$  is closed  $\iff$  for all compact sets  $C \subseteq X$  the intersection  $A \cap C$  is closed in  $C$ .

*Proof.* We only show that forward direction as the backward direction follows in the same way. Of the equivalence that we now need to show the forward direction is trivial. Thus let  $A \subseteq X$  be a set such that for all compact sets  $C \subseteq X$  the intersection  $A \cap C$  is closed in  $C$ . It is enough to show that  $A^c$  is open. By definition of the k-space that is the case if for every compact set  $C \subseteq X$  the intersection  $A^c \cap C$  is open in  $C$ . Take any compact  $C \subseteq X$ . By assumption  $A \cap C$  is closed in  $C$ . Since  $A \cap C$  is the complement of  $A^c \cap C$  in  $C$ , this immediately gives us that  $A^c \cap C$  is open in  $C$ .  $\square$

We also define a way to make any topological space into a k-space which we call the k-ification:

**Definition 4.1.3.** Let  $X$  be a topological space. We can define another topological space  $X_c$  on the same set by setting

$A \subseteq X_c$  is open  $\iff$  for all compact sets  $C \subseteq X$  the intersection  $A \cap C$  is open in  $C$ .

We call  $X_c$  the *k-ification* of  $X$ .

It is easy to see that this gives us a finer topology:

**Lemma 4.1.4.**  $A \subseteq X$  is open  $\implies A \subseteq X_c$  is open.

Again it is useful to characterise the closed sets in the k-ification:

**Lemma 4.1.5.**  $A \subseteq X_c$  is closed  $\iff A \cap C$  is closed in  $C$  for all compact sets  $C \subseteq X$ .

*Proof.* Completely analogue to the proof of lemma 4.1.2.  $\square$

To show that the k-ification actually fulfils its purpose of turning any space into a k-space, we first need the following lemma:

**Lemma 4.1.6.**  $A \subseteq X$  is compact  $\iff A \subseteq X_c$  is compact.

*Proof.* For the backward direction notice that lemma 4.1.4 is another way of stating that the map  $\text{id}: X_c \rightarrow X$  is continuous. As the image of a compact set under a continuous map, that makes  $A \subseteq X$  compact.

For the forward direction take  $A \subseteq X$  compact. To show that  $A \subseteq X_c$  is compact, take an open cover  $(U_i)_{i \in \iota}$  of  $A$  in  $X_c$ . For every  $i \in \iota$  there is by definition of the k-ification an open set  $V_i \subseteq X$  such that  $V_i \cap A = U_i \cap A$ .  $(V_i)_{i \in \iota}$  is an (open) cover of  $A$  in  $X$ :

$$A = A \cap \bigcup_{i \in \iota} U_i = \bigcup_{i \in \iota} (A \cap U_i) = \bigcup_{i \in \iota} (A \cap V_i) = A \cap \bigcup_{i \in \iota} V_i \subseteq \bigcup_{i \in \iota} V_i.$$

Thus there is a finite subcover  $(V_i)_{i \in \iota'}$  of  $A$  in  $X$ .  $(U_i)_{i \in \iota'}$  is now a finite subcover of  $A$  in  $X_c$ :

$$A = A \cap \bigcup_{i \in \iota'} V_i = \bigcup_{i \in \iota'} (A \cap V_i) = \bigcup_{i \in \iota'} (A \cap U_i) = A \cap \bigcup_{i \in \iota'} U_i \subseteq \bigcup_{i \in \iota'} U_i.$$

$\square$

Now we are ready to move on to the promised lemma:

**Lemma 4.1.7.**  $X_c$  is a k-space for every topological space  $X$ .

*Proof.* We need to show that a set  $A \subseteq X_c$  is open iff  $A \cap C$  is open in  $C$  for every compact set  $C \subseteq X_c$ . The forward direction is again trivial.

For the backward direction take a set  $A \subseteq X_c$  such that for every compact set  $C \subseteq X_c$  the intersection  $A \cap C$  is open in  $C$ . By the definition of the k-ification it is enough to show that for every compact set  $C \subseteq X$  the intersection  $A \cap C$  is open in  $C$ . So let  $C \subseteq X$  be a compact set. By 4.1.6  $C$  is also compact in  $X$ . By assumption this means that  $A \cap C$  is open in  $C \subseteq X_c$  (in the subspace topology of the k-ification). Thus there is an open set  $B \subset X_c$  such that  $A \cap C = B \cap C$ . By the definition of the k-ification  $B \cap C$  is open in  $C \subseteq X$ . That means there is an open set  $E \subseteq X$  such that  $B \cap C = E \cap C$ . But that now gives us  $A \cap C = B \cap C = E \cap C$  with which we can conclude that  $A \cap C$  is open in  $C \subseteq X$  (in the subspace topology of the original topology of  $X$ ).  $\square$

If we already have a k-space, then the k-ification just maintains the topology of our space:

**Lemma 4.1.8.** Let  $X$  be a k-space. Then the topologies of  $X$  and  $X_c$  coincide.

*Proof.* Notice that the characterisation of open sets in  $X$  and  $X_c$  respectively agree in this setting.  $\square$



**Corollary 4.1.9.** *The k-ification is idempotent.*

Now we will characterise continuous maps to and from the k-ification. Going from the k-ification is not a big issue:

**Lemma 4.1.10.** *Let  $f: X \rightarrow Y$  be a continuous map of topological spaces. Then  $f: X_c \rightarrow Y$  is continuous.*

*Proof.* This follows easily from lemma 4.1.4. □

More interesting questions are when a map to the k-ification or a map from a k-ification to a k-ification is continuous. The following two lemmas and proofs that answer these questions are based on lemma 46.4 of [Mun14]. The next lemma is the first step towards the answer:

**Lemma 4.1.11.** *Let  $X$  be a compact space and  $f: X \rightarrow Y$  be a continuous map. Then  $f: X \rightarrow Y_c$  is continuous.*

*Proof.* We want to show that for every closed  $A \subseteq Y_c$  the preimage  $f^{-1}(A)$  is closed in  $X$ . Take any closed set  $A \subseteq Y_c$ . We know by lemma 4.1.5 that  $A \cap C$  is closed in  $C$  for every compact  $C \subseteq Y$ . As the image of a compact set  $f(X)$  is compact. Thus  $A \cap f(X)$  is closed in  $f(X) \subseteq Y$ . By the definition of the subspace topology there is a closed set  $B \subseteq Y$  such that  $A \cap f(X) = B \cap f(X)$ . Now we have

$$f^{-1}(A) = f^{-1}(A \cap f(X)) = f^{-1}(B \cap f(X)) = f^{-1}(B)$$

which is closed as the preimage of a closed set under a continuous map. □

Now this helps us get the following lemma:

**Lemma 4.1.12.** *Let  $f: X \rightarrow Y$  be a map of topological spaces such that for every compact  $C \subseteq X$  the restriction  $f|_C: C \rightarrow Y$  is continuous. Then  $f: X_c \rightarrow Y_c$  is continuous.*

*Proof.* The last lemma together with our assumption tells us that for every compact  $C \subseteq X$  the restriction  $f|_C: C \rightarrow Y_c$  is continuous. To show the claim take any open  $A \subseteq Y_c$ . We need to show that  $f^{-1}(A) \subseteq X_c$  is open. By definition of the k-ification this set is open if for all compact sets  $C \subseteq X$  the intersection  $f^{-1}(A) \cap C$  is open in  $C$ . Take any compact set  $C \subseteq X$ . As noted above we now know that  $f|_C: C \rightarrow Y_c$  is continuous. Or in other words we know that for every open  $B \subseteq Y_c$  there is an open set  $E \subseteq X$  such that  $f^{-1}(B) \cap C = E \cap C$ . Applying this to the set  $A \subseteq Y_c$  gives us an open set  $E \subseteq X$  such that  $f^{-1}(A) \cap C = E \cap C$ . But that is just another way of stating that  $f^{-1}(A)$  is open in  $C \subseteq X$ . □

That yields the following corollary:

**Corollary 4.1.13.** *Let  $f: X \rightarrow Y$  be a continuous map of topological spaces. Then  $f: X_c \rightarrow Y_c$  is continuous.*

*Proof.* This situation trivially fulfils the conditions of the previous lemma. □

If you look at the discussion of the product of CW-complexes in some topology books, for example [Hat01] and [Lüc05], you will notice that the k-ification rarely gets discussed in detail. One possible reason for this is that most common spaces that you encounter are already k-spaces. Lemma 4.1.8 then allows you to ignore the k-ification entirely. We will therefore discuss in the remainder of this section which spaces are k-spaces and which are not. The first example are weakly locally compact spaces.

**Definition 4.1.14.** Let  $X$  be a topological space. We call  $X$  *weakly locally compact* if every point  $x \in X$  has some compact neighbourhood.

This property is in some sources just called locally compact. The following proof is from Lemma 46.3 in [Mun14].

**Lemma 4.1.15.** *Weakly locally compact spaces are k-spaces.*

*Proof.* Let  $X$  be a weakly locally compact space. Let  $A \subseteq X$ . We need to show that  $A$  is open iff  $A \cap C$  is open in  $C$  for every compact set  $C$ . The forward direction is trivial. So assume that for every compact set  $C$  the intersection  $A \cap C$  is open in  $C$ .  $A$  is open if it is a neighbourhood of every point  $x \in A$ . So fix any  $x \in A$ . Since  $X$  is weakly locally compact,  $x$  has a compact neighbourhood  $C$ . By definition of neighbourhoods there is an open set  $U \subseteq C$  such that  $x \in U$  and we need to find an open set  $V \subseteq A$  such that  $x \in V$ . We show that  $A \cap U$  fulfils these conditions. It is obvious that  $A \cap U \subseteq A$  and  $x \in A \cap U$ . So it is left to show that  $A \cap U$  is open. By assumption  $A \cap C$  is open in  $C$ . That means that there is an open set  $B$  such that  $A \cap C = B \cap C$ . This now gives us

$$A \cap U = A \cap C \cap U = B \cap C \cap U = B \cap U$$

which is open as the intersection of two open sets. □

Another big class of spaces which are k-spaces are sequential spaces.

**Definition 4.1.16.** A set  $A$  in a topological space  $X$  is *sequentially closed* if for every convergent sequence contained in  $A$  its limit point is also in  $A$ . The *sequential closure* of a set  $A$  in  $X$  is defined as  $\text{scl}(A) = \{x \in X \mid \text{there is a sequence } (a_n)_{n \in \mathbb{N}} \subseteq A \text{ such that } (a_n)_{n \in \mathbb{N}} \text{ converges to } x\}$ . A *sequential space* is a space in which all sequentially closed sets are closed.

We will need the following characterisation of sequentially closed sets:

**Lemma 4.1.17.** *A set  $A \subseteq X$  is sequentially closed iff  $A = \text{scl}(A)$ .*

*Proof.* This is easy to see from the definitions. □

The following proof is based on [Sco16] and Lemma 46.3 in [Mun14].

**Lemma 4.1.18.** *Sequential Spaces are k-spaces.*

*Proof.* Let  $X$  be a Sequential Space. By lemma 4.1.2 it is enough to show that

$$A \subseteq X \text{ is closed} \iff \text{for all compact sets } C \subseteq X \text{ the intersection } A \cap C \text{ is closed in } C.$$

The forward direction is trivial. Let  $A$  be a set such that  $A \cap C$  is closed in  $C$  for every compact set  $C$ . Since  $X$  is a sequential space it is enough to show that  $A$  is sequentially closed or by the previous lemma  $A = \text{scl}(A)$ . The inclusion  $A \subseteq \text{scl}(A)$  is obvious. For the backward inclusion take  $x \in \text{scl}(A)$ . We need to show that  $x \in A$ . By definition there is a sequence  $(a_n)_{n \in \mathbb{N}} \subseteq A$  that converges to  $x$ . It is well known (and can be shown directly from the definition of compactness) that the set  $\{a_n \mid n \in \mathbb{N}\} \cup x$  is compact as the set of terms of a sequence together with the limit point of that sequence. By assumption that gives us that  $A \cap (\{a_n \mid n \in \mathbb{N}\} \cup x)$  is closed in  $\{a_n \mid n \in \mathbb{N}\} \cup x$ . In other words there is a closed set  $B$  such that

$$A \cap (\{a_n \mid n \in \mathbb{N}\} \cup x) = B \cap (\{a_n \mid n \in \mathbb{N}\} \cup x).$$

With that we get

$$x \in A \iff x \in A \cap (\{a_n \mid n \in \mathbb{N}\} \cup x) = B \cap (\{a_n \mid n \in \mathbb{N}\} \cup x) \iff x \in B$$

and for all  $n \in \mathbb{N}$  we get  $a_n \in B$  in the exact same way. Thus  $(a_n)_{n \in \mathbb{N}} \subseteq B$ . Since  $B$  is in particular sequentially closed this gives us  $x \in B$  which is enough by the above equivalence.  $\square$

In particular sequential spaces include metric spaces:

**Lemma 4.1.19.** *Metric spaces are sequential spaces.*

*Proof.* Let  $X$  be a metric space and  $A$  be a sequentially closed set. We need to show that that  $A$  is closed which is equivalent to  $A^c$  being open. Assume towards a contradiction that  $A^c$  is not open. Then there is a point  $x \in A^c$  such that for every  $n \in \mathbb{N}$  the open ball  $B_{1/n}(x)$  contains a point  $x_n \in A$ . But then we have a sequence  $(x_n)_{n \in \mathbb{N}} \subseteq A$  that converges to  $x \in A^c$ . Thus  $A$  is not sequentially closed. Contradiction.  $\square$

**Corollary 4.1.20.** *Metric spaces are k-spaces.*

Lastly we will discuss spaces that are not k-spaces:

**Lemma 4.1.21.** *Let  $X$  be an anti-compact  $T_1$  space. Then  $X_c$  has discrete topology.*

*Proof.* Let  $A \subseteq X_c$  be any set. We need to show that it is open. By the definition of the k-ification it is enough to show that  $A \cap C$  is open in  $C$  for every compact set  $C \subseteq X$ . Since  $X$  is anti-compact  $C$  is finite. And by  $T_1$  every finite set has discrete topology. Thus  $A \cap C$  is open in  $C$  and  $X_c$  has discrete topology.  $\square$

**Corollary 4.1.22.** *Let  $X$  be a non-discrete anti-compact  $T_1$  space. Then  $X$  is not a k-space.*

*Proof.* This follows easily from the previous lemma and lemma 4.1.8.  $\square$

That leads us to our first concrete example of a space that is not a k-space:

**Example 4.1.23.** Let  $X$  be any uncountable set. Equip  $X$  with the cocountable topology, i.e. let a set  $A \subseteq X$  be open iff  $A = \emptyset$  or  $A^c$  is countable. Then  $X$  is not a k-space.

*Proof.* It is easy to see by going through the axioms that the cocountable topology is indeed a topology. We will now show that this space satisfies the conditions of the previous corollary.  $X$  is clearly non-discrete. To see that  $X$  is a  $T_1$  space take two distinct points  $a$  and  $b$ . Now let  $A$  be the set  $X \setminus \{b\}$ . This set is open since  $\{b\}$  is countable and it obviously does not contain  $b$ . We lastly need to show that that  $X$  is anti-compact. To do that take any set  $A \subseteq X$ . Pick an (if possible infinite) countable subset  $B \subseteq A$ . Now for every  $b \in B$  define  $U_b = (X \setminus B) \cup \{b\}$ . Since  $U_b^c = B \setminus \{b\}$  is countable  $U_b$  is open for every  $b \in B$ . It is also easy to see that  $A \subseteq \bigcup_{b \in B} U_b$ . Thus  $(U_b)_{b \in B}$  is an open cover of  $A$ . But since for every  $b \in B$  there is no  $b' \in B$  with  $b \neq b'$  and  $b \in U_{b'}$ ,  $(U_b)_{b \in B}$  cannot have a proper subcover. Therefore  $A$  can only be compact if all these possible covers are already finite. That can only be the case if  $B$  and with that  $A$  are finite.  $\square$

Other examples can be found on  $\pi$ -base [PiB24].

## 4.2. The product of CW-complexes

We can now move on to discuss the product. We want to prove the following theorem:

**Theorem 4.2.1.**

**Lemma 4.2.2.**  $(X \times Y)_c$  has weak topology, i.e.  $A \subseteq (X \times Y)_c$  is closed iff  $(Q_i^n \times P_j^m)(D^{n+m}) \cap A$  is closed for all  $n, m \in \mathbb{N}$ ,  $i \in I_n$  and  $j \in J_m$ .

*Proof.*

" $\Rightarrow$ " Since  $D^{n+m}$  is compact, its image is compact and therefore closed. As the intersection of two closed sets  $(Q_i^n \times P_j^m)(D^{n+m}) \cap A$  is closed as well.

" $\Leftarrow$ " We know by definition of the k-ification that  $A$  is closed if for every compact set  $C \subseteq X \times Y$   $A \cap C$  is closed in  $C$ . Take such a compact set  $C$ . The projections  $\text{pr}_1(C)$  and  $\text{pr}_2(C)$  are compact as images of a compact set. By ? there are finite sets  $E \subseteq \{e_i^n \mid n \in \mathbb{N}, i \in I_n\}$  and  $F \subseteq \{f_j^m \mid m \in \mathbb{N}, j \in J_m\}$  s.t  $\text{pr}_1(C) \subseteq \bigcup_{e \in E} e$  and  $\text{pr}_2(C) \subseteq \bigcup_{f \in F} f$ . Thus

$$C \subseteq \text{pr}_1(C) \times \text{pr}_2(C) \subseteq \bigcup_{e \in E} e \times \bigcup_{f \in F} f = \bigcup_{e \in E} \bigcup_{f \in F} e \times f.$$

So  $C$  is included in a finite union of cells of  $(X \times Y)_c$ . Therefore

$$A \cap C = A \cap \left( \bigcup_{e \in E} \bigcup_{f \in F} e \times f \right) \cap C = \left( \bigcup_{e \in E} \bigcup_{f \in F} A \cap (e \times f) \right) \cap C$$

is closed since by assumption  $A \cap (e \times f)$  is closed for every  $e$  and  $f$  and the union is finite. Thus  $A \cap C$  is in particular closed in  $C$ .  $\square$

# Appendix

A. a

B. b



# Symbol Index

$D^n$	The closed unit disk in $\mathbb{R}^n$ , i.e. $D^n := \{x \in \mathbb{R}^n \mid \ x\  \leq 1\}$ .
$S^n$	The boundary of the unit disk in $\mathbb{R}^n$ , i.e. $S^n := \{x \in \mathbb{R}^n \mid \ x\  = 1\}$ .
$\bar{e}^n$	A closed $n$ -cell, i.e. $\bar{e}^n := Q^n(D^n)$ . See definition 3.1.2.
$\partial e^n$	The edge of an $n$ -cell, i.e. $\partial e^n := Q^n(\partial D^n)$ . See definition 3.1.2.
$e^n$	An (open) $n$ -cell, i.e. $e^n := Q^n(\text{int}(D^n))$ . See definition 3.1.2.
$\mathbb{R}$	The real numbers
$F$	A topological or Riemann surface.





# Index

vectorspace, 7



# Bibliography

- [Abh05] Jochen Abhau. “Die Homologie von Modulräumen Riemannscher Flächen – Berechnungen für  $g \leq 2$ ”. Diplomarbeit. Rheinische Friedrich-Wilhelms-Universität Bonn, 2005.
- [Dis06] Margherita Disertori. “Constructive Renormalization for Interacting Fermions”. In: *Lett Math Phys* 78 (2006), pp. 263–277. DOI: 10.1007/s11005-006-0124-0. URL: <https://doi.org/10.1007/s11005-006-0124-0>.
- [Hat01] Allen Hatcher. *Algebraic topology*. 16. printing 2016. Cambridge: Cambridge Univ. Press, 2001. ISBN: 9780521795401.
- [Lüc05] Wolfgang Lück. *Algebraische Topologie Homologie und Mannigfaltigkeiten*. ger. 1. Aufl. Vieweg Studium : Aufbaukurs Mathematik. Wiesbaden: Vieweg, 2005. ISBN: 3528032189.
- [Mun14] James Raymond Munkres. *Topology*. 2. ed., Pearson new internat. ed. Harlow: Pearson, 2014. ISBN: 9781292023625.
- [PiB24] The pi-Base Community. *Search for ‘Generated by compact subspaces’*.  $\pi$ -base. Accessed: 25.07.2024. 2024. URL: <https://topology.pi-base.org/spaces?q=%7EGenerated+by+compact+subspaces>.
- [Sco16] Brian M. Scott (<https://math.stackexchange.com/users/12042/brian-m-scott>). *A first countable Hausdorff space is compactly generated*. Mathematics Stack Exchange. Version: 24.11.2016. 2016. URL: <https://math.stackexchange.com/q/2026250>.
- [Whi18] J. H. C. Whitehead. “Combinatorial homotopy. I”. In: *Bulletin (new series) of the American Mathematical Society* 55.3 (2018), pp. 213–245. ISSN: 0273-0979.
- [Wik24] Wikipedia contributors. *Compactly generated space*. Wikipedia, The Free Encyclopedia. Accessed: 23.07.2024. 2024. URL: [https://en.wikipedia.org/w/index.php?title=Compactly\\_generated\\_space&oldid=1233649259](https://en.wikipedia.org/w/index.php?title=Compactly_generated_space&oldid=1233649259).