

STA219 Assignment 6

12312706 Zhou Liangyu

1. Standard normal distribution is symmetric, i.e. the probability only depends on the distance from the origin. Therefore, we can consider using the polar coordinate system to describe it.

Let $R = \sqrt{-2 \ln U_1}$, $\theta = 2\pi U_2$, we have $\theta \sim U(0, 2\pi)$, $f_\theta(\omega) = \frac{1}{2\pi}$, and $Z_1 = R \cos \theta$, $Z_2 = R \sin \theta$.

$$\therefore P(R^2 \leq t) = P(-2 \ln U_1 \leq t) = P(U_1 \geq e^{-\frac{t}{2}}) = 1 - e^{-\frac{t}{2}}$$

$$\therefore R^2 \sim \text{Exp}\left(\frac{1}{2}\right).$$

$$\therefore F_R(r) = F_{R^2}(r^2) = 1 - e^{-\frac{r^2}{2}}, f_R(r) = \frac{dF_R(r)}{dr} = re^{-\frac{r^2}{2}}.$$

$$\therefore R \text{ and } \theta \text{ are independent, } f_{R,\theta}(r, \omega) = f_R(r)f_\theta(\omega) = \frac{re^{-\frac{r^2}{2}}}{2\pi}.$$

$$\therefore Z_1 = R \cos \theta, Z_2 = R \sin \theta$$

$$\therefore R = \sqrt{Z_1^2 + Z_2^2}, f_{Z_1, Z_2}(z_1, z_2) = \frac{1}{r} f_{R,\theta}(r, \omega) = \frac{e^{-\frac{z_1^2 + z_2^2}{2}}}{2\pi} = \frac{1}{\sqrt{2\pi}} e^{-\frac{z_1^2}{2}} \cdot \frac{1}{\sqrt{2\pi}} e^{-\frac{z_2^2}{2}}.$$

$$\therefore Z_1 \text{ and } Z_2 \text{ are independent, and } f_{Z_1}(z_1) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z_1^2}{2}}, f_{Z_2}(z_2) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z_2^2}{2}}.$$

$$\therefore P(Z_1 \leq a, Z_2 \leq b) = \int_{-\infty}^b \int_{-\infty}^a f_{Z_1, Z_2}(z_1, z_2) dz_1 dz_2 = \int_{-\infty}^a f_{Z_1}(z_1) dz_1 \int_{-\infty}^b f_{Z_2}(z_2) dz_2 = \Phi(a)\Phi(b).$$

$\therefore Z_1$ and Z_2 are a pair of independent standard normal random variables.

2. (1) Sample mean: $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i = \frac{1}{10}(17.2 + 22.1 + 18.5 + 17.2 + 18.6 + 14.8 + 21.7 + 15.8 + 16.3 + 22.8) = 18.5$.

$$\begin{aligned} \text{Sample variance: } S^2 &= \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 = \frac{1}{9} [(17.2 - 18.5)^2 + (22.1 - 18.5)^2 + (18.5 - 18.5)^2 + (17.2 - 18.5)^2 + \\ &\quad (18.6 - 18.5)^2 + (14.8 - 18.5)^2 + (21.7 - 18.5)^2 + (15.8 - 18.5)^2 + (16.3 - 18.5)^2 + (22.8 - 18.5)^2] \\ &\approx 7.88. \end{aligned}$$

$$\text{Sample standard deviation: } S = \sqrt{S^2} \approx 2.81.$$

(2) Sample lower quartile: $Q_{0.25} = X_{(\lfloor 10 \times 0.25 + 1 \rfloor)} = X_{(3)} = 16.3$.

Sample upper quartile: $Q_{0.75} = X_{(\lfloor 10 \times 0.75 + 1 \rfloor)} = X_{(8)} = 21.7$.

Sample interquartile range: $Q_{0.75} - Q_{0.25} = 21.7 - 16.3 = 5.4$.

3. (1) $\therefore X \sim U(0, \theta)$

$$\therefore E(X) = \frac{\theta}{2}, \text{Var}(X) = \frac{\theta^2}{12}, f_X(x) = \frac{1}{\theta}, F_X(x) = \frac{x}{\theta}.$$

$$\therefore f_{\min}(x) = nf(x)[1 - F(x)]^{n-1} = \frac{3}{\theta} \left(1 - \frac{x}{\theta}\right)^2 = \frac{2(\theta - x)^2}{\theta^3}, f_{\max}(x) = nf(x)F(x)^2 = \frac{3}{\theta} \left(\frac{x}{\theta}\right)^2 = \frac{3x^2}{\theta^3}.$$

$$\therefore E(X_{(1)}) = \int_0^\theta x f_{\min}(x) dx = \int_0^\theta \frac{3x(\theta - x)^2}{\theta^3} dx = -\frac{1}{\theta^3} \int_0^\theta x d(\theta - x)^3 = -\frac{1}{\theta^3} [0 - \int_0^\theta (\theta - x)^3 dx] = \frac{\theta}{4},$$

$$E(X_{(3)}) = \int_0^\theta x f_{\max}(x) dx = \int_0^\theta \frac{3x^3}{\theta^3} dx = \frac{3x^4}{4\theta^3} \Big|_0^\theta = \frac{3\theta}{4}.$$

$$\therefore E(\hat{\theta}_1) = \frac{4}{3} E(X_{(3)}) = \theta, E(\hat{\theta}_2) = 4E(X_{(1)}) = \theta.$$

$\therefore \hat{\theta}_1$ and $\hat{\theta}_2$ are both unbiased estimators of θ .

$$(2) \therefore E(X_{(1)}^2) = \int_0^\theta x^2 f_{\min}(x) dx = \int_0^\theta \frac{3x^2(\theta-x)^2}{\theta^3} dx = \frac{1}{\theta^3} \int_0^\theta (\theta-x)^2 dx^3 = \frac{1}{\theta^3} [0 - \int_0^\theta x^3 d(\theta-x)^2] = \frac{\theta^2}{10},$$

$$E(X_{(3)}^2) = \int_0^\theta x^2 f_{\max}(x) dx = \int_0^\theta \frac{3x^4}{\theta^3} dx = \frac{3x^5}{5\theta^3} \Big|_0^\theta = \frac{3\theta^2}{5}.$$

$$\therefore \text{Var}(X_{(1)}) = E(X_{(1)}^2) - E(X_{(1)})^2 = \frac{\theta^2}{10} - \frac{\theta^2}{16} = \frac{3\theta^2}{80}, \text{Var}(X_{(3)}) = E(X_{(3)}^2) - E(X_{(3)})^2 = \frac{3\theta^2}{5} - \frac{9\theta^2}{16} = \frac{3\theta^2}{80}.$$

$$\therefore \text{Var}(\hat{\theta}_1) = \frac{16}{9} \text{Var}(X_{(3)}) = \frac{\theta^2}{15}, \text{Var}(\hat{\theta}_2) = 16 \text{Var}(X_{(1)}) = \frac{3\theta^2}{5}.$$

$$\therefore \text{Var}(\hat{\theta}_1) < \text{Var}(\hat{\theta}_2)$$

$$\therefore \hat{\theta}_1 \text{ is more efficient than } \hat{\theta}_2.$$

4. Python Code:

```

1  import numpy as np
2  import matplotlib.pyplot as plt
3  from scipy.stats import geom, cauchy
4
5  # Set seed
6  np.random.seed(20250510)
7
8  # Generate uniform samples
9  n_samples = 10000
10 unif_samples1 = np.random.uniform(0, 1, n_samples)
11 unif_samples2 = np.random.uniform(0, 1, n_samples)
12
13 # Transform
14 p_geom = 0.5
15 geo_samples = np.ceil(np.log(1 - unif_samples1) / np.log(1 - p_geom))
16 stdcauchy_samples = np.tan(np.pi * (unif_samples2 - 0.5))
17
18 # Theoretical PMF/PDF
19 geo_x = np.arange(min(geo_samples), max(geo_samples))
20 geo_pmf = geom.pmf(geo_x, p_geom)
21 stdcauchy_x = np.linspace(-10, 10, 1000)
22 stdcauchy_pdf = cauchy.pdf(stdcauchy_x)
23
24 # Adjust the plots
25 plt.figure(figsize=(19.5, 9), dpi=143.4)
26 plt.rcParams['font.size'] = 12
27 plt.tight_layout()
28
29 # Plot - Geometric(0.5)
30 plt.subplot(1, 2, 1)
31 plt.hist(geo_samples, bins=np.arange(1, 17)-0.5, density=True, alpha=0.6, color='skyblue',
32          label='Generated Geometric')
33 plt.scatter(geo_x, geo_pmf, color='red', s=20, marker='o', label='Theoretical PMF', zorder=2)
34 plt.xlabel('x')
35 plt.ylabel('Probability')
36 plt.title('Geometric(0.5)')
37 plt.legend()
38
39 # Plot - Standard Cauchy Distribution
40 plt.subplot(1, 2, 2)
41 plt.hist(stdcauchy_samples, bins=100, density=True, range=(-10, 10), alpha=0.6, color='skyblue',
42          label='Generated Standard Cauchy')
43 plt.plot(stdcauchy_x, stdcauchy_pdf, 'r-', label='Theoretical PDF')
44 plt.xlabel('x')


```

```

43 plt.ylabel('Density')
44 plt.title('Standard Cauchy Distribution')
45 plt.legend()
46
47 # Save the plot for assignment
48 plt.savefig('Geometric_StdCauchy_Generate.png')
49
50 # Show
51 plt.show()

```

Plots:



5. (1) (2) (3) Proposal distribution: $N(-3, 16)$. Acceptance proportion: 28.64%.

See the following Python code and results:

```


1  import numpy as np
2  import matplotlib.pyplot as plt
3  from scipy.stats import norm
4  from scipy.integrate import quad
5  from scipy.optimize import minimize_scalar
6
7  # Reject sampling
8  def reject_sampling(n_samples, mu, sigma, c):
9      samples = []
10     accepted = 0
11     total_trials = 0
12
13     while accepted < n_samples:
14         batch_size = 100000
15         x_prop = np.random.normal(loc=mu, scale=sigma, size=batch_size)
16         u = np.random.rand(batch_size)
17
18         g_x = norm.pdf(x_prop, loc=mu, scale=sigma)
19         accept_prob = fs(x_prop) / (c * g_x)
20
21         accept_mask = u <= accept_prob
22         accepted_samples = x_prop[accept_mask]
23
24         samples.extend(accepted_samples)

```

```

25         accepted += len(accepted_samples)
26         total_trials += batch_size
27
28     accept_rate = accepted / total_trials
29     return np.array(samples[:n_samples]), accept_rate
30
31 fs = lambda x: 0.6 * np.exp(-((x + 5) ** 2) / 2) + 0.4 * np.exp(-((x - 1) ** 2) / 0.5)
32 c1, _ = quad(fs, -np.inf, np.inf)
33 mu = -3
34 sigma = 4
35 g = lambda x: norm.pdf(x, loc=mu, scale=sigma)
36 c = 7 # by trial and error
37
38 n_samples = 500000
39 samples, accept_rate = reject_sampling(n_samples, mu, sigma, c)
40 x = np.linspace(-15, 10, 1000)
41
42 # Set plot parameters
43 plt.figure(figsize=(14.3, 10), dpi=143.4)
44 plt.rcParams['font.size'] = 16
45 plt.tight_layout()
46
47 # Plot to show that it covers the target distribution, the histogram of the generated samples, and
48 # compare it with the theoretical PDF
49 plt.plot(x, fs(x), label='Target f*(x)')
50 plt.plot(x, c*g(x), label=f'Proposal cg(x) (c={c})')
51 plt.hist(samples, bins=200, density=True, alpha=0.6, label='Generated Samples')
52 plt.plot(x, fs(x)/c1, 'r-', label='Normalized f*(x)', linewidth=2)
53 plt.xlabel('x')
54 plt.ylabel('Density')
55 plt.legend()
56 plt.title("Target and Proposal Distribution, Sampled Histogram with Theoretical PDF")
57 plt.grid(True)
58 plt.savefig('RejectSampling.png')
59 plt.show()
60 print(f"Acceptance Rate = {accept_rate:.2%}")

```



Code running result:

```

PS C:\Users\Chaos Zhou\PyCharmProjects\STA219> python -u 'c:\Users\Chaos Zhou\PyCharmProjects\STA219\Assignment6_P5.py'
Acceptance Rate = 28.65%
PS C:\Users\Chaos Zhou\PyCharmProjects\STA219> python -u 'c:\Users\Chaos Zhou\PyCharmProjects\STA219\Assignment6_P5.py'
Acceptance Rate = 28.62%
PS C:\Users\Chaos Zhou\PyCharmProjects\STA219> python -u 'c:\Users\Chaos Zhou\PyCharmProjects\STA219\Assignment6_P5.py'
Acceptance Rate = 28.64%
PS C:\Users\Chaos Zhou\PyCharmProjects\STA219> python -u 'c:\Users\Chaos Zhou\PyCharmProjects\STA219\Assignment6_P5.py'
Acceptance Rate = 28.64%
PS C:\Users\Chaos Zhou\PyCharmProjects\STA219> python -u 'c:\Users\Chaos Zhou\PyCharmProjects\STA219\Assignment6_P5.py'
Acceptance Rate = 28.64%
PS C:\Users\Chaos Zhou\PyCharmProjects\STA219> python -u 'c:\Users\Chaos Zhou\PyCharmProjects\STA219\Assignment6_P5.py'
Acceptance Rate = 28.60%
PS C:\Users\Chaos Zhou\PyCharmProjects\STA219> 

```

行 16, 列 39 空格: 4

6. (1) Let \hat{p}_n be the frequency of that more than 30% of the forest will eventually be burning.

By the CLT, we have $\hat{p}_n \overset{\text{approx.}}{\sim} N(p, \frac{p(1-p)}{n})$.

$$\therefore P(|\hat{p}_n - p| \leq 0.005) \approx 2\Phi\left(\frac{0.005\sqrt{n}}{\sqrt{p(1-p)}}\right) - 1 \geq 0.95 \Rightarrow \frac{0.005\sqrt{n}}{\sqrt{p(1-p)}} \geq \Phi^{-1}(0.975) \approx 1.96.$$

$$\therefore n \geq \frac{1.96^2 p(1-p)}{0.005^2} \geq \frac{1.96^2 \cdot 0.25}{0.005^2} = 38416.$$

Python code:

```

1 import random as rd
2 from collections import deque
3 import numpy as np
4
5 def forest_mc():
6     rows, cols = 20, 50
7     forest = [[False] * cols for _ in range(rows)]

```

```

8     forest[0][0] = True
9     q = deque([(0, 0)])
10    burning_tree_count = 1
11    while q:
12        i, j = q.popleft()
13        # right
14        if j + 1 < cols and not forest[i][j + 1]:
15            if rd.random() < 0.8:
16                forest[i][j + 1] = True
17                burning_tree_count += 1
18                q.append((i, j + 1))
19        # down
20        if i + 1 < rows and not forest[i + 1][j]:
21            if rd.random() < 0.3:
22                forest[i + 1][j] = True
23                burning_tree_count += 1
24                q.append((i + 1, j))
25    return burning_tree_count
26
27    n = 38416
28    samples = []
29    count = 0
30    for _ in range(n):
31        x = forest_mc()
32        samples.append(x)
33        if x > 300:
34            count += 1
35    prob = count / n
36    mean_x = np.mean(samples)
37    std_x = np.std(samples, ddof=1)
38
39    print(f"(1) The probability of X > 30%: {prob:.4f}")
40    print(f"(2) Mean of X: {mean_x:.2f}")
41    print(f"(3) Standard deviation of X: {std_x:.2f}")

```

The probability that more than 30% of the forest will eventually be burning: 0.0035.

- (2) The total number of affected trees X : 43.
- (3) The corresponding standard deviation of X : 62.99.

Code running result:

```

PS C:\Users\Chaos Zhou\PyCharmProjects\STA219> python -u "c:\Users\Chaos Zhou\PyCharmProjects\STA219\A.py"
(1) The probability of X > 30%: 0.0035
(2) Mean of X: 43
(3) Standard deviation of X: 62.89
PS C:\Users\Chaos Zhou\PyCharmProjects\STA219> python -u "c:\Users\Chaos Zhou\PyCharmProjects\STA219\A.py"
(1) The probability of X > 30%: 0.0036
(2) Mean of X: 44
(3) Standard deviation of X: 63.24
PS C:\Users\Chaos Zhou\PyCharmProjects\STA219> python -u "c:\Users\Chaos Zhou\PyCharmProjects\STA219\A.py"
(1) The probability of X > 30%: 0.0037
(2) Mean of X: 43
(3) Standard deviation of X: 63.11
PS C:\Users\Chaos Zhou\PyCharmProjects\STA219> python -u "c:\Users\Chaos Zhou\PyCharmProjects\STA219\A.py"
(1) The probability of X > 30%: 0.0034
(2) Mean of X: 43
(3) Standard deviation of X: 62.70
PS C:\Users\Chaos Zhou\PyCharmProjects\STA219>

```