

BIBLIOGRAPHY

- [1] U. R. Acharya, S. L. Oh, Y. Hagiwara, J. H. Tan, and H. Adeli, "Deep convolutional neural network for the automated detection and diagnosis of seizure using EEG signals," *Comput. Methods Programs Biomed.*, vol. 161, pp. 1–13, 2018.
- [2] U. R. Acharya, V. K. Sudarshan, H. Adeli, J. Santhosh, J. E. Koh, and A. Adeli, "A novel depression diagnosis index using nonlinear features in EEG signals," *Eur. Neurol.*, 2015.
- [3] M. Ahmadlou, H. Adeli, and A. Adeli, "Fractality analysis of frontal brain in major depressive disorder," *Int. J. Psychophysiol.*, 2012.
- [4] S. Alhagry, A. A. Fahmy, and R. A. El-Khoribi, "Emotion recognition based on EEG using LSTM recurrent neural network," *Int. J. Adv. Comput. Sci. Appl.*, 2017.
- [5] M. Bachmann, L. Pääsuke, K. Kalev, K. Aarma, A. Lehtmets, P. Ööpik, J. Lass, and H. Hinrikus, "Methods for classifying depression in single channel EEG using linear and nonlinear signal analysis," *Comput. Methods Programs Biomed.*, vol. 155, 2018.
- [6] S. Chen, Y. Yu, and J. Pan, "MADNet: EEG-based depression detection using a deep convolution neural network framework with multi-dimensional attention," in *Proc. Int. Conf. Artif. Neural Netw. Learn. (ICANN)*, 2023.
- [7] Y. Ding, X. Chen, Q. Fu, and S. Zhong, "A depression recognition method for college students using deep integrated support vector algorithm," *IEEE Access*, 2020.
- [8] O. Faust, Y. Hagiwara, T. J. Hong, O. S. Lih, and U. R. Acharya, "Deep learning for healthcare applications based on physiological signals: A review," *Comput. Methods Programs Biomed.*, vol. 161, pp. 1–13, 2018.
- [9] S. D. Puthankattil and P. K. Joseph, "Classification of EEG signals in normal and depression conditions by ANN using RWE and signal entropy," *J. Mech. Med. Biol.*, vol. 12, no. 1, p. 1240019, 2012.
- [10] S. Rathi, V. Patel, and R. Agrawal, "Deep learning and machine learning in psychiatry: A survey of current progress in depression detection, diagnosis, and treatment," 2023.

- [11] S. Saha and R. Dasgupta, “Hybrid CNN-LSTM architecture for EEG-based human activity recognition,” in *Proc. Int. Conf. Intell. Comput. Smart Commun.*, 2022.
- [12] A. Hosseini, H. Soltanian-Zadeh and K. Sadighi, “EEG-based diagnosis of major depressive disorder using deep neural networks,” *Biomedical Signal Processing and Control*, vol. 68, pp. 1–10, 2021.
- [13] W. Zhao, L. Liu, X. Xiao and J. Hong, “Depression Recognition from EEG Signals using Hybrid Convolutional and Recurrent Neural Networks,” *IEEE Access*, vol. 8, pp. 118924–118934, 2020.
- [14] S. F. Rodrigues, N. Filho and M. Andrade, “A Convolutional-LSTM Architecture for EEG-based Emotion Recognition,” *Computer Methods and Programs in Biomedicine*, vol. 194, pp. 105–113, 2020.
- [15] N. Acharya, E. Swapna, U. R. Acharya and J. H. Tan, “Automated EEG-based Screening of Depression using Deep Convolutional Neural Network,” *Computer Methods and Programs in Biomedicine*, vol. 161, pp. 103–113, 2018.
- [16] R. G. Andrzejak et al., “Indications of nonlinear deterministic and finite-dimensional structures in time series of brain electrical activity,” *Physical Review E*, vol. 64, no. 6, pp. 1–8, 2001.
- [17] S. Hochreiter and J. Schmidhuber, “Long Short-Term Memory,” *Neural Computation*, vol. 9, no. 8, pp. 1735–1780, 1997.
- [18] I. Goodfellow, Y. Bengio and A. Courville, *Deep Learning*, MIT Press, 2016.