

Pingala Assignment

Mannem Charan AI21BTECH11019

CONTENTS

1.2

Abstract—This manual provides a simple introduction to Transforms

1 JEE 2019

Let

$$a_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad n \geq 1 \quad (1.1)$$

$$b_n = a_{n-1} + a_{n+1}, \quad n \geq 2, \quad b_1 = 1 \quad (1.2)$$

Verify the following using a python code.

1.1

$$\sum_{k=1}^n a_k = a_{n+2} - 1, \quad n \geq 1 \quad (1.3)$$

Solution: Download the following python code from the below link,

```
wget https://github.com/Charanyash/EE3900-Digital_Signal_Processing/tree/master/pingala/Codes/1.py
```

Then run the following command,

```
python3 1.py
```




Fig. 1.1

$$\sum_{k=1}^{\infty} \frac{a_k}{10^k} = \frac{10}{89} \quad (1.4)$$

Solution:




Fig. 1.2

1.3

$$b_n = \alpha^n + \beta^n, \quad n \geq 1 \quad (1.5)$$

Solution:




Fig. 1.3

1.4

$$\sum_{k=1}^{\infty} \frac{b_k}{10^k} = \frac{8}{89} \quad (1.6)$$

Solution: As you can see in the 1.4, the summation is converging to $\frac{12}{89}$ but not $\frac{8}{89}$.




Fig. 1.4

2 PINGALA SERIES

2.1 The *one sided* Z-transform of $x(n)$ is defined as

$$X^+(z) = \sum_{n=0}^{\infty} x(n)z^{-n}, \quad z \in \mathbb{C} \quad (2.1)$$

2.2 The *Pingala* series is generated using the difference equation

$$x(n+2) = x(n+1) + x(n), \quad x(0) = x(1) = 1, n \geq 0 \quad (2.2)$$

Generate a stem plot for $x(n)$.

Solution: Download the below python code,

```
wget https://github.com/Charanyash/EE3900-Digital_Signal_Processing/tree/master/pingala/Codes/2.py
```

Then run the following command,

```
python3 2.py
```

2.3 Find $X^+(z)$.

Solution: Consider the eq :(2.2),

$$x(n+2) = x(n+1) + x(n) \quad (2.3)$$




Fig. 2.2

Now apply one-sided Z-transform on both sides,(using linearity)

$$\sum_{n=0}^{\infty} x(n+2)z^{-n} = \sum_{n=0}^{\infty} x(n+1)z^{-n} + \sum_{n=0}^{\infty} x(n)z^{-n} \quad (2.4)$$

$$z^2 (X^+(z) - x(1)z^{-1} - x(0)) = z(X^+(z) - x(0)) + X^+(z) \quad (2.5)$$

$$X^+(z) = \frac{x(1)z + x(0)(z^2 - z)}{z^2 - z - 1} \quad (2.6)$$

$$\Rightarrow X^+(z) = \frac{1}{1 - z^{-1} - z^{-2}} \quad (|z| \neq 0) \quad (2.7)$$

2.4 Find $x(n)$.

Solution: We know that,

$$X^+(z) = \frac{1}{1 - z^{-1} - z^{-2}} \quad (2.8)$$

Using partial fractions,

$$= \frac{1}{(1 - \alpha z^{-1})(1 - \beta z^{-1})} \quad (2.9)$$

$$= \frac{1}{\alpha - \beta} \left(\frac{\alpha}{1 - \alpha z^{-1}} - \frac{\beta}{1 - \beta z^{-1}} \right) \quad (2.10)$$

where α, β are the roots of the equation,

$$z^2 - z - 1 = 0 \quad (2.11)$$

Using the result of Z-transform,

$$a^n u(n) \xleftrightarrow{Z} \frac{1}{1 - az^{-1}} \quad |z| > |a| \quad (2.12)$$

$$(2.13)$$

We can write,

$$x(n) = \frac{\alpha^{n+1} - \beta^{n+1}}{\alpha - \beta} \quad (2.14)$$

with ROC as,

$$|z| > \max \{\alpha, \beta\} \quad (2.15)$$

2.5 Sketch

$$y(n) = x(n-1) + x(n+1), \quad n \geq 0 \quad (2.16)$$

Solution: Download the below python code,

```
wget https://github.com/Charanyash/EE3900-Digital_Signal_Processing/tree/master/pingala/Codes/2.py
```

Then run the following command,

```
python3 2.py
```




Fig. 2.5

2.6 Find $Y^+(z)$.

Solution: Apply the one-sided Z-transform on (??),

$$\sum_{n=0}^{\infty} y(n)z^{-n} = \sum_{n=0}^{\infty} x(n-1)z^{-n} + \sum_{n=0}^{\infty} x(n+1)z^{-n} \quad (2.17)$$

$$= z^{-1} (X^+(z) - x(-1)) + z (X^+(z) - x(0)) \quad (2.18)$$

Using (2.2) and (2.7),

$$Y^+(z) = \frac{z^{-1}}{1 - z^{-1} - z^{-2}} + \frac{1 + z^{-1}}{1 - z^{-1} - z^{-2}} \quad (2.19)$$

$$\Rightarrow Y^+ z = \frac{1 + 2z^{-1}}{1 - z^{-1} - z^{-2}} \quad (2.20)$$

2.7 Find $y(n)$.

Solution: Consider (??),

$$Y^+ z = \frac{1 + 2z^{-1}}{1 - z^{-1} - z^{-2}} \quad (2.21)$$

$$= X^+(z) + 2z^{-1}X^+(z) \quad (2.22)$$

Using (??) and the property of z-transform that,

$$x(n-1) \xleftrightarrow{Z} z^{-1}X^+(z) \quad (2.23)$$

we can write the inverse z-transform of $Y^+(z)$ (using ROC as $|z| > \max \{\alpha, \beta\}$) as,

$$y(n) = x(n) + 2x(n-1) \quad (2.24)$$

$$= \frac{\alpha^{n+1} - \beta^{n+1}}{\alpha - \beta} + 2 \frac{\alpha^n - \beta^n}{\alpha - \beta} \quad (2.25)$$

$$= \frac{\alpha^{n+2} + \alpha^n - \beta^{n+2} - \beta^n}{\alpha - \beta} \quad (2.26)$$

$$= \frac{\alpha^{n+2} - \beta^{n+2} - \beta\alpha^{n+1} + \beta^{n+1}\alpha}{\alpha - \beta} \quad (\because \alpha\beta = -1) \quad (2.27)$$

$$\Rightarrow y(n) = \alpha^{n+1} + \beta^{n+1} \quad (2.28)$$

3 POWER OF THE Z TRANSFORM

3.1 Show that

$$\sum_{k=1}^n a_k = \sum_{k=0}^{n-1} x(n) = x(n) * u(n-1) \quad (3.1)$$

3.2 Show that

$$a_{n+2} - 1, \quad n \geq 1 \quad (3.2)$$

can be expressed as

$$[x(n+1) - 1]u(n) \quad (3.3)$$

3.3 Show that

$$\sum_{k=1}^{\infty} \frac{a_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{x(k)}{10^k} = \frac{1}{10} X^+(10) \quad (3.4)$$

3.4 Show that

$$\alpha^n + \beta^n, \quad n \geq 1 \quad (3.5)$$

can be expressed as

$$w(n) = (\alpha^{n+1} + \beta^{n+1})u(n) \quad (3.6)$$

and find $W(z)$.

3.5 Show that

$$\sum_{k=1}^{\infty} \frac{b_k}{10^k} = \frac{1}{10} \sum_{k=0}^{\infty} \frac{y(k)}{10^k} = \frac{1}{10} Y^+(10) \quad (3.7)$$

3.6 Solve the JEE 2019 problem.