

Chapter 06 Part 1: Ideal Gas Law

1. A 2 L container will hold about 4 g of which of the following gases at 0°C and 1 atm?


(A) SO_2

(B) N_2

(C) CO_2


(D) C_4H_8

(E) NH_3



At 27°C, five identical rigid 2.0 L vessels are filled with $\text{N}_2(g)$ and sealed. Four of the five vessels also contain a 0.050 mol sample of $\text{NaHCO}_3(s)$, $\text{NaBr}(s)$, $\text{Cu}(s)$, or $\text{I}_2(s)$, as shown in the diagram above. The volume taken up by the solids is negligible, and the initial pressure of $\text{N}_2(g)$ in each vessel is 720 mm Hg. All four vessels are heated to 127°C and allowed to reach a constant pressure.


2.




The gas particles in vessel 3 at 27°C are represented in the diagram above. The lengths of the arrows represent the speeds of the particles. Which of the following diagrams best represents the particles when vessel 3 is heated to 127°C?

Chapter 06 Part 1: Ideal Gas Law


(A)




(B)



(C)



(D)



3. At 127°C , the pressure in vessel 1 is found to be higher than that in vessel 2. Which of the following reactions best accounts for the observation?


Chapter 06 Part 1: Ideal Gas Law

- (A) $\text{NaHCO}_3(s) \rightarrow \text{Na}(s) + \text{HCO}_3(s)$
- (B) $\text{NaHCO}_3(s) \rightarrow \text{NaH}(s) + \text{CO}_3(s)$
- (C) $2 \text{NaHCO}_3(s) \rightarrow \text{Na}_2\text{CO}_3(s) + \text{CO}_2(g) + \text{H}_2\text{O}(g)$ ✓
- (D) $2 \text{NaHCO}_3(s) + \text{N}_2(g) \rightarrow 2 \text{NaNO}_3(s) + \text{C}_2\text{H}_2(g)$

4. At standard temperature and pressure, a 0.50 mol sample of H_2 gas and a separate 1.0 mol sample of O_2 gas have the same

- (A) average molecular kinetic energy ✓
- (B) average molecular speed
- (C) volume
- (D) effusion rate
- (E) density






5.



The distribution of speeds of $\text{H}_2(g)$ molecules at 273 K and 1 atm is shown in the diagram above. Which of the following best shows the speed distribution of $\text{He}(g)$ atoms under the same conditions of temperature and pressure?





———— $\text{H}_2(g)$ - - - - - $\text{He}(g)$

Chapter 06 Part 1: Ideal Gas Law




The graph above shows the speed distribution of molecules in a sample of a gas at a certain temperature. Which of the following graphs shows the speed distribution of the same molecules at a lower temperature (as a dashed curve) ?

Chapter 06 Part 1: Ideal Gas Law



7.



The Maxwell-Boltzmann distributions of molecular speeds in samples of two different gases at the same temperature are shown above. Which gas has the greater molar mass?

(A) Gas A



(B) Gas B

(C) Both gases have the same molar mass.

(D) It cannot be determined unless the pressure of each sample is known.

Chapter 06 Part 1: Ideal Gas Law

Compound	Molar Mass	Balanced Equation for the Combustion of One Mole of the Compound	ΔH°_{comb} (kJ/mol _{rxn})
$\begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H}-\text{C}-\text{C}-\text{H} \\ \quad \\ \text{H} \quad \text{H} \end{array}$ Ethane	30. g/mol	$\text{C}_2\text{H}_6(g) + \frac{7}{2}\text{O}_2(g) \rightarrow 2\text{CO}_2(g) + 3\text{H}_2\text{O}(l)$	-1560
$\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\ddot{\text{O}}-\text{H} \\ \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \end{array}$ Propanol	60. g/mol	$\text{C}_3\text{H}_8\text{O}(l) + \frac{9}{2}\text{O}_2(g) \rightarrow 3\text{CO}_2(g) + 4\text{H}_2\text{O}(l)$	-2020
Unknown	?	$? + 5\text{O}_2(g) \rightarrow 4\text{CO}_2(g) + 4\text{H}_2\text{O}(l)$	-2240

8. A sealed 10.0 L flask at 400 K contains equimolar amounts of ethane and propanol in gaseous form. Which of the following statements concerning the average molecular speed of ethane and propanol is true?
- (A) The average molecular speed of ethane is less than the average molecular speed of propanol.
- (B) The average molecular speed of ethane is greater than the average molecular speed of propanol. ✓
- (C) The average molecular speed of ethane is equal to the average molecular speed of propanol.
- (D) The average molecular speeds of ethane and propanol cannot be compared without knowing the total pressure of the gas mixture.

Refer to three gases in identical rigid containers under the conditions given in the table below.

Container	A	B	C
Gas	Methane	Ethane	Butane
Formula	CH ₄	C ₂ H ₆	C ₄ H ₁₀
Molar mass (g/mol)	16	30.	58
Temperature (°C)	27	27	27
Pressure (atm)	2.0	4.0	2.0

9. The average kinetic energy of the gas molecules is
- (A) greatest in container A
- (B) greatest in container B
- (C) greatest in container C
- (D) the same in all three containers ✓


10. The density of the gas, in g/L, is

Chapter 06 Part 1: Ideal Gas Law

- (A) greatest in container A
- (B) greatest in container B
- (C) greatest in container C
- (D) the same in all three containers



11.




The experimental apparatus represented above is used to demonstrate the rates at which gases diffuse. When the cotton balls are placed in the ends of a tube at the same time, the gases diffuse from each end and meet somewhere in between, where they react to form a white solid. Which of the following combinations will produce a solid closest to the center of the tube?

- (A) HCl and CH_3NH_2
- (B) HCl and NH_3
- (C) HBr and CH_3NH_2
- (D) HBr and NH_3




Chapter 06 Part 1: Ideal Gas Law

12.



The diagram above shows the distribution of speeds for a sample of $\text{N}_2(g)$ at 25°C . Which of the following graphs shows the distribution of speeds for a sample of $\text{O}_2(g)$ at 25°C (dashed line) ?

Chapter 06 Part 1: Ideal Gas Law




13. When a sample of oxygen gas in a closed container of constant volume is heated until its absolute temperature is doubled, which of the following is also doubled?

- (A) The density of the gas
- (B) The pressure of the gas
- (C) The average velocity of the gas molecules
- (D) The number of molecules per cm^3
- (E) The potential energy of the molecules

Chapter 06 Part 1: Ideal Gas Law

14.



The volume of a sample of air in a cylinder with a movable piston is 2.0 L at a pressure P_1 , as shown in the diagram above. The volume is increased to 5.0 L as the temperature is held constant. The pressure of the air in the cylinder is now P_2 . What effect do the volume and pressure changes have on the average kinetic energy of the molecules in the sample?


- (A) The average kinetic energy increases.
- (B) The average kinetic energy decreases.

(C) The average kinetic energy stays the same.




- (D) It cannot be determined how the kinetic energy is affected without knowing P_1 and P_2 .

Chapter 06 Part 1: Ideal Gas Law



A sample of $C_4H_{10}(g)$ is placed in a rigid vessel with a movable piston. The volume of the gas is reduced by moving the piston downward while the temperature is maintained at $25^\circ C$, as shown above. The pressure of the gas versus its volume is plotted in the graph below.



15. Which of the following best explains the increase in pressure of the $C_4H_{10}(g)$ when its volume is decreased from 0.50 L to 0.21 L ?
- (A) The molecules strike the walls of the vessel more frequently. ✓
- (B) The molecules strike the walls of the vessel with different orientations.
- (C) The molecules strike the walls of the vessel with greater force.
- (D) The molecules decompose to produce a greater number of particles.
16. Which of the following best accounts for the pressure in the vessel remaining constant as the volume decreases below 0.21 L?

Chapter 06 Part 1: Ideal Gas Law

- (A) The density of the $\text{C}_4\text{H}_{10}(g)$ increases.
- (B) The surface area of the interior of the vessel decreases.
- (C) The temperature of the $\text{C}_4\text{H}_{10}(g)$ decreases.

(D) Some $\text{C}_4\text{H}_{10}(g)$ condenses to form $\text{C}_4\text{H}_{10}(l)$.



17. When 4.0 L of $\text{He}(g)$, 6.0 L of $\text{N}_2(g)$, and 10. L of $\text{Ar}(g)$, all at 0°C and 1.0 atm, are pumped into an evacuated 8.0 L rigid container, the final pressure in the container at 0°C is

- (A) 0.5 atm
- (B) 1.0 atm

(C) 2.5 atm



(D) 4.0 atm

18. When 6.0 L of $\text{He}(g)$ and 10. L of $\text{N}_2(g)$, both at 0°C and 1.0 atm, are pumped into an evacuated 4.0 L rigid container, the final pressure in the container at 0°C is

(A) 2.0 atm

(B) 4.0 atm



(C) 6.4 atm

(D) 8.8 atm

(E) 16 atm

19. A sample of an ideal gas in a sealed container occupies a volume of 50.0 L at 200 K. The temperature of the gas is changed to 400 K, and the pressure doubles. Which of the following must occur and why?

- (A) Volume must decrease because the pressure is higher.
- (B) Volume must increase because the temperature is higher.

(C) Volume must remain the same because the pressure and temperature change proportionately.



(D) Volume cannot be predicted without knowing the number of moles of gas in the sample.

Chapter 06 Part 1: Ideal Gas Law

The following questions refer to the below.

The table below contains information about samples of four different gases at 273 K. The samples are in four identical rigid containers numbered 1 through 4.

Container	Gas	Pressure (atm)	Mass of Sample (g)
1	He	2.00	?
2	Ne	2.00	?
3	?	2.00	16.0
4	SO ₂	1.96	64.1

20. Under the conditions given, consider containers 1, 2, and 4 only. The average speed of the gas particles is

- (A) greatest in container 1
- (B) greatest in container 2
- (C) greatest in container 4
- (D) the same in containers 1, 2, and 4



21. On the basis of the data provided above, the gas in container 3 could be

- (A) CH₄
- (B) O₂
- (C) Ar
- (D) CO₂



22. Two flexible containers for gases are at the same temperature and pressure. One holds 0.50 gram of hydrogen and the other holds 8.0 grams of oxygen. Which of the following statements regarding these gas samples is FALSE?

- (A) The volume of the hydrogen container is the same as the volume of the oxygen container.
- (B) The number of molecules in the hydrogen container is the same as the number of molecules in the oxygen container.
- (C) The density of the hydrogen sample is less than that of the oxygen sample.
- (D) The average kinetic energy of the hydrogen molecules is the same as the average kinetic energy of the oxygen molecules.

- (E) The average speed of the hydrogen molecules is the same as the average speed of the oxygen molecules.




Chapter 06 Part 1: Ideal Gas Law

23. A 0.5 mol sample of $\text{He}(g)$ and a 0.5 mol sample of $\text{Ne}(g)$ are placed separately in two 10.0 L rigid containers at 25°C . Each container has a pinhole opening. Which of the gases, $\text{He}(g)$ or $\text{Ne}(g)$, will escape faster through the pinhole and why?

- (A) $\text{He}(g)$ will escape faster because the $\text{He}(g)$ atoms are moving at a higher average speed than the $\text{Ne}(g)$ atoms. ✓
- (B) $\text{Ne}(g)$ will escape faster because its initial pressure in the container is higher.
- (C) $\text{Ne}(g)$ will escape faster because the $\text{Ne}(g)$ atoms have a higher average kinetic energy than the $\text{He}(g)$ atoms.
- (D) Both gases will escape at the same rate because the atoms of both gases have the same average kinetic energy.

24.




Based on the Maxwell-Boltzmann distributions at 300 K shown above, which of the following gases could be represented by curve X?

- (A) $\text{Br}_2(g)$ ✓
- (B) $\text{CH}_4(g)$
- (C) $\text{He}(g)$
- (D) $\text{N}_2(g)$
25. A 1 L sample of helium gas at 25°C and 1 atm is combined with a 1 L sample of neon gas at 25°C and 1 atm. The temperature is kept constant. Which of the following statements about combining the gases is correct?
- (A) The average speed of the helium atoms increases when the gases are combined.
- (B) The average speed of the neon atoms increases when the gases are combined.
- (C) The average kinetic energy of the helium atoms increases when the gases are combined.
- (D) The average kinetic energy of the helium atoms and neon atoms do not change when the gases are combined. ✓

Chapter 06 Part 1: Ideal Gas Law

26.



A sample of an unknown gas from a cylinder is collected over water in the apparatus shown above. After all the gas sample has been collected, the water levels inside and outside the gas collection tube are made the same.

Measurements that must be made to calculate the molar mass of the gas include all of the following EXCEPT

- (A) atmospheric pressure
- (B) temperature of the water
- (C) volume of gas in the gas-collection tube
- (D) initial and final mass of the gas cylinder
- (E) mass of the water in the apparatus



Directions: Each set of lettered choices below refers to the numbered statements immediately following it. Select the one lettered choice that best fits the statement. A choice may be used once, more than once, or not at all.

The following questions refer to the 1 L flasks shown below.

- | | | | | |
|---|---|---|---|---|
| (A) | (B) | (C) | (D) | (E) |
| | | | | |
| $T = 30^{\circ}\text{C}$
$P = 1 \text{ atm}$ | $T = 50^{\circ}\text{C}$
$P = 0.5 \text{ atm}$ | $T = 40^{\circ}\text{C}$
$P = 2 \text{ atm}$ | $T = 50^{\circ}\text{C}$
$P = 1 \text{ atm}$ | $T = 40^{\circ}\text{C}$
$P = 0.5 \text{ atm}$ |

27. Which flask contains the sample with the greatest density?

Chapter 06 Part 1: Ideal Gas Law

(A) A

(B) B

(C) C



(D) D

(E) E

28. In which flask do the molecules have the greatest average speed?

(A) A

(B) B



(C) C

(D) D

(E) E

29. Which flask contains the smallest number of moles of gas?

(A) A

(B) B



(C) C

(D) D

(E) E

30. A gas mixture at 0°C and 1.0 atm contains 0.010 mol of H_2 , 0.015 mol of O_2 , and 0.025 mol of N_2 . Assuming ideal behavior, what is the partial pressure of hydrogen gas (H_2) in the mixture?

(A) About 0.010 atm, because there is 0.010 mol of H_2 in the sample.(B) About 0.050 atm, because there is 0.050 mol of gases at 0°C and 1.0 atm.(C) About 0.20 atm, because H_2 comprises 20% of the total number of moles of gas.(D) About 0.40 atm, because the mole ratio of $\text{H}_2:\text{O}_2:\text{N}_2$ is 0.4 : 0.6 : 1.

31. Equal masses of three different ideal gases, X, Y, and Z, are mixed in a sealed rigid container. If the temperature of the system remains constant, which of the following statements about the partial pressure of gas X is correct?

(A) It is equal to $\frac{1}{3}$ the total pressure

(B) It depends on the intermolecular forces of attraction between molecules of X, Y, and Z.

(C) It depends on the relative molecular masses of X, Y, and Z.



(D) It depends on the average distance traveled between molecular collisions.

(E) It can be calculated with knowledge only of the volume of the container.

Chapter 06 Part 1: Ideal Gas Law

32.





Diagram 1 above shows equimolar samples of two gases inside a container fitted with a removable barrier placed so that each gas occupies the same volume. The barrier is carefully removed as the temperature is held constant. Diagram 2 above shows the gases soon after the barrier is removed. Which statement describes the changes to the initial pressure of each gas and the final partial pressure of each gas in the mixture and also indicates the final total pressure?

- (A) The partial pressure of each gas in the mixture is double its initial pressure; the final total pressure is half the sum of the initial pressures of the two gases.
- (B) The partial pressure of each gas in the mixture is double its initial pressure; the final total pressure is twice the sum of the initial pressures of the two gases.
- (C) The partial pressure of each gas in the mixture is half its initial pressure; the final total pressure is half the sum of the initial pressures of the two gases. ✓
- (D) The partial pressure of each gas in the mixture is half its initial pressure; the final total pressure is the same as the sum of the initial pressures of the two gases.

33.



The reaction between $\text{NO}(\text{g})$ and $\text{O}_2(\text{g})$ to produce $\text{NO}_2(\text{g})$ in a rigid reaction vessel is represented in the diagram above. The pressure inside the container is recorded using a pressure gauge. Which of the following statements correctly predicts the change in pressure as the reaction goes to completion at constant temperature, and provides the correct explanation?

Chapter 06 Part 1: Ideal Gas Law

- (A) The pressure will increase because the product molecules have a greater mass than either of the reactant molecules.
- (B) The pressure will decrease because there are fewer molecules of product than of reactants. ✓
- (C) The pressure will decrease because the product molecules have a lower average speed than the reactant molecules.
- (D) The pressure will not change because the total mass of the product molecules is the same as the total mass of the reactant molecules.

Refer to the following.



$\text{PCl}_5(\text{g})$ decomposes into $\text{PCl}_3(\text{g})$ and $\text{Cl}_2(\text{g})$ according to the equation above. A pure sample of $\text{PCl}_5(\text{g})$ is placed in a rigid, evacuated 1.00 L container. The initial pressure of the $\text{PCl}_5(\text{g})$ is 1.00 atm. The temperature is held constant until the $\text{PCl}_5(\text{g})$ reaches equilibrium with its decomposition products. The figures below show the initial and equilibrium conditions of the system.




Figure 1: Initial




Figure 2: Equilibrium


34. Which of the following is the most likely cause for the increase in pressure observed in the container as the reaction reaches equilibrium?
- (A) A decrease in the strength of intermolecular attractions among molecules in the flask
- (B) An increase in the strength of intermolecular attractions among molecules in the flask
- (C) An increase in the number of molecules, which increases the frequency of collisions with the walls of the container. ✓
- (D) An increase in the speed of the molecules that then collide with the walls of the container with greater force
35. The pressure, in atm, exerted by 1.85 mol of an ideal gas placed in a 3.00 L container at 35.0°C is given by which of the following expressions?
- (A) $\frac{(1.85)(0.0821)(308)}{3.00} \text{ atm}$ ✓
- (B) $\frac{(1.85)(35.0)}{(0.0821)(3.00)} \text{ atm}$
- (C) $\frac{3.00}{(1.85)(0.0821)(308)} \text{ atm}$
- (D) $\frac{(1.85)(8.314)(308)}{3.00} \text{ atm}$
- (E) $\frac{(3.00)(1.85)}{(0.0821)(35.0)} \text{ atm}$

Chapter 06 Part 1: Ideal Gas Law

36. A sealed vessel contains 0.200 mol of oxygen gas, 0.100 mol of nitrogen gas, and 0.200 mol of argon gas. The total pressure of the gas mixture is 5.00 atm. The partial pressure of the argon is
- (A) 0.200 atm
 - (B) 0.500 atm
 - (C) 1.00 atm
 - (D) 2.00 atm
 - (E) 5.00 atm
37. Equal masses of He and Ne are placed in a sealed container. What is the partial pressure of He if the total pressure in the container is 6 atm?
- (A) 1 atm
 - (B) 2 atm
 - (C) 3 atm
 - (D) 4 atm
 - (E) 5 atm
38. A flask contains 0.25 mole of $\text{SO}_2(g)$, 0.50 mole of $\text{CH}_4(g)$, and 0.50 mole of $\text{O}_2(g)$. The total pressure of the gases in the flask is 800 mm Hg. What is the partial pressure of the $\text{SO}_2(g)$ in the flask?
- (A) 800 mm Hg
 - (B) 600 mm Hg
 - (C) 250 mm Hg
 - (D) 200 mm Hg
 - (E) 160 mm Hg
39. Equal numbers of moles of $\text{He}(g)$, $\text{Ar}(g)$, and $\text{Ne}(g)$ are placed in a glass vessel at room temperature. If the vessel has a pinhole-sized leak, which of the following will be true regarding the relative values of the partial pressures of the gases remaining in the vessel after some of the gas mixture has effused?
- (A) $P_{\text{He}} < P_{\text{Ne}} < P_{\text{Ar}}$
 - (B) $P_{\text{He}} < P_{\text{Ar}} < P_{\text{Ne}}$
 - (C) $P_{\text{Ne}} < P_{\text{Ar}} < P_{\text{He}}$
 - (D) $P_{\text{Ar}} < P_{\text{He}} < P_{\text{Ne}}$
 - (E) $P_{\text{He}} = P_{\text{Ar}} = P_{\text{Ne}}$

Chapter 06 Part 1: Ideal Gas Law

40.




The diagram above shows the distribution of molecular speed in a sample of gas at two different temperatures, T_1 and T_2 . Which of the following is a correct interpretation of the information provided by the diagram?

- (A) The average speed of the molecules is greater at T_1 than at T_2 .
- (B) The taller peak corresponds to the higher temperature, T_1 .
- (C) There are fewer molecules of gas at T_2 than at T_1 .
- (D) The average kinetic energy of the molecules is greater at T_2 than at T_1 .



41.



The graph above shows the distribution of molecular speeds for four different gases at the same temperature. What property of the different gases can be correctly ranked using information from the graph, and why?

Chapter 06 Part 1: Ideal Gas Law


- (A) The densities of the gases, because as the density of a gas increases, the average speed of its molecules decreases.
- (B) The pressures of the gases, because the pressure exerted by a gas depends on the average speed with which its molecules are moving.
- (C) The volumes of the gases, because at a fixed temperature the volume of a gas can be calculated using the equation $PV = nRT$.
- (D) The molecular masses of the gases, because the gas molecules have the same average kinetic energy and mass can be calculated using the equation $KE_{\text{avg}} = \frac{1}{2}mv^2$. ✓

42. A 2 L sample of $\text{N}_2(g)$ and a 1 L sample of $\text{Ar}(g)$, each originally at 1 atm and 0°C , are combined in a 1 L tank. If the temperature is held constant, what is the total pressure of the gases in the tank?

- (A) 1 atm
- (B) 2 atm
- (C) 3 atm ✓
- (D) 4 atm
- (E) 5 atm

43. An equimolar mixture of $\text{N}_2(g)$ and $\text{Ar}(g)$ is kept inside a rigid container at a constant temperature of 300 K. The initial partial pressure of Ar in the mixture is 0.75 atm. An additional amount of Ar was added to the container, enough to double the number of moles of Ar gas in the mixture. Assuming ideal behavior, what is the final pressure of the gas mixture after the addition of the Ar gas?

- (A) 0.75 atm, because increasing the partial pressure of Ar decreases the partial pressure of N_2 .
- (B) 1.13 atm, because 33% of the moles of gas are N_2 .
- (C) 1.50 atm, because the number of moles of N_2 did not change.
- (D) 2.25 atm, because doubling the number of moles of Ar doubles its partial pressure. ✓




The figure above represents three sealed 1.0 L vessels, each containing a different inert gas at 298 K. The pressure of Ar in the first vessel is 2.0 atm. The ratio of the numbers of Ar, Ne, and He atoms in the vessels is 2:1:6, respectively. After all the gases are combined in a previously evacuated 2.0 L vessel, what is the total pressure of the gases at 298 K?


- (A) 3.0 atm
- (B) 4.5 atm ✓
- (C) 9.0 atm
- (D) 18 atm

Chapter 06 Part 1: Ideal Gas Law

45.



A mixture of two gases, 0.01 mol of $\text{C}_4\text{H}_{10}(g)$ and 0.065 mol of $\text{O}_2(g)$, is pumped into a cylinder with a movable piston, as shown above. The mixture, originally at 200°C and 1.0 atm, is sparked and the reaction represented below occurs.



Which of the following is true after the product gases return to the original temperature and pressure, and why will the change occur? (Assume all gases behave ideally.)

- (A) The piston will be higher than its original position, because the cylinder will contain a greater number of gas molecules. ✓
- (B) The position of the piston will be unchanged, because the total mass of the gases in the cylinder does not change.
- (C) The position of the piston will be unchanged, because the temperature and pressure of the contents of the cylinder remain the same.
- (D) The piston will be lower than its original position, because the product molecules are smaller than the reactant molecules.
46. A balloon filled with 0.25 mol of $\text{He}(g)$ at 273 K and 1 atm is allowed to rise through the atmosphere. Which of the following explains what happens to the volume of the balloon as it rises from ground level to an altitude where the air temperature is 220 K and the air pressure is 0.1 atm?

Chapter 06 Part 1: Ideal Gas Law

- (A) The volume will increase because the decrease in air pressure will have a greater effect than the decrease in temperature. ✓
- (B) The volume will remain unchanged because of the counteracting effects of the decrease in temperature and the decrease in air pressure.
- (C) The volume will decrease because the decrease in temperature will have a greater effect than the decrease in air pressure.
- (D) It cannot be determined whether the volume of the balloon will increase, decrease, or remain the same without knowing the initial volume of the balloon.
47. A rigid metal tank contains oxygen gas. Which of the following applies to the gas in the tank when additional oxygen is added at constant temperature?
- (A) The volume of the gas increases.
- (B) The pressure of the gas decreases.
- (C) The average speed of the gas molecules remains the same. ✓
- (D) The total number of gas molecules remains the same.
- (E) The average distance between the gas molecules increases.
48. A sample of an ideal gas is cooled from 50.0°C to 25.0°C in a sealed container of constant volume. Which of the following values for the gas will decrease?
- I. The average molecular mass of the gas
- II. The average distance between the molecules
- III. The average speed of the molecules
- (A) I only
- (B) II only
- (C) III only ✓
- (D) I and III
- (E) II and III

Chapter 06 Part 1: Ideal Gas Law

49. A student was assigned the task of determining the molar mass of an unknown gas. The student measured the mass of a sealed 843 mL rigid flask that contained dry air. The student then flushed the flask with the unknown gas, resealed it, and measured the mass again. Both the air and the unknown gas were at 23.0°C and 750. torr. The data for the experiment are shown in the table below.

Volume of sealed flask	843 mL
Mass of sealed flask and dry air	157.70 g
Mass of sealed flask and unknown gas	158.08 g

- Calculate the mass, in grams, of the dry air that was in the sealed flask. (The density of dry air is 1.18 g L^{-1} at 23.0°C and 750. torr.)
- Calculate the mass, in grams, of the sealed flask itself (i.e., if it had no air in it).
- Calculate the mass, in grams, of the unknown gas that was added to the sealed flask.
- Using the information above, calculate the value of the molar mass of the unknown gas.

After the experiment was completed, the instructor informed the student that the unknown gas was carbon dioxide (44.0 g mol^{-1}).

- Calculate the percent error in the value of the molar mass calculated in part (d).
- For each of the following two possible occurrences, indicate whether it by itself could have been responsible for the error in the student's experimental result. You need not include any calculations with your answer. For each of the possible occurrences, justify your answer.

Occurrence 1: The flask was incompletely flushed with $\text{CO}_2(\text{g})$, resulting in some dry air remaining in the flask.

Occurrence 2: The temperature of the air was 23.0°C, but the temperature of the $\text{CO}_2(\text{g})$ was lower than the reported 23.0°C.

- Describe the steps of a laboratory method that the student could use to verify that the volume of the rigid flask is 843 mL at 23.0°C. You need not include any calculations with your answer.

Part A

1 point is earned for the correct setup and calculation of mass.

$$m = D \times V = (1.18 \text{ g L}^{-1})(0.843 \text{ L}) = 0.995 \text{ g}$$

Note: Significant figures were checked in this problem: parts (a) and (d) were scored with ± 1 significant figure needed, and parts (b) and (c) were scored with the correct number of significant figures needed for the subtraction.

Chapter 06 Part 1: Ideal Gas Law

0

1

The student response earns one of the following points:

1 point is earned for the correct setup and calculation of mass.

$$m = D \times V = (1.18 \text{ g L}^{-1})(0.843 \text{ L}) = 0.995 \text{ g}$$

Note: Significant figures were checked in this problem: parts (a) and (d) were scored with ± 1 significant figure needed, and parts (b) and (c) were scored with the correct number of significant figures needed for the subtraction.

Part B

1 point is earned for subtracting the answer in part (a) from 157.70 g.

$$157.70 \text{ g} - 0.995 \text{ g} = 156.71 \text{ g}$$

Note: Significant figures were checked in this problem: parts (a) and (d) were scored with ± 1 significant figure needed, and parts (b) and (c) were scored with the correct number of significant figures needed for the subtraction.



0

1

The student response earns one of the following points:

1 point is earned for subtracting the answer in part (a) from 157.70 g.

$$157.70 \text{ g} - 0.995 \text{ g} = 156.71 \text{ g}$$

Note: Significant figures were checked in this problem: parts (a) and (d) were scored with ± 1 significant figure needed, and parts (b) and (c) were scored with the correct number of significant figures needed for the subtraction.

Part C

1 point is earned for subtracting the answer in part (b) from 158.08 g.

$$158.08 \text{ g} - 156.71 \text{ g} = 1.37 \text{ g}$$

Note: Significant figures were checked in this problem: parts (a) and (d) were scored with ± 1 significant figure needed, and parts (b) and (c) were scored with the correct number of significant figures needed for the subtraction.

Chapter 06 Part 1: Ideal Gas Law



0	1
---	---

The student response earns one of the following points:

1 point is earned for subtracting the answer in part (b) from 158.08 g.

$$158.08 \text{ g} - 156.71 \text{ g} = 1.37 \text{ g}$$

Note: Significant figures were checked in this problem: parts (a) and (d) were scored with ± 1 significant figure needed, and parts (b) and (c) were scored with the correct number of significant figures needed for the subtraction.

Part D

1 point is earned for the conversion of pressure (if necessary) and temperature and the use of the appropriate R .

1 point is earned for the correct setup and calculation of moles of gas.

$$n = \frac{PV}{RT} = \frac{\left(\frac{750}{760} \text{ atm}\right)(0.843 \text{ L})}{(0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1})(296 \text{ K})} = 0.0342 \text{ mol}$$

1 point is earned for the correct setup and calculation of molar mass.

OR

If calculation is done in a single step, 1 point is earned for the correct P and T , 1 point is earned for the correct density, and 1 point is earned for the correct answer.

$$\text{molar mass} = \frac{1.37 \text{ g}}{0.0342 \text{ mol}} = 40.1 \text{ g mol}^{-1}$$

OR

$$\text{molar mass} = \frac{DRT}{P} = \frac{\left(\frac{1.37 \text{ g}}{0.843 \text{ L}}\right)(0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1})(296 \text{ K})}{\left(\frac{750}{760} \text{ atm}\right)}$$

$$= 40.0 \text{ g mol}^{-1}$$

Note: Significant figures were checked in this problem: parts (a) and (d) were scored with ± 1 significant figure needed, and parts (b) and (c) were scored with the correct number of significant figures needed for the subtraction.



0	1	2	3
---	---	---	---

The student response earns three of the following points:

Chapter 06 Part 1: Ideal Gas Law

1 point is earned for the conversion of pressure (if necessary) and temperature and the use of the appropriate R .

1 point is earned for the correct setup and calculation of moles of gas.

$$n = \frac{PV}{RT} = \frac{\left(\frac{750}{760} \text{ atm}\right)(0.843 \text{ L})}{(0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1})(296 \text{ K})} = 0.0342 \text{ mol}$$

1 point is earned for the correct setup and calculation of molar mass.

OR

If calculation is done in a single step, 1 point is earned for the correct P and T , 1 point is earned for the correct density, and 1 point is earned for the correct answer.

$$\text{molar mass} = \frac{1.37 \text{ g}}{0.0342 \text{ mol}} = 40.1 \text{ g mol}^{-1}$$

OR

$$\begin{aligned} \text{molar mass} &= \frac{DRT}{P} \\ &= \frac{\left(\frac{1.37 \text{ g}}{0.843 \text{ L}}\right)(0.0821 \text{ L atm mol}^{-1} \text{ K}^{-1})(296 \text{ K})}{\left(\frac{750}{760} \text{ atm}\right)} \\ &= 40.0 \text{ g mol}^{-1} \end{aligned}$$

Note: Significant figures were checked in this problem: parts (a) and (d) were scored with ± 1 significant figure needed, and parts (b) and (c) were scored with the correct number of significant figures needed for the subtraction.

Part E

1 point is earned for the correct setup and answer.

$$\text{percent error} = \frac{|44.0 \text{ g mol}^{-1} - 40.1 \text{ g mol}^{-1}|}{44.0 \text{ g mol}^{-1}} \times 100 = 8.9\%$$



0	1
---	---

The student response earns one of the following points:

1 point is earned for the correct setup and answer.

$$\text{percent error} = \frac{|44.0 \text{ g mol}^{-1} - 40.1 \text{ g mol}^{-1}|}{44.0 \text{ g mol}^{-1}} \times 100 = 8.9\%$$

Part F

Occurrence 1:

1 point is earned for the correct reasoning and conclusion.

Chapter 06 Part 1: Ideal Gas Law

This occurrence could have been responsible.

The dry air left in the flask is less dense (or has a lower molar mass) than CO₂ gas at the given T and P . This would result in a lower mass of gas in the flask and a lower result for the molar mass of the unknown gas.

Occurrence 2:

1 point is earned for the correct reasoning and conclusion.

This occurrence could not have been responsible.

The density of CO₂ is greater at the lower temperature. A larger mass of CO₂ would be in the flask than if the CO₂ had been at 23.0°C, resulting in a higher calculated molar mass for the unknown gas.



0	1	2
---	---	---

The student response earns two of the following points:

Occurrence 1:

1 point is earned for the correct reasoning and conclusion.

This occurrence could have been responsible.

The dry air left in the flask is less dense (or has a lower molar mass) than CO₂ gas at the given T and P . This would result in a lower mass of gas in the flask and a lower result for the molar mass of the unknown gas.

Occurrence 2:

1 point is earned for the correct reasoning and conclusion.

This occurrence could not have been responsible.

The density of CO₂ is greater at the lower temperature. A larger mass of CO₂ would be in the flask than if the CO₂ had been at 23.0°C, resulting in a higher calculated molar mass for the unknown gas.

Part G

1 point is earned for a valid method.

Valid methods include the following:

- Find the mass of the empty flask. Fill the flask with a liquid of known density (e.g., water at 23°C), and measure the mass of the liquid-filled flask. Subtract to find the mass of the liquid. Using the known density and mass, calculate the volume.
- Measure 843 mL of a liquid (e.g., water) in a 1,000 mL graduated cylinder and transfer the liquid quantitatively into the flask to see if the water fills the flask completely.

Chapter 06 Part 1: Ideal Gas Law

0	1
---	---

The student response earns one of the following points:

1 point is earned for a valid method.

Valid methods include the following:

- Find the mass of the empty flask. Fill the flask with a liquid of known density (e.g., water at 23°C), and measure the mass of the liquid-filled flask. Subtract to find the mass of the liquid. Using the known density and mass, calculate the volume.
- Measure 843 mL of a liquid (e.g., water) in a 1,000 mL graduated cylinder and transfer the liquid quantitatively into the flask to see if the water fills the flask completely.


Chapter 06 Part 1: Ideal Gas Law

- 50.** For parts of the free response question that require calculations, clearly show the method used and the steps involved in arriving at your answers. You must show your work to receive credit for your answer. Examples and equations may be included in your answers where appropriate.

Dry air is comprised of N_2 , O_2 , and traces of other gases.

A student obtains a 2.00 L container filled with a sample of dry air at a total pressure of 1.30 atm and a temperature of 295 K.

- (a) Calculate the total number of moles of gas in the container.
- (b) The student cools the gaseous mixture until either N_2 or O_2 condenses. Which gas will condense first? Justify your answer in terms of the types and relative strengths of the intermolecular forces in N_2 and O_2 .
- (c) Using the legend given, draw at least three molecules of each type of gas in the sealed container represented below when one of the substances has condensed, but the other substance is still a gas.



- (d) In this sample, 78% of the molecules are N_2 and 21% of the molecules are O_2 . Calculate the total number of grams of the substance that condenses first.

Part A

Select a point value to view scoring criteria, solutions, and/or examples and to score the response.

Chapter 06 Part 1: Ideal Gas Law



0	1
---	---

The response provides the correct calculated value.

- $PV = nRT$

$$n = \frac{PV}{RT} = \frac{(1.30 \text{ atm})(2.00 \text{ L})}{(0.08206 \text{ L atm mol}^{-1} \text{ K}^{-1})(295 \text{ K})} = 0.107 \text{ mol}$$

Part B

Select a point value to view scoring criteria, solutions, and/or examples and to score the response.



0	1
---	---

The response provides the correct answer and a valid justification.

- O_2 will condense first. Both N_2 and O_2 have only dispersion forces as their intermolecular forces. Because O_2 has more electrons, it has a higher polarizability. Therefore, O_2 will have stronger dispersion forces and thus will condense at the higher temperature.

Part C

Select a point value to view scoring criteria, solutions, and/or examples and to score the response.




0	1
---	---

The response provides a correct diagram, consistent with part (b).

- Diagram should show one substance is a gas and the other substance is a liquid.

Chapter 06 Part 1: Ideal Gas Law



Part D

Select a point value to view scoring criteria, solutions, and/or examples and to score the response.



0

1

The response provides the correct calculated value, consistent with part (b).

- $(0.21)(0.107 \text{ mol})(32.00 \text{ g/mol}) = 0.72 \text{ g O}_2$