Stochastic Calculus

MATH-GA 2903

Yao Xiao

Spring 2023

Contents

1 Construction of Brownian Motion **1**

1.1 Brownian Motion and Its Motivation . . . 1

1.2 Symmetric Random Walk . . . 2

1.2.1 Increments of the SRW . . . 2

1.2.2 Martingale Property of the SRW . . . 2

1.2.3 Quadratic Variation of the SRW . . . 3

1.3 Scaled Symmetric Random Walk . . . 3

1.3.1 Limiting Distribution of the Scaled SRW . . . 4

1.4 Brownian Motion: A Formal Definition . . . 5

2 Properties of the Brownian Motion **5**

2.1 Covariance of Brownian Motion . . . 5

2.2 Non-differentiability of Brownian Paths . . . 6

2.3 Scaling Properties of the Brownian Motion . . . 6

2.4 Quadratic Variation of Brownian Motion . . . 7

3/27 Lecture

1 Construction of Brownian Motion

1.1 Brownian Motion and Its Motivation

Loosely speaking, Brownian motion is a "random path" that satisfies the following properties.

- The path is continuous.
- At any given point in time, where the path goes next is *independent* of where the path came from.
- At any given point in time, where the path ends up in any finite time is normally distributed.

1.2 Symmetric Random Walk

It is always easier to start with a discrete case. Consider a symmetric random walk (SRW). Let X_j's be independent and identically distributed such that $\mathbb{P}[X_j = 1] = \mathbb{P}[X_j = -1] = 1/2$. We now define $M(n)$ such that

$$M(0) = 0, \qquad M(n) = \sum_{j=1}^{n} X_j. \tag{1}$$

The process $M(n)$ is called a symmetric random walk. With each step, it either steps up or steps down one unit, and each of the two possible moves is equally likely.

1.2.1 Increments of the SRW

SRW has independent increments, which means that if we choose nonnegative integers $0 = n_0 < n_1 < \cdots < n_k$, then the random variables

$$M(n_1) = M(n_1) - M(n_0),\ M(n_2) - M(n_1),\ \cdots,\ M(n_k) - M(n_{k-1}) \tag{2}$$

are independent. Each of these random variables

$$M(n_{i+1}) - M(n_i) = \sum_{j=n_i+1}^{n_{i+1}} X_j \tag{3}$$

is called an increment of the random walk. It is the change in the position of the random walk between times n_i and n_{i+1}. Increments over non-overlapping time intervals are indeed independent, since X_j's are independent. Moreover,

$$\mathbb{E}[M(n_{i+1}) - M(n_i)] = \sum_{j=n_i+1}^{n_{i+1}} \mathbb{E}[X_j] = 0, \tag{4}$$

and

$$\mathrm{Var}[M(n_{i+1}) - M(n_i)] = \sum_{j=n_i+1}^{n_{i+1}} \mathrm{Var}[X_j] = n_{i+1} - n_i, \tag{5}$$

since $\mathrm{Var}[X_j] = 1$ and this holds by independence.

1.2.2 Martingale Property of the SRW

In order to define martingale we need the notion of σ-algebra. There is a very important, nontechnical reason to include σ-algebras in the study of stochastic processes, and that is to keep track of the amount of information. The temporal feature of a stochastic process suggests a flow of time, in which at each moment $t \geq 0$, we can talk about the past, the present, and the future, and can ask about how much an observer of the process knows about it at present, as compared to how much he know at some point in the past or will know at some time in the future.

Definition 1.1. A *filtration* on $(\Omega, \mathscr{F})$is a family $\{\mathscr{F}_t\}_{t\geq 0}$ of σ-algebras $\mathscr{F}_t \subseteq \mathscr{F}$, such that

$$0 \leq s \leq t \implies \mathscr{F}_s \subseteq \mathscr{F}_t, \tag{6}$$

i.e., $\{\mathscr{F}_t\}_{t\geq 0}$ is increasing.

Thus informally, one can think about the σ-algebra $\mathscr{F}_t$ in the above definition as the "knowledge" available at time t. The fact that $\{\mathscr{F}_t\}_{t\geq 0}$ is monotonically increasing just reflects the fact that as time goes on, the amount of available information cannot decrease.

Definition 1.2. A stochastic process $\{Z_t\}_{t\geq 0}$ is called a *martingale* with respect to a filtration $\{\mathscr{F}_T\}_{t\geq 0}$ if

(1) Z_t is $\mathscr{F}_t$-measurable for all t.

(2) $\mathbb{E}[|Z_t|] < \infty$ for all t.

(3) $\mathbb{E}[Z_t|\mathscr{F}_s] = Z_s$ for all $s \leq t$.

The first two conditions are for technical reasons, since we can only define measures on measurable sets, and we assume that the first moments are finite. The most important property in the above definition is the third property. Simply put, it states that the "best guess" for X_t given information available at moment s is X_s, and thus the process X_t has neither upward or downward drift.

Now we want to check that the SRW is a martingale. First, we need to specify a filtration $\{\mathscr{F}_t\}_{t\geq 0}$. Let $\mathscr{F}_n$ be the σ-algebra (read information) generated by the set of random variables $X_1, X_2, \cdots, X_n$.

(1) $M(n)$ is $\mathscr{F}_n$-measurable since it depends only on X_j, $j \leq n$, *i.e.*, on the information available at time n.

(2) From the fact that $|M(t)| \leq t$, we can easily see that $\mathbb{E}[|M(t)|] \leq t < \infty$.

(3) Using the fact that $\mathbb{E}[M(s)|\mathscr{F}_s] = M(s)$ and the fact that the random variable $M(t) - M(s)$ is independent of σ-algebra $\mathscr{F}_s$, we can deduce that

$$\begin{aligned}\mathbb{E}[M(t)|\mathscr{F}_s] &= \mathbb{E}[M(t) - M(s) + M(s)|\mathscr{F}_s] = \mathbb{E}[M(t) - M(s)|\mathscr{F}_s] + \mathbb{E}[M(s)|\mathscr{F}_s] \\ &= \mathbb{E}[M(t) - M(s)] + M(s) = 0 + M(s) = M(s).\end{aligned} \tag{7}$$

1.2.3 Quadratic Variation of the SRW

The quadratic variation of a discrete stochastic process up to time t is defined as

$$\langle M, M\rangle_t = \sum_{j=1}^{t} (M(j) - M(j-1))^2. \tag{8}$$

The quadratic variation up to time t along the path is computed by taking all the one-step increments $M(j)-M(j-1)$ along that path, squaring these increments and then summing them up. Clearly, the SRM increments can only take values ± 1, and thus $\langle M, M\rangle_t = t$. Note that in general, the quadratic variation of a stochastic process is stochastic itself. The fact that the quadratic variation in this case is deterministic is a special feature of the SRW.

1.3 Scaled Symmetric Random Walk

To approximate a Brownian motion, we speed up time and scale down the step size of an SRW. More precisely, we fix a positive integer n and define the scaled SRW at rational points k/n as

$$B^{(n)}\left(\frac{k}{n}\right) = \frac{1}{\sqrt{n}} M(k). \tag{9}$$

At all other points we define $B^{(n)}(t)$ as the linear interpolation between its values at the nearest points of the form k/n. The following properties of the scaled SRW could be easily proved using the corresponding properties of the SRW and we leave their proof as an exercise.

(1) Independence of increments, *i.e.*, for all rational numbers $0 = t_0 < t_1 < \cdots < t_n$ of the form k/n, the random variables

$$B^{(n)}(t_1) - B^{(n)}(t_0),\ B^{(n)}(t_2) - B^{(n)}(t_1),\ \cdots,\ B^{(n)}(t_n) - B^{(n)}(t_{n-1}) \tag{10}$$

are independent.

(2) $\mathbb{E}[B^{(n)}(t) - B^{(n)}(s)] = 0$, $\mathrm{Var}[B^{(n)}(t) - B^{(n)}(s)] = t - s$.

(3) $\mathbb{E}[B^{(n)}|\mathscr{F}_s] = B^{(n)}(s)$.

(4) The quadratic variation of the scaled SRW $\left\langle B^{(n)}, B^{(n)}\right\rangle_t = t$.

1.3.1 Limiting Distribution of the Scaled SRW

Now the idea is to let $n \to \infty$ in the scaled SRW and in the limit we obtain something satisfying the intuitive properties outlined for Brownian motion. By definition,

$$B^{(n)}(t) = \frac{1}{\sqrt{n}} M_{nt}. \tag{11}$$

Since $M_k = \sum_{j=1}^{k} X_j$ has a binomial distribution with parameters k and $1/2$, we can easily calculate the distribution of $B^{(n)}(t)$. In particular, one can draw the histogram of $B^{(n)}(t)$. We know that the random variable $B^{(n)}(t)$ has mean 0 and variance t. If we draw on top of the histogram of $B^{(n)}(t)$ the graph of normal density with mean 0 and variance t, we will see that the distribution of $B^{(n)}(t)$ is nearly normal. In fact, the central limit theorem asserts that $B^{(n)}(t)$ converges *in distribution* to the normal random variable $\mathscr{N}(0,t)$, but let us not use this theorem and prove the convergence of $b^{(n)}(t)$ from scratch. The whole point of this exercise is to learn a very powerful mathematical tool known as *characteristic functions*.

4/3 Lecture

Definition 1.3. Let X be a random variable with distribution $\mathbb{P}$. The *characteristic function* of X is defined as

$$\phi_X(u) = \mathbb{E}[e^{iuX}] = \int e^{iux} d\mathbb{P}[x] \tag{12}$$

for every $u \in \mathbb{R}$.

Example 1.4. (1) If $X = \mathrm{Ber}(p)$, then

$$\phi_X(u) = p(e^{iu} - 1) + 1. \tag{13}$$

(2) If $X = \mathrm{Bin}(n,p)$, then $\mathbb{P}[X = k] = \binom{n}{k} p^k (1-p)^{n-k}$, and

$$\phi_X(u) = (p(e^{iu} - 1) + 1)^n. \tag{14}$$

(3) If $X = \mathrm{Pois}(\lambda)$, then $\mathbb{P}[X = k] = e^{-\lambda}\lambda^k/k!$, and

$$\phi_X(u) = \sum_{k=0}^{\infty} \frac{e^{-\lambda}\lambda^k}{k!} e^{iuk} = e^{-\lambda} \sum_{k=0}^{\infty} \frac{(\lambda e^{iu})^k}{k!} = \exp\left(\lambda(e^{it} - 1)\right). \tag{15}$$

(4) If X is a Gaussian random variable $\mathscr{N}(m, \sigma^2)$, then

$$\phi_X(u) = \exp\left(ium - \frac{u^2\sigma^2}{2}\right). \tag{16}$$

The importance of the characteristics functions could be summarized in the following two theorems.

Theorem 1.5. If two random variables have the same characteristic function, then they have the same distribution. Of course, the converse also holds, in that if two random variables have the same distribution, then they have the same characteristic function.

Theorem 1.6 (Paul Lévy theorem)**.** Let X_n be a sequence of random variables on $\mathbb{R}$. Then

(1) If X_n converges to X in distribution, then $\phi_{X_n}(u) \to \phi_X(u)$ as $n \to \infty$ for every u, and in fact this convergence is uniform.

(2) If $\phi_{X_n}(u) \to \phi_X(u)$ for every u pointwise, then X_n converges to X in distribution.

(3) If $\phi_{X_n}(u)$ converges to some function $\phi(u)$ for every u pointwise, and $\phi(u)$ is continuous at 0, then there exists a unique random variable X, such that ϕ is the characteristic function of X and X_n converges to X in distribution.

We use Part (2) of Paul Lévy theorem to prove that $B^{(n)}(t)$ converges to the normal distribution with expectation 0 and variance t. We have that

$$\phi_{B^{(n)}(t)}(u) = \mathbb{E}\left[\exp\left(iuB^{(n)}(t)\right)\right] = \mathbb{E}\left[\exp\left(\frac{iuM_{nt}}{\sqrt{n}}\right)\right] = \mathbb{E}\left[\exp\left(\frac{iu}{\sqrt{n}}\sum_{j=1}^{nt} X_j\right)\right]$$
$$= \left(\mathbb{E}\left[\exp\left(\frac{iuX_j}{\sqrt{n}}\right)\right]\right)^{nt} = \left(\frac{1}{2}\exp\left(\frac{iu}{\sqrt{n}}\right) + \frac{1}{2}\exp\left(\frac{-iu}{\sqrt{n}}\right)\right)^{nt}. \tag{17}$$

Now we need to show that as $n \to \infty$, the above result converges to the characteristic function of the normal random variable with mean 0 and variance t, *i.e.*, to $e^{-u^2t/2}$. Expanding in Taylor series, we can see that

$$\frac{1}{2}\exp\left(\frac{iu}{\sqrt{n}}\right) + \frac{1}{2}\exp\left(\frac{-iu}{\sqrt{n}}\right) = 1 - \frac{u^2}{2n} + O(n^{-3/2}), \tag{18}$$

and thus

$$\phi_n(u) = \left(1 - \frac{u^2}{2n} + O(n^{-3/2})\right)^{nt} \to \exp\left(-\frac{u^2}{2n}\cdot nt\right) = \exp\left(-\frac{u^2t}{2}\right). \tag{19}$$

Therefore, we can conclude that $B^{(n)}(t)$ converges in distribution to a normal variable with expectation 0 and variance t. Strictly speaking, we have just demonstrated that for a fixed moment t, the distribution of $B^{(n)}(t)$ converges to $\mathscr{N}(0,t)$, but we did not prove the convergence of the whole stochastic process to Brownian motion. To do this, one has to further prove the convergence of the finite dimensional distributions and use *Prokhorov's theorem*.

1.4 Brownian Motion: A Formal Definition

A one-dimensional Brownian motion is a continuous-time stochastic process that satisfies the following properties.

- **Independence of increments:** If $t_0 < t_1 < \cdots < t_n$, then the random variables
$$B(t_0), B(t_1) - B(t_0), \cdots, B(t_n) - B(t_{n-1}) \tag{20}$$
are independent.
- **Normally distributed increments:** If $s, t \geq 0$, then
$$\mathbb{P}[B(t+s) - B(s) \in A] = \int_A \frac{1}{\sqrt{2\pi t}} e^{-x^2/2t} dx. \tag{21}$$
- **Continuous trajectories:** With probability 1, $B_0 = 0$ and $t \mapsto B_t$ is continuous.

2 Properties of the Brownian Motion

The limiting object described above is called a *standard Brownian motion*. It will be our main object of study, and the main building block in many applications.

2.1 Covariance of Brownian Motion

Let $0 \leq s \leq t$, then

$$\mathbb{E}[B(s)B(t)] = \mathbb{E}[B(s)(B(t) - B(s) + B(s))] = \mathbb{E}[B(s)(B(t) - B(s))] + \mathbb{E}[B^2(s)]. \tag{22}$$

By independence of increments, we have that $\mathbb{E}[B(s)(B(t)-B(s))] = \mathbb{E}[B(s)]\mathbb{E}[B(t)-B(s)] = 0$. Moreover, from the fact that $B(s)$is a normally distributed random variable with expectation 0 and variance s, we have that $\mathbb{E}[B^2(s)] = s$. Bringing these results into the previous expression, we have that

$$\mathbb{E}[B(t)B(s)] = 0 + s = s. \tag{23}$$

Note that it is important to know how to compute the moments of a normal random variable.

2.2 Non-differentiability of Brownian Paths

In the following theorem we prove that the trajectories of Brownian motion are very special, in that they are non-differentiable at every point.

Theorem 2.1. With probability 1, Brownian paths are not differentiable at any point.

Proof. Let $W = \{\omega;\ \exists t \in (0,1)$, such that $B(t)$ is differentiable at $t\}$. We want to show that $\mathbb{P}[W] = 0$. We shall use the following lemma, whose proof is fairly straightforwards and will be left as an exercise.

> *Lemma. If $f : (0,1) \to \mathbb{R}$ is differentiable at $x \in (0,1)$, then there exists $C > 0$ and $\delta > 0$, such that $f(x) - f(s) \le C|x - s|$, for any $s \in [x - \delta, x + \delta]$.*

Now let $M_n(k) = \max_i \{|B((i+1)/n) - B(i/n)|;\ i = k-1, k, k+1\}$, and $M_n = \min_k \{M_n(k);\ k = 1, \cdots, n\}$. By the above lemma, for any $\omega \in W$, we can find $C > 0$ and $\delta > 0$, such that

$$|B(t) - B(s)| \le C|t - s|, \qquad \forall s \in [t - \delta, t + \delta]. \tag{24}$$

Choose n sufficiently large such that $n > \max\{t, 4/\delta\}$. We also choose k such that $(k-1)/n \le t \le k/n$. The choices of k and n imply that $|i/n - t| < \delta$ for $i = k-1, k, k+1, k+2$. Furthermore, we can see that

$$\left|B\left(\frac{i+1}{n}\right) - B\left(\frac{i}{n}\right)\right| \le \left|B\left(\frac{i+1}{n}\right) - B(t)\right| + \left|B\left(\frac{i}{n}\right) - B(t)\right| \le C\left|\frac{i+1}{n} - t\right| + C\left|\frac{i}{n} - t\right| < 2C\delta < \frac{8C}{n}. \tag{25}$$

Note that $B((i+1)/n) - B(i/n)$ are independent and normally distributed with mean 0 and variance $1/n$, so that

$$\mathbb{P}\left[|M_n| \le \frac{8C}{n}\right] \le n\mathbb{P}\left[|M_n(k)| \le \frac{8C}{n}\right] = n\left(\mathbb{P}\left[|z| \le \frac{8C}{\sqrt{n}}\right]\right)^3 \le n\left(\frac{16C}{\sqrt{2\pi n}}\right)^3 = \frac{16^3C^3}{\sqrt{2\pi n}}, \tag{26}$$

where z is the standard normal. Let $A_n := \{|M_n| \le 8C/n\}$, then the above inequality means that $\lim_{n\to\infty} \mathbb{P}[A_n] = 0$. By Fatou's lemma, we can conclude that

$$\mathbb{P}\left[\liminf_{n\to\infty} A_n\right] \le \liminf_{n\to\infty} \mathbb{P}[A_n] = \lim_{n\to\infty} \mathbb{P}[A_n] = 0. \tag{27}$$

Finally, we note that $W \subseteq \liminf_{n\to\infty} A_n$, so $\mathbb{P}[W] = 0$, and the proof is complete. □

2.3 Scaling Properties of the Brownian Motion

If we fix time t, then the distribution of Brownian motion $B(t)$ is normal with expectation 0 and variance t. In fact, for any fixed set of times $t_1 < t_2 < \cdots < t_n$, the random variables $B(t_1), B(t_2), \cdots, B(t_n)$ are jointly normal with mean 0 and covariance given by $\mathbb{E}[B(s)B(t)] = \min\{s, t\}$.

Theorem 2.2. If $B(0) = 0$, then for any $\lambda > 0$, the stochastic process $B(\lambda t)/\sqrt{\lambda}$, $t \ge 0$ is a Brownian motion.

Proof. First, note that $X(t) = B(\lambda t)/\sqrt{\lambda}$ is a Gaussian process, *i.e.*, for any set $t_1 < t_2 < \cdots < t_n$, the joint distribution of $X(t_1), X(t_2), \cdots, X(t_n)$ is a multivariate Gaussian distribution. This property is clearly inherited from Brownian motion properties. Since normal distribution is characterized by its mean and covariance, we have to check that the mean and covariance of the process $X(t)$ coincide with those of Brownian motion. Indeed,

$$\mathbb{E}[X(t)] = \mathbb{E}\left[\frac{B(\lambda t)}{\sqrt{\lambda}}\right] = \frac{\mathbb{E}[B(\lambda t)]}{\sqrt{\lambda}} = 0, \tag{28}$$

and for $s < t$, we have that

$$\mathbb{E}[X(s)X(t)] = \mathbb{E}\left[\frac{B(\lambda s)}{\sqrt{\lambda}} \cdot \frac{B(\lambda t)}{\sqrt{\lambda}}\right] = \frac{\mathbb{E}[B(\lambda s)B(\lambda t)]}{\lambda} = \frac{\min\{\lambda s, \lambda t\}}{\lambda} = s. \tag{29}$$

Therefore, we can conclude that $B(\lambda t)/\sqrt{\lambda}$ has the same distribution as a Brownian motion, and thus a Brownian motion. The proof is thus complete. □

Theorem 2.3. If $B(t)$ is a Brownian motion starting at 0, then so is the process defined by $X(0) = 0$ and $X(t) = tB(1/t)$ for $t > 0$.

Proof. Fix $t_1 < t_2 < \cdots < t_n$, then clearly $X(t_1), X(t_2), \cdots, X(t_n)$ has a multivariate Gaussian distribution. We just have to check that it has the same mean and covariance structure. First of all, we have that

$$\mathbb{E}[X(t)] = \mathbb{E}[tB(1/t)] = t\mathbb{E}[B(1/t)] = 0. \tag{30}$$

Moreover, for $0 < s < t$, we can deduce that

$$\mathbb{E}[X(s)X(t)] = \mathbb{E}[sB(1/s) \cdot tB(1/t)] = st\mathbb{E}[B(1/s)B(1/t)] = st \min\{1/s, 1/t\} = s. \tag{31}$$

Therefore, $X(t)$ is indeed a Brownian motion, so the proof is complete. □

2.4 Quadratic Variation of Brownian Motion

Last time we computed quadratic variation of scaled SRW and it turned out to be t. In the following theorem we prove that the quadratic variation of the Brownian motion is also t. Let me again emphasize that the paths of Brownian motion are unusual in that their quadratic variation is not zero. This makes stochastic calculus different from ordinary calculus. In fact, non-zero quadratic variation is the source of the *volatility* term in the *Black-Scholes equation*.

To see how Brownian motion is different from functions we are used to in ordinary calculus, consider a function $f : \mathbb{R} \to \mathbb{R}$ that is continuously differentiable and compute its quadratic variation up to time T. Let us introduce the norm of the partition $0 = t_0 < t_1 < \cdots < t_n = T$, such that

$$\|\Pi\| = \max_{0 \le j \le n-1} (t_{j+1} - t_j). \tag{32}$$

Then, its quadratic variation can be computed as

$$\langle f, f \rangle_T = \lim_{\|\Pi\| \to 0} \sum_{j=0}^{n-1} (f(t_{j+1}) - f(t_j))^2. \tag{33}$$

We will use the mean value theorem which says that in each interval (x_1, x_2), there exists a point x_0, such that

$$f(x_2) - f(x_1) = f'(x_0)(x_2 - x_1). \tag{34}$$

Applying to each interval (t_j, t_{j+1}), we get that there is a point t_j^* in each such interval, such that

$$f(t_{j+1}) - f(t_j) = f'(t_j^*)(t_{j+1} - t_j). \tag{35}$$

Therefore, we can compute that

$$\begin{aligned} \sum_{j=0}^{n-1} (f(t_{j+1}) - f(t_j))^2 &= \sum_{j=0}^{n-1} |f'(t_j^*)|^2 (t_{j+1} - t_j)^2 \le \max_{0 \le j \le n-1} |t_{j+1} - t_j| \sum_{j=0}^{n-1} |f'(t_j^*)|^2 (t_{j+1} - t_j) \\ &= \|\Pi\| \sum_{j=0}^{n-1} |f'(t_j^*)|^2 (t_{j+1} - t_j) \to \|\Pi\| \int_0^T |f'(t)|^2 dt \to 0, \qquad \text{as } \|\Pi\| \to 0. \end{aligned} \tag{36}$$

Therefore, we have shown that continuously differentiable function has zero quadratic variation. For this reason we rarely consider quadratic variation in ordinary calculus.

Theorem 2.4. Let $B(t)$ be a Brownian motion. Then $\langle B, B \rangle_T = T$ almost surely.

Proof. Define

$$Q_\Pi = \sum_{j=0}^{n-1} (B(t_{j+1}) - B(t_j))^2. \tag{37}$$

We shall prove that $\mathbb{E}[Q_\Pi] \to T$ and $\mathrm{Var}[Q_\Pi] \to 0$ as $\|\Pi\| \to 0$. In other words, we shall prove the L^2 convergence of Q_Π, which is stronger than *almost sure* convergence. First, notice that

$$\mathbb{E}[(B(t_{j+1}) - B(t_j))^2] = \mathrm{Var}[B(t_{j+1}) - B(t_j)] = t_{j+1} - t_j. \tag{38}$$

□

Therefore, we can compute that

$$\mathbb{E}[Q_\Pi] = \sum_{j=0}^{n-1} \mathbb{E}[(B(t_{j+1}) - B(t_j))^2] = \sum_{j=0}^{n-1} (t_{j+1} - t_j) = T. \tag{39}$$

Now, in order to show that $\mathrm{Var}[Q_\Pi] \to 0$ as $\|\Pi\| \to 0$, note that

$$\begin{aligned}
\mathrm{Var}[(B(t_{j+1}) - B(t_j))^2] &= \mathbb{E}[((B(t_{j+1}) - B(t_j))^2 - (t_{j+1} - t_j))^2] \\
&= \mathbb{E}[(B(t_{j+1}) - B(t_j))^4 - 2(t_{j+1} - t_j)(B(t_{j+1}) - B(t_j))^2 + (t_{j+1} - t_j)^2] \\
&= \mathbb{E}[(B(t_{j+1}) - B(t_j))^4] - 2(t_{j+1} - t_j)\mathbb{E}[(B(t_{j+1}) - B(t_j))^2] + (t_{j+1} - t_j)^2 \\
&= 3(t_{j+1} - t_j)^2 - 2(t_{j+1} - t_j)^2 + (t_{j+1} - t_j)^2 = 2(t_{j+1} - t_j)^2.
\end{aligned} \tag{40}$$

In the previous deduction, we used the fourth moment of a normal random variable. Now, we can see that

$$\mathrm{Var}[Q_\Pi] = \sum_{j=0}^{n-1} \mathrm{Var}[(B(t_{j+1}) - B(t_j))^2] = \sum_{j=0}^{n-1} 2(t_{j+1} - t_j)^2 \le 2\,\|\Pi\| \sum_{j=0}^{n-1} (t_{j+1} - t_j) = 2\,\|\Pi\|\, T, \tag{41}$$

which clearly tends to 0 as $\|\Pi\| \to 0$. Therefore, we can conclude that $\langle B, B\rangle_T = \lim_{\|\Pi\|\to 0} Q_\Pi$ which has expectation T and variance 0, thus being T almost surely. Our proof is thus complete.

Remark 2.5. $\langle B, B\rangle_t = t$ can be informally written as

$$dB(t)dB(t) = dt. \tag{42}$$

Remark 2.6. In addition to computing the quadratic variation of Brownian motion, we can also compute the cross variation of $B(t)$ with t and the quadratic variation of t itself, which are

$$\lim_{\|\Pi\|\to 0} \sum_{j=0}^{n-1} (B(t_{j+1}) - B(t_j))(t_{j+1} - t_j) = 0, \qquad \lim_{\|\Pi\|\to 0} \sum_{j=0}^{n-1} (t_{j+1} - t_j)^2 = 0. \tag{43}$$

These can be informally written as

$$dB(t)dt = 0, \qquad dtdt = 0. \tag{44}$$