



Machine Learning


# Linear regression with one variable

---

## Model representation

# Housing Prices (Portland, OR)

Price  
(in 1000s  
of dollars)



## Supervised Learning

Given the “right answer” for each example in the data.

## Regression Problem

Predict real-valued output

Training set of housing prices (Portland, OR)	Size in feet <sup>2</sup> (x)	Price (\$) in 1000's (y)
	2104	460
	1416	232
	1534	315
	852	178
	...	...

Notation:

**m** = Number of training examples

**x**'s = "input" variable / features

**y**'s = "output" variable / "target" variable

Training Set



Learning Algorithm



Size of  
house



$h$




Estimated  
price

**How do we represent  $h$  ?**

Linear regression with one variable.  
Univariate linear regression.





Machine Learning

Linear regression  
with one variable

---

Cost function


Training Set	Size in feet <sup>2</sup> (x)	Price (\$) in 1000's (y)
	2104	460
	1416	232
	1534	315
	852	178
	...	...

Hypothesis:  $h_{\theta}(x) = \theta_0 + \theta_1 x$


$\theta_i$ 's: Parameters

How to choose  $\theta_i$ 's ?


$$h_{\theta}(x) = \theta_0 + \theta_1 x$$



$$\theta_0 = 1.5$$
$$\theta_1 = 0$$




$$\theta_0 = 0$$
$$\theta_1 = 0.5$$




$$\theta_0 = 1$$
$$\theta_1 = 0.5$$





Idea: Choose  $\theta_0, \theta_1$  so that  $h_{\theta}(x)$  is close to  $y$  for our training examples  $(x, y)$





Machine Learning

Linear regression  
with one variable

---

Cost function  
intuition I

## Simplified

Hypothesis:

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

Parameters:

$$\theta_0, \theta_1$$

Cost Function:

$$J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

Goal: minimize  $J(\theta_0, \theta_1)$   
 $\theta_0, \theta_1$

$$h_{\theta}(x) = \theta_1 x$$


$$\theta_1$$

$$J(\theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

minimize  $J(\theta_1)$   
 $\theta_1$


$$h_{\theta}(x)$$

(for fixed  $\theta_1$ , this is a function of  $x$ )




$$J(\theta_1)$$

(function of the parameter  $\theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_1$ , this is a function of  $x$ )




$$J(\theta_1)$$

(function of the parameter  $\theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_1$ , this is a function of  $x$ )



$$J(\theta_1)$$

(function of the parameter  $\theta_1$ )









Machine Learning

Linear regression  
with one variable

---

Cost function  
intuition II

Hypothesis:  $h_{\theta}(x) = \theta_0 + \theta_1 x$


Parameters:  $\theta_0, \theta_1$

Cost Function:  $J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$

Goal: minimize  $J(\theta_0, \theta_1)$   
 $\theta_0, \theta_1$

$$h_{\theta}(x)$$


(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )



$$h_{\theta}(x) = 50 + 0.06x$$


$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )




$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )




$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )




$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )




$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )









Machine Learning

Linear regression  
with one variable

---


Gradient  
descent


Have some function  $J(\theta_0, \theta_1)$

Want  $\min_{\theta_0, \theta_1} J(\theta_0, \theta_1)$

## Outline:

- Start with some  $\theta_0, \theta_1$
- Keep changing  $\theta_0, \theta_1$  to reduce  $J(\theta_0, \theta_1)$   
until we hopefully end up at a minimum





# Gradient descent algorithm

repeat until convergence {  
     $\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$     (for  $j = 0$  and  $j = 1$ )  
}

---


Correct: Simultaneous update

```
temp0 :=  $\theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$   
temp1 :=  $\theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$   
 $\theta_0 :=$  temp0  
 $\theta_1 :=$  temp1
```

Incorrect:

```
temp0 :=  $\theta_0 - \alpha \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1)$   
 $\theta_0 :=$  temp0  
temp1 :=  $\theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1)$   
 $\theta_1 :=$  temp1
```





Machine Learning

Linear regression  
with one variable

---

Gradient descent  
intuition



# Gradient descent algorithm


repeat until convergence {  
     $\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$       (simultaneously update  
     $j = 0$  and  $j = 1$ )  
}




$$\theta_1 := \theta_1 - \alpha \frac{\partial}{\partial \theta_1} J(\theta_1)$$

If  $\alpha$  is too small, gradient descent can be slow.

If  $\alpha$  is too large, gradient descent can overshoot the minimum. It may fail to converge, or even diverge.






Current value of  $\theta_1$

$$\theta_1 := \theta_1 - \alpha \frac{d}{d\theta_1} J(\theta_1)$$


Gradient descent can converge to a local minimum, even with the learning rate  $\alpha$  fixed.

$$\theta_1 := \theta_1 - \alpha \frac{d}{d\theta_1} J(\theta_1)$$

As we approach a local minimum, gradient descent will automatically take smaller steps. So, no need to decrease  $\alpha$  over time.







Machine Learning

# Linear regression with one variable

---

## Gradient descent for linear regression

## Gradient descent algorithm

repeat until convergence {  
     $\theta_j := \theta_j - \alpha \frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1)$   
    (for  $j = 1$  and  $j = 0$ )  
}

## Linear Regression Model

$$h_{\theta}(x) = \theta_0 + \theta_1 x$$

$$J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$



$$\frac{\partial}{\partial \theta_j} J(\theta_0, \theta_1) =$$

$$j = 0 : \frac{\partial}{\partial \theta_0} J(\theta_0, \theta_1) =$$

$$j = 1 : \frac{\partial}{\partial \theta_1} J(\theta_0, \theta_1) =$$

# Gradient descent algorithm


repeat until convergence {


$$\theta_0 := \theta_0 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})$$

$$\theta_1 := \theta_1 - \alpha \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) \cdot x^{(i)}$$

}


update  
 $\theta_0$  and  $\theta_1$   
simultaneously






$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )




$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )




$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )




$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )



$$J(\theta_0, \theta_1)$$


(function of the parameters  $\theta_0, \theta_1$ )






$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )




$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )




$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )




$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )




$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )




$$h_{\theta}(x)$$

(for fixed  $\theta_0, \theta_1$ , this is a function of  $x$ )



$$J(\theta_0, \theta_1)$$

(function of the parameters  $\theta_0, \theta_1$ )



# **“Batch” Gradient Descent**

“Batch”: Each step of gradient descent uses all the training examples.

