Variable Name	Variable Description	Variable Attribute
Dataset 1: **Pandemic** (pandemicPRL.csv)		
study_id	a unique id to help users match behavioral data to raw task data* *See Erin et.al., 2020 for the first 72 subjects' raw task data	e.g., TP117
mturk_id	a unique, alphanumeric identifier for each MTurk worker	e.g., A2541C8MY0BYV3
month	month group based on *date_collected* *the present study analyzed data collected from 3/19/2020 - 7/17/2020	**Group** / **Date Range** 0 / 2017-2018 1 / 1/28-1/30 2 / 3/19-4/1 3 / 4/3-4/5 4 / 6/2-6/27 5 / 7/1-7/17
date_collected	date of data collection	MM/DD/YYYY
dataset	grouping of dataset	**Group** / **Dataset** elife2020 / data collected in Erin et. al., 2020 pandemic / data collected during the pandemic replication / data collected to replicate effects
period	grouping with respect to lockdown *the present study analyzed data collected during the lockdown and postlockdown periods	**Group** / **Period** prelockdown / before the lockdown lockdown / during the lockdown postlockdown / after the lockdown

<table>
<tr><td>demo_1</td><td>gender</td><td><table><tr><th>Group</th><th>Gender</th></tr><tr><td>1</td><td>Male</td></tr><tr><td>2</td><td>Female</td></tr><tr><td>3</td><td>Other</td></tr><tr><td>N/A</td><td>Not specified</td></tr></table></td></tr>
<tr><td>demo_2</td><td>age</td><td></td></tr>
<tr><td>demo_3</td><td>Hispanic</td><td><table><tr><th>Group</th><th>Hispanic</th></tr><tr><td>1</td><td>No</td></tr><tr><td>2</td><td>Yes</td></tr><tr><td>N/A</td><td>Not specified</td></tr></table></td></tr>
<tr><td>demo_4</td><td>race</td><td><table><tr><th>Group</th><th>Race</th></tr><tr><td>1</td><td>White</td></tr><tr><td>2</td><td>Black or African American</td></tr><tr><td>3</td><td>Asian</td></tr><tr><td>4</td><td>American Indian or Alaska Native</td></tr><tr><td>5</td><td>Multiracial</td></tr><tr><td>N/A</td><td>Not specified</td></tr></table></td></tr>
<tr><td>demo_5</td><td>education</td><td><table><tr><th>Group</th><th>Education</th></tr><tr><td>1</td><td>Less than a high school diploma</td></tr><tr><td>2</td><td>High school degree or equivalent (e.g., GED)</td></tr></table></td></tr>
</table>

		3 — Some college or university, no degree 4 — Associate degree (e.g., AA, AS) 5 — Bachelor's degree (e.g., BA, BS) 6 — Master's degree (e.g., MA, MS, MEd) 7 — Completed some postgraduate 8 — Doctorate or professional degree (e.g., PhD, MD, JD) N/A — Not specified
demo_6	employment	**Group** — **Employment** 1 — Employed full time (40 or more hours per week) 2 — Employed part time (up to 39 hours per week) 3 — Unemployed and currently looking for work 4 — Unemployed and not currently looking for work 5 — Student 6 — Retired 7 — Homemaker 8 — Self-employed 9 — Unable to work N/A — Not specified
demo_7	income	**Group** — **Income** 1 — Less than $20,000

<table>
<tr><td></td><td></td><td>
<table>
<tr><td>2</td><td>$20,000 to $34,999</td></tr>
<tr><td>3</td><td>$35,000 to $49,999</td></tr>
<tr><td>4</td><td>$50,000 to $74,999</td></tr>
<tr><td>5</td><td>$75,000 to $99,999</td></tr>
<tr><td>6</td><td>Over $100,000</td></tr>
<tr><td>N/A</td><td>Not specified</td></tr>
</table>
</td></tr>
<tr><td>demo_8</td><td>Are you currently taking any prescription medications, over-the-counter medications, or supplements? (self-report medication; drug)</td><td>
<table>
<tr><th>Group</th><th>Drug</th></tr>
<tr><td>0</td><td>No medication</td></tr>
<tr><td>1</td><td>Medicated – other (possibly psychoactive)</td></tr>
<tr><td>2</td><td>Medicated – psychiatric</td></tr>
<tr><td>N/A</td><td>Not specified</td></tr>
</table>
</td></tr>
<tr><td>demo_9</td><td>Have you ever been diagnosed with any neurological or psychiatric conditions (e.g., concussion/head trauma, epilepsy, depression, anxiety, schizophrenia? (self-report mental health; disorder)</td><td>
<table>
<tr><th>Group</th><th>Disorder</th></tr>
<tr><td>0</td><td>No psychiatric diagnosis</td></tr>
<tr><td>1</td><td>Neurological</td></tr>
<tr><td>2</td><td>Mood disorder</td></tr>
<tr><td>3</td><td>Schizophrenia spectrum</td></tr>
<tr><td>N/A</td><td>Not specified</td></tr>
</table>
</td></tr>
<tr><td>demo_10</td><td>living situation</td><td>
<table>
<tr><th>Group</th><th>Living Situation</th></tr>
<tr><td>1</td><td>Lockdown not leaving home</td></tr>
<tr><td>2</td><td>Staying at home, but only allowed out for essential things</td></tr>
<tr><td>3</td><td>Staying mostly at home, but can go out</td></tr>
<tr><td>4</td><td>Going to work and going out,</td></tr>
</table>
</td></tr>
</table>

<table>
<tr><td></td><td></td><td><table><tr><td></td><td>but with social distancing</td></tr><tr><td>5</td><td>Life largely unchanged</td></tr><tr><td>N/A</td><td>Not specified</td></tr></table></td></tr>
<tr><td>bai_</td><td>21-item beck anxiety inventory*

*See github repo (…/questionnaires/bai)</td><td><table><tr><th>Group</th><th>BAI</th></tr><tr><td>0</td><td>Not at all</td></tr><tr><td>1</td><td>Mildly but it didn’t bother me much</td></tr><tr><td>2</td><td>Moderately – it wasn’t pleasant at times</td></tr><tr><td>3</td><td>Severely – it bothered me a lot</td></tr><tr><td>N/A</td><td>Not specified</td></tr></table></td></tr>
<tr><td>bdi_</td><td>21-item beck depression inventory*

NOTE: avoided asking question 9 to participants as it related to suicide</td><td>*See github repo (…/questionnaires/bdi)</td></tr>
<tr><td>rgpts_</td><td>18-item revised green et. al paranoid thought scale, where the first 8-items assess social reference paranoia (rgpts_ref) and the last 10-items assess persecutory ideations (rgpts_per)*</td><td><table><tr><th>Group</th><th>R-GPTS</th></tr><tr><td>0</td><td>Not at all</td></tr><tr><td>1</td><td>-</td></tr><tr><td>2</td><td>-</td></tr><tr><td>3</td><td>-</td></tr><tr><td>4</td><td>Totally</td></tr><tr><td>N/A</td><td>Not specified</td></tr></table>
*See github repo (…/questionnaires/rgpts)</td></tr>
<tr><td>scid_group</td><td>paranoia grouping* based on the Structured Clinical Interview for DSM-IV Axis II Disorder (SCID-II)

*See Reed et.al, 2020</td><td><table><tr><th>Group</th><th>SCID</th></tr><tr><td>low</td><td>below clinical cutoff on SCID-II paranoia</td></tr><tr><td>high</td><td>above clinical cutoff on SCID-II paranoia</td></tr></table></td></tr>
</table>

docs_	20-item dimensional obsessive compulsive scale*	*See github repo (…/questionnaires/docs)
avg_rt_total	avg_rt_total represents averaged reaction time of the task	avg_rt_total is refered to as “decision time”
reversals_	number of reversals achieved in the first half (reversals_block1) and second half (reversals_block2) of the task	
wsr_	win-switch rate behavior in the first half (wsr_block1) and second half (wsr_block2) of the task	wsr_mean is the average of wsr_block1 and wsr_block2
lsr_	lose-stay rate behavior in the first half (lsr_block1) and second half (lsr_block2) of the task	lsr_mean is the average of lsr_block1 and lsr_block2
mu02_	initial prior of the second-level contingency belief in the first half (mu02_1) and second half (mu02_2) of the task	μ_2^0 is the average of mu02_block1 and mu02_block2
mu03_	initial prior of the third-level volatility belief in the first half (mu03_1) and second half (mu03_2) of the task	μ_3^0 is the average of mu03_block1 and mu03_block2
kappa2_	sensitivity measure of unexpected changes in the first half (kappa2_1) and second half (kappa2_2) of the task	κ is the average of kappa2_block1 and kappa2_block2
omega2_	evolution rate of the second-level reward contingency belief in the first half (omega2_1) and second half (omega2_2) of the task	ω_2^0 is the average of omega2_block1 and omega2_block2
omega3_	evolution rate of the third-level volatility belief in the first half (omega3_1) and second half (omega3_2) of the task	ω_3^0 is the average of omega3_block1 and omega3_block2