

A sharp winding bound for the self-similar collapse of three point vortices in the α -models

Chase Hendrick

Independent Researcher

chasewhendrick@gmail.com

Abstract

In the α -models of two-dimensional flow a point vortex of circulation Γ induces the velocity $\Gamma r^{-\alpha-1}/(2\pi)$ at distance r ; $\alpha = 0$ is the Euler equation and $\alpha = 1$ the surface quasi-geostrophic equation. In a self-similar collapse of three point vortices every vortex moves on a logarithmic spiral about the collision point, and the rotation per unit decrease of the logarithm of the squared size, $P = |\operatorname{Im} \kappa|/(2|\operatorname{Re} \kappa|)$, measures how tightly the spiral winds. We prove that $P > \sqrt{3} + \alpha/(2 + \alpha)$ for every self-similar collapse of three point vortices for every $\alpha > -1$, that is, whenever the velocity induced by a vortex decays with distance, and that the constant is sharp: it is approached, but not attained, as two weak vortices of opposite sign merge next to a strong one. Equivalently, every vortex travels more than $(4 + \alpha)/(2 + \alpha)$ times its initial distance from the collision point, and its path makes an angle larger than $\arccos((2 + \alpha)/(4 + \alpha))$ with the direction to that point. The bound decreases from $\sqrt{3}/2$ in the Euler case to $2/3$ in the surface quasi-geostrophic case and tends to 0 as $\alpha \rightarrow \infty$. The proof rests on two elementary identities: the circulations of a collapsing triangle are fixed, up to a factor, by its side lengths, and P is a symmetric function of the side lengths alone. For $\alpha = 0$ it gives a short proof of the Euler bound $\sqrt{3}/2$. For four or more vortices in the Euler case the bound fails: numerically, four vortices reach $P = 0.7978967838\dots$ and sixty-one vortices $P = 0.4981844392\dots$

Keywords: point vortices; vortex collapse; self-similar motion; surface quasi-geostrophic equation; generalized Euler equation.

MSC 2020: 76B47, 76U60, 37N10.

1 Introduction

Three point vortices whose angular impulse and harmonic combination of circulations vanish can shrink to a point in finite time while the triangle they form keeps its shape and rotates. Each vortex then moves on a logarithmic spiral about the collision point. In the Euler case this self-similar collapse goes back to Gröbli; Conte and de Seze [3] and Kimura [11] wrote the motion in closed form, and the companion preprint [8] shows that the winding of the spiral, measured by the dimensionless number P of (3) below, satisfies $P > \sqrt{3}/2$ for every self-similar collapse of three vortices, with a sharp constant.

Self-similar collapse of three vortices also occurs in the family of active-scalar models in which the streamfunction is obtained from the vorticity by a fractional inverse Laplacian, the generalized Euler or α -models, which include the surface quasi-geostrophic (SQG) equation [1, 7, 13, 14]. Their point vortices move by

$$\frac{dz_j}{dt} = \frac{i}{2\pi} \sum_{k \neq j} \Gamma_k \frac{z_j - z_k}{|z_j - z_k|^{2\beta}}, \quad \beta = 1 + \frac{\alpha}{2}, \quad (1)$$

with positions $z_j \in \mathbb{C}$ and real circulations Γ_j . The case $\alpha = 0$ ($\beta = 1$) is the Euler system, and $\alpha = 1$ ($\beta = 3/2$) is SQG. A constant factor in front of the sum, which differs between papers, only rescales time and does not change P . In the conventions of Badin and Barry [1] and of Grotto and Pappalettera [7] the same system has the parameter $2 - \alpha$.

This note extends the winding bound to these models. With $B(\alpha) = \sqrt{3 + \alpha}/(2 + \alpha)$ we prove:

Theorem 1. *Let $\alpha > -1$. Every self-similar collapse of three point vortices under (1) has $P > B(\alpha)$, and for every $\delta > 0$ there is one with $P < B(\alpha) + \delta$.*

The proof (Section 4) needs two facts proved in Section 3: a collapsing triangle is scalene and carries circulations fixed up to a factor by its side lengths (Lemma 2), and P is a symmetric function of the side lengths alone (Lemma 3). For $\alpha = 0$ it gives a short proof of the Euler bound of [8], though not the minimum for each ratio of circulations obtained there. Section 5 records that the bound does not extend to more than three vortices in the Euler case, and Section 6 lists the numerical checks.

2 Self-similar motion

Call a motion of (1) *self-similar* if the triangle keeps its shape and orientation, that is, $z_j(t) = a(t) + b(t)z_j(0)$ with complex $a(t)$ and $b(t) \neq 0$, and *collapsing* if moreover $b(t) \rightarrow 0$ in finite time.

Lemma 1. *Let a self-similar motion of (1) change size. Then $\Gamma_1 + \Gamma_2 + \Gamma_3 \neq 0$, and with $z_c = \sum_j \Gamma_j z_j / \sum_j \Gamma_j$ the center of vorticity, there is $\kappa_0 \in \mathbb{C}$ with $\text{Re } \kappa_0 \neq 0$ such that $\dot{z}_j = \kappa(t)(z_j - z_c)$ with $\kappa(t) = \kappa_0 \lambda^{-2\beta}$, and*

$$z_j(t) - z_c = \lambda(t)e^{i\phi(t)}(z_j(0) - z_c), \quad \lambda^{2\beta} = 1 + 2\beta \text{Re } \kappa_0 t, \quad \phi = \frac{\text{Im } \kappa_0}{2 \text{Re } \kappa_0} \ln \lambda^2. \quad (2)$$

The motion collapses if and only if $\text{Re } \kappa_0 < 0$, at $t_c = -1/(2\beta \text{Re } \kappa_0)$. Every vortex moves on a logarithmic spiral about z_c ; the number

$$P = \frac{|\text{Im } \kappa|}{2|\text{Re } \kappa|} \quad (3)$$

does not depend on t , and it is the angle through which the triangle turns while its squared size decreases by the factor e . The path of each vortex makes the angle $\arctan 2P$ with the direction to z_c , a vortex at distance r_0 from z_c travels the distance $r_0\sqrt{1 + 4P^2}$ before the collapse, and the initial angular velocity $\omega_0 = \text{Im } \kappa_0$ satisfies $|\omega_0|t_c = P/\beta$.

Proof. The linear impulse $Q = \sum_j \Gamma_j z_j$ is conserved. If $\sum_j \Gamma_j = 0$, then $Q(t) = b(t)Q(0)$, and since $|b|$ changes, $Q(0) = 0$; then $\Gamma_1(z_1 - z_3) + \Gamma_2(z_2 - z_3) = 0$ with real coefficients not both zero, so the vortices are collinear, and a collinear triangle does not change size under (1) (Lemma 2 below). Hence $\sum_j \Gamma_j \neq 0$, z_c is conserved, and $z_c = a + bz_c$, so $z_j(t) - z_c = b(t)w_j$ with $w_j = z_j(0) - z_c$. The right side of (1) is homogeneous: replacing $z_j - z_c$ by bw_j multiplies it by $b|b|^{-2\beta}$. Writing $\dot{z}_j(0) = \kappa_0 w_j$, which holds for the same κ_0 for all j because $\dot{b}(0)w_j = \dot{z}_j(0)$, we get $\dot{b} = \kappa_0 b|b|^{-2\beta}$. With $b = \lambda e^{i\phi}$ this reads $\lambda^{2\beta-1}\dot{\lambda} = \text{Re } \kappa_0$ and $\dot{\phi} = \text{Im } \kappa_0 \lambda^{-2\beta}$, which integrate to (2). The size changes, so $\text{Re } \kappa_0 \neq 0$. Along the spiral $r = r_0\lambda$ and $d\phi/d \ln \lambda = 2P$ in absolute value, so the tangent of the angle between the path and the radius is $2P$ and the arc length from r_0 to 0 is $r_0\sqrt{1 + 4P^2}$. Finally $|\omega_0|t_c = |\text{Im } \kappa_0|/(2\beta|\text{Re } \kappa_0|) = P/\beta$. $\square$

The spiral form (2) of a self-similar collapse in the α -models is in [15] and [9, Eqs. (3)–(6)], in [5, (A.19)], whose exponent $1/(\alpha + 1)$, in a parametrization where the Euler case is $\alpha = 1$, is our $1/(2\beta)$, and in [7, Prop. 2.1]. For $\beta = 1$, $P = |\omega_0|t_c$ is the product studied in [8]. For $\beta \neq 1$ the product $|\omega_0|t_c$ is the rotation per unit decrease of $\ln(t_c - t)$, and it differs from P by the factor $1/\beta$; we state the bound for P , which measures the geometry of the spiral, and for $|\omega_0|t_c$ in Corollary 1.

3 Circulations and rates of a collapsing triangle

Label the sides by the opposite vertices: $r_1 = |z_2 - z_3|$, $r_2 = |z_3 - z_1|$, $r_3 = |z_1 - z_2|$, and put $f_i = r_i^{-2\beta}$. Let $A_s = \text{Im}((z_2 - z_1)(z_3 - z_1))/2$ be the signed area, positive when z_1, z_2, z_3 are counterclockwise, and $A = |A_s|$. Indices (i, j, k) run over the cyclic permutations of $(1, 2, 3)$.

Lemma 2. *Under (1), $d(r_i^2)/dt = (2A_s/\pi)\Gamma_i(f_j - f_k)$. A motion is self-similar and changes size if and only if the triangle is not collinear, its sides are pairwise different, and*

$$\Gamma_i = c \frac{r_i^2}{f_j - f_k} \quad (i = 1, 2, 3) \quad (4)$$

for some real $c \neq 0$. In that case $\text{Re } \kappa = cA_s/\pi$, and exactly one of a triangle and its mirror image collapses.

Proof. For $i = 3$, the terms of $\dot{z}_1 - \dot{z}_2$ proportional to $z_1 - z_2$ do not change $|z_1 - z_2|$, and the rest is $(i\Gamma_3/2\pi)((z_1 - z_3)f_2 - (z_2 - z_3)f_1)$. Since $\text{Im}((z_1 - z_2)(z_1 - z_3)) = \text{Im}((z_1 - z_2)(z_2 - z_3)) = 2A_s$, $d(r_3^2)/dt = 2\text{Re}((z_1 - z_2)(\dot{z}_1 - \dot{z}_2)) = (2A_s/\pi)\Gamma_3(f_1 - f_2)$, and the other sides follow by cyclic relabeling. In a self-similar motion $d \ln r_i^2/dt = 2\text{Re } \kappa$ for every i , so $(2A_s/\pi)\Gamma_i(f_j - f_k)/r_i^2$ is the same for all i ; if the size changes, this common value is not 0, so $A_s \neq 0$, $f_j \neq f_k$ for all pairs, and (4) holds with $c = \pi \text{Re } \kappa/A_s$. Conversely, if (4) holds with $c \neq 0$ on a scalene triangle that is not collinear, the three logarithmic rates coincide, so the ratios of the squared sides are stationary at this shape; the rates of these ratios are functions of the shape alone, multiplied by a power of the size, so the shape is an equilibrium of the reduced motion and is kept for all time, with the orientation, since the area cannot pass through zero. The motion is therefore self-similar, and its size changes because $\text{Re } \kappa = cA_s/\pi \neq 0$. Reflecting the triangle changes the sign of A_s and keeps (4), so it changes the sign of $\text{Re } \kappa$. $\square$

In particular, isosceles, equilateral and collinear triangles never collapse, and every scalene triangle collapses in exactly one orientation, with the circulations (4) fixed up to a real factor. On a scalene triangle, (4) says that the vector $(1/\Gamma_i)$ is proportional to the cross product of (r_i^2) and $(r_i^{2-2\beta})$, that is, orthogonal to both; so it is equivalent to the vanishing of the angular impulse, $\sum_i r_i^2/\Gamma_i = 0$, together with $\sum_i r_i^{-\alpha}/\Gamma_i = 0$, which is the vanishing of $\sum_{j < k} \Gamma_j \Gamma_k r_i^{-\alpha}$ for $\alpha \neq 0$ and the classical condition $\sum_{j < k} \Gamma_j \Gamma_k = 0$ for $\alpha = 0$. Conditions of this kind are given in [1, 2, 7, 13, 14]; Lemma 2 is a short self-contained form of them, with the circulations explicit.

Lemma 3. *For a self-similar motion that changes size, with (4),*

$$2\pi \text{Im } \kappa = \frac{1}{2} \sum_{i=1}^3 \Gamma_i(f_j + f_k) = \frac{c}{2} S, \quad S = \sum_{i=1}^3 r_i^2 \coth\left(\beta \ln \frac{r_k}{r_j}\right), \quad P = \frac{|S|}{8A}. \quad (5)$$

Proof. Since $\dot{z}_1 - \dot{z}_2 = \kappa(z_1 - z_2)$, $\text{Im } \kappa = \text{Im}((\dot{z}_1 - \dot{z}_2)/(z_1 - z_2))$. By (1),

$$2\pi \frac{\dot{z}_1 - \dot{z}_2}{z_1 - z_2} = i(\Gamma_1 + \Gamma_2)f_3 + i\Gamma_3 \frac{f_2(z_1 - z_3) - f_1(z_2 - z_3)}{z_1 - z_2},$$

and by the law of cosines $\operatorname{Re}((z_1 - z_3)/(z_1 - z_2)) = (r_2^2 + r_3^2 - r_1^2)/(2r_3^2)$ and $\operatorname{Re}((z_2 - z_3)/(z_1 - z_2)) = (r_2^2 - r_1^2 - r_3^2)/(2r_3^2)$. Hence

$$2\pi \operatorname{Im} \kappa = (\Gamma_1 + \Gamma_2)f_3 + \Gamma_3 \frac{f_1 + f_2}{2} + \Gamma_3 \frac{(f_2 - f_1)(r_2^2 - r_1^2)}{2r_3^2}.$$

By (4) the last term is $c(r_1^2 - r_2^2)/2$, and also $c(r_1^2 - r_2^2) = \Gamma_1(f_2 - f_3) + \Gamma_2(f_1 - f_3)$. Substituting gives $2\pi \operatorname{Im} \kappa = \frac{1}{2} \sum_i \Gamma_i(f_j + f_k)$, and by (4), $\Gamma_i(f_j + f_k) = cr_i^2(f_j + f_k)/(f_j - f_k) = cr_i^2 \coth(\beta \ln(r_k/r_j))$. With $\operatorname{Re} \kappa = cA_s/\pi$ from Lemma 2, (3) gives $P = |S|/(8A)$. $\square$

The circulations drop out: P depends only on the shape of the triangle. Interchanging two labels reverses the cyclic order, which changes the sign of every term of S , so $|S|$ is a symmetric function of the side lengths. The first identity says that the rotation rate is half the sum of the rates at which the three pairs would rotate in isolation.

4 The bound

Throughout, $\alpha > -1$, that is, $\beta > 1/2$. We use the elementary inequality

$$\frac{1}{\ln(1+x)} \geq \frac{1}{x} + \frac{1}{2} - \frac{x}{12} \quad (x > 0). \quad (6)$$

It is trivial where the right side is negative, that is, for $x \geq 3 + \sqrt{21}$. For $0 < x < 3 + \sqrt{21}$, the function $g(x) = 12x/(12 + 6x - x^2) - \ln(1+x)$ has $g(0) = 0$ and $g'(x) = x^3(24 - x)/((12 + 6x - x^2)^2(1+x)) > 0$, so $\ln(1+x) \leq 12x/(12 + 6x - x^2)$, which is (6).

Proof of Theorem 1. By Lemma 2 a collapsing triangle is scalene, and by Lemma 3, $P = |S|/(8A)$ is symmetric in the sides and invariant under scaling. Order and scale the sides as $r_2 = \rho < r_3 = 1 < r_1$, and write $r_1^2 = 1 + \rho m$. With ψ the angle at z_1 , between the sides of lengths ρ and 1, the law of cosines gives $2 \cos \psi = \rho - m$, so $0 < m < 2 + \rho$, $0 < \rho < 1$, $A = (\rho/2) \sin \psi$ and $4 \sin^2 \psi = 4 - (m - \rho)^2$. Put $X = \rho^{2\beta}$, $Y = (1 + \rho m)^\beta$ and $b = (\beta/2) \ln(1 + \rho m)$. Then $0 < X < 1 < Y$ and, from (5) with $(i, j, k) = (1, 2, 3)$ and its cyclic shifts,

$$S = (1 + \rho m) \frac{1 + X}{1 - X} + \rho^2 \coth b - \frac{Y + X}{Y - X}.$$

(i) The function $t \mapsto (t + X)/(t - X)$ decreases for $t > X$, so $(Y + X)/(Y - X) \leq (1 + X)/(1 - X)$ and $S \geq \rho m + \rho^2 \coth b$. (ii) $\coth b > 1/b$, and (6) with $x = \rho m$ gives $\rho^2/b \geq (\rho/\beta)(2/m + \rho - \rho^2 m/6)$. Hence

$$S > \frac{\rho}{\beta} \Lambda, \quad \Lambda = \beta m + \frac{2}{m} + \rho - \frac{\rho^2 m}{6} > 0,$$

the last because $\rho^2 m/6 < \rho/2$. In particular $S > 0$, and

$$P = \frac{S}{4\rho \sin \psi} > \frac{\Lambda}{2\beta \sqrt{4 - (m - \rho)^2}}.$$

(iii) The identity

$$\Lambda^2 - (1 + 2\beta)(4 - (m - \rho)^2) = \left((1 + \beta)m - \frac{2}{m} - \rho \right)^2 + \frac{\rho^2}{3} \left(1 + 6\beta - \beta m^2 - \rho m + \frac{\rho^2 m^2}{12} \right) \quad (7)$$

holds for all ρ, m, β , and its right side is positive. If $m < 2$, the second bracket exceeds $1+6\beta-4\beta-2 = 2\beta-1 > 0$. If $m \geq 2$, then $2/m \leq 1$ and $\rho < 1$, so $(1+\beta)m-2/m-\rho > (1+\beta)m-2 > 0$, and the second bracket is at least $b(m) = 1+6\beta-\beta m^2-m$. If $b(m) \geq 0$ the right side is positive; otherwise $\rho^2 b(m) > b(m)$, and the right side exceeds $D(m) = ((1+\beta)m-2)^2 + b(m)/3$. Now $D'' = 2\beta^2 + 10\beta/3 + 2 > 0$ and $D'(2) = 4\beta^2 + 8\beta/3 - 1/3 > 0$, so D increases on $[2, \infty)$, and $D(2) = 4\beta^2 + (2\beta-1)/3 > 0$. Hence $\Lambda > \sqrt{1+2\beta}\sqrt{4-(m-\rho)^2}$ and $P > \sqrt{1+2\beta}/(2\beta) = B(\alpha)$.

Sharpness. Let $\rho \rightarrow 0$ with m fixed in $(0, 2)$. Since $2\beta > 1$, $X = o(\rho)$, so the first and last terms of S are $1 + \rho m + o(\rho)$ and $1 + o(\rho)$, while $\rho^2 \coth b = 2\rho/(\beta m) + O(\rho^2)$. Hence $S/\rho \rightarrow m + 2/(\beta m)$ and

$$P \rightarrow P_0(m) = \frac{\beta m^2 + 2}{2\beta m \sqrt{4 - m^2}},$$

whose only critical point on $(0, 2)$ is $m_* = \sqrt{2/(1+\beta)}$, with $P_0(m_*) = \sqrt{1+2\beta}/(2\beta)$. By Lemma 2 each of these triangles, or its mirror image, collapses with the circulations (4), so P comes arbitrarily close to $B(\alpha)$. $\square$

In the extremal limit $\rho \rightarrow 0$ the circulations (4) are, up to a factor, $\Gamma_2 \approx 1$ at the vertex opposite the shortest side and $\Gamma_1 \approx -\Gamma_3$ of order $\rho^{2\beta-1}$ at its ends: two weak vortices of opposite sign, at distance ρ , spiral into a strong vortex, and the line through them makes the angle $\arccos(1/\sqrt{2(1+\beta)})$ with the line to the strong vortex. For $\alpha = 0$ this is the limit $\mu \rightarrow 0$ of [8, Corollary 1].

Corollary 1. *Let $\alpha > -1$. In every self-similar collapse of three point vortices under (1), each vortex travels more than $(4+\alpha)/(2+\alpha)$ times its initial distance from the collision point, the angle between its path and the direction to that point exceeds $\arccos((2+\alpha)/(4+\alpha))$, and $|\omega_0|t_c > 2\sqrt{3+\alpha}/(2+\alpha)^2$. None of these constants can be improved.*

Proof. By Lemma 1 the three quantities are $\sqrt{1+4P^2}$, $\arctan 2P$ and P/β , which increase with P , and $\sqrt{1+4B^2} = (4+\alpha)/(2+\alpha)$. $\square$

	$\alpha = 0$ (Euler)	$\alpha = 1/2$	$\alpha = 1$ (SQG)	$\alpha = 3/2$	$\alpha = 2$
$B(\alpha)$	$\sqrt{3}/2 = 0.8660$	$\sqrt{14}/5 = 0.7483$	$2/3$	0.6061	$\sqrt{5}/4 = 0.5590$
path ratio	2	$9/5$	$5/3$	$11/7$	$3/2$
angle	60°	56.25°	53.13°	50.48°	48.19°

Table 1: The bound of Theorem 1 and the two constants of Corollary 1 for some values of α .

Since $d \ln B/d\alpha = -(4+\alpha)/(2(3+\alpha)(2+\alpha)) < 0$, the bound decreases strictly in α , from $\sqrt{3}/2$ in the Euler case through $2/3$ for SQG to 0 as $\alpha \rightarrow \infty$: more local interactions allow tighter spirals, but never a collapse without rotation (Figure 1). The restriction $\alpha > -1$, that is, $2\beta > 1$, is the range in which the velocity (1) induced by a vortex decays with distance; the proof uses it in step (iii).

Remark 1 (below $\alpha = -1$). The sharpness argument holds for every $\alpha > -2$, since exactly $S - \rho m - \rho^2 \coth b = 2X((1+\rho m)/(1-X) - 1/(Y-X))$, which is $O(X\rho)$ for every $\beta > 0$. Steps (i) and (ii) hold for every $\beta > 0$, and so does the case $m \geq 2$ of step (iii) as soon as $D(2) = (12\beta^2 + 2\beta - 1)/3 > 0$. In the case $m < 2$, if $b(m) = 1 + 6\beta - \beta m^2 - m > 0$ the right side of (7) is positive; otherwise, when $(1+\beta)m - 2/m - 1 \geq 0$, it exceeds $((1+\beta)m - 2/m - 1)^2 + b(m)/3$. An interval-arithmetic subdivision of $0.2622 \leq \beta \leq 1/2$, $3/2 \leq m \leq 2$ (the program `verify_alpha_extension.py`) shows that one of these holds on every box, and $b(m) \geq b(3/2) = 15\beta/4 - 1/2 > 0$ for $m \leq 3/2$. Hence




Figure 1: (a) The bound $B(\alpha)$ of Theorem 1 (curve) and $P = |S|/(8A)$ for 24395 random collapsing triangles at 61 values of α (dots, spread slightly in α for visibility); none lies below the curve, which is approached as the shortest side tends to 0. (b) The paths of the three vortices of an SQG collapse ($\alpha = 1$) with sides 0.45, 1 and $(1 + 0.45m_*)^{1/2}$ and the circulations of Lemma 2, integrated from (1) to $10^{-6}t_c$ before the collapse; dots mark the initial positions and the cross the collision point. The program is `plot_alpha_winding.py`.

Theorem 1 and Corollary 1 hold for every $\alpha \geq -59/40$, with this computer-assisted step. At $\alpha = -1.5$ and $\alpha = -1.8$ numerical minimization of (5), in arithmetic whose precision grows with $|\ln \rho|$, finds no value below $\sqrt{1 + 2\beta}/(2\beta)$ either, but we do not claim a bound there.

Remark 2 (two equal circulations). For circulations $(1, -\gamma, 1)$ the collapsing triangles form a single family. For $\alpha = 2$ it is $r_1 r_3 = r_2^2$, where r_2 joins the two equal vortices, the ratio is $\gamma = r_2^2/(r_1^2 + r_3^2)$, and Lemma 3 gives

$$P^2 = \frac{(1 + \gamma)(2\gamma^2 - 2\gamma + 1)^2}{4(1 - 2\gamma)(1 + 2\gamma)(3\gamma - 1)}, \quad \frac{1}{3} < \gamma < \frac{1}{2},$$

whose minimum, $1.1039451679\dots$, is a root of $27648y^8 - 29952y^6 - 3200y^4 - 1556y^2 - 125$; at $\gamma = 1/3$ the triangle degenerates to the collinear configuration with sides in the ratio $\varphi : 1 : 1/\varphi$, φ the golden ratio. For SQG the interval $0.387464\dots < \gamma < 1/2$ of $[1, 2]$ has as its lower end the root in $(0, 1/2)$ of $4\gamma^4 - 8\gamma^3 + \gamma^2 - 2\gamma + 1$, where the triangle becomes collinear.

5 More than three vortices

Self-similar motion of any number of vortices has the same form [4, 6, 12], $\dot{z}_j = \kappa(z_j - z_c)$ and P defined by (3). In the Euler case the bound of [8] does not extend to more vortices: the seven-vortex collapse of Demina and Kudryashov [4, Table 1] has $P = 12433/(1240\sqrt{155}) = 0.805\dots$ [8]. A numerical minimization of $\text{Im } \kappa/(-2 \text{Re } \kappa)$ over the circulations and positions of four vortices, from more than two hundred random starts, ends at a single configuration up to relabeling, with

$$P_4 = 0.79789678387986348076760587182\dots,$$

a strict local minimum (the Hessian of the Lagrangian on the three-dimensional tangent space of the constraints has eigenvalues 2.21, 12.58 and 18.75). In the normalization $\Gamma_1 = 1$, $z_c = 0$, $\kappa = -1 + ib$ it has $b = 1.5957935677597269615 \dots$ and

$$\begin{aligned}\Gamma &= (1, 0.1673914318538, 0.1041331880362, -0.2272513003829), \\ z &= (0.0166495669172, -0.2191226402232 - 0.3290676220068i, \\ &\quad -0.1579874300122 - 0.2864759934272i, -0.1605333843048 - 0.3736601672808i).\end{aligned}$$

Families of larger numerical minimizers go lower still: $P = 0.5172291322 \dots$ for 33 vortices and $P = 0.4981844392 \dots$ for 61 vortices, so a self-similar collapse can turn by less than half a radian per unit decrease of $\ln \lambda^2$. Whether P can be arbitrarily small for large numbers of vortices is open; the families computed so far extrapolate to about 0.477. The same happens for $\alpha \neq 0$: minimizing over four vortices in the same way gives $P_4 = 0.5499151189 \dots$ for SQG, against $B(1) = 2/3$, and $P_4 = 0.4116738600 \dots$ for $\alpha = 2$, against $B(2) = \sqrt{5}/4$, both strict local minima with one strong and three weak vortices, as for $\alpha = 0$. These are numerical results, not proofs.

6 Numerical verification

The program `verify_alpha_winding.py` checks each step in high-precision arithmetic, and its output is `data/verify_alpha_winding.txt`. On 700 random scalene triangles with $\alpha \in \{0, 0.3, 1, 1.7, 3\}$, the circulations (4) make the Biot–Savart velocities (1) self-similar to 10^{-38} , and P computed from the velocities agrees with (5) to $3 \cdot 10^{-38}$. The derivative of g , the identity (7) and the factorization of $h(\rho) - h(1)$ are verified symbolically, and the critical point and value of P_0 as well. The chain of inequalities in the proof holds at 20000 random points with β from just above $1/2$ to 20 and ρ down to 10^{-8} , computed with precision adapted to $\rho^{2\beta}$; in double precision the first step appears to fail at small ρ and large β , where its margin, of order $\rho^{2\beta}$, is below the rounding error. For α from -1.8 to -1 , outside the theorem, the minimum of $P - B(\alpha)$ over the ordered triangles is positive in every run, again only with precision adapted to ρ . For $\alpha = 1$, the scalene triangles on which (4) gives two equal circulations carry the ratios $-\Gamma_2/\Gamma_1$ from 0.3876 to 0.49999 on a grid, in agreement with the interval of [1, Lemma 1], and at $\Gamma = 0.49$ the ratio of sides is 0.7515, against 0.751484 in [1]. The program `verify_alpha_extension.py` performs the interval-arithmetic step of Remark 1, and `symmetric_family.py` in `research/generalizations-2026-09-24/checks/` checks Remark 2. Near-extremal triangles at $\rho = 10^{-6}$ and $m = m_*$ collapse, and $P - B(\alpha)$ is $1.3 \cdot 10^{-12}$, $1.8 \cdot 10^{-13}$, $1.6 \cdot 10^{-13}$ and $1.6 \cdot 10^{-13}$ for $\alpha = 0, 1/2, 1, 2$. The four-vortex minima for $\alpha = 1, 2$ were refined by Newton’s method on the Lagrange conditions in 80-digit arithmetic (residual 10^{-56}), and 800 constrained perturbations of each lower P in no case. The four-, thirty-three- and sixty-one-vortex configurations of Section 5 satisfy the self-similarity equations to 10^{-29} , 10^{-39} and 10^{-28} relative in independent Biot–Savart checks.

7 Open problems

1. Is there a positive lower bound for P over self-similar collapses of any number of vortices in the Euler case? The smallest values found decrease with the number of vortices, to 0.4981844392 $\dots$ for 61 vortices, and the families computed extrapolate to about 0.477.
2. Is $P_4 = 0.7978967838 \dots$ of Section 5 the minimum over all collapses of four vortices, and is it algebraic? No integer polynomial of degree at most 16 with coefficients below 10^7 vanishes at P_4 or P_4^2 .

3. Does Theorem 1 hold for $-2 < \alpha < -59/40$? Numerically it does.
4. Are the four-vortex values $0.5499151189\dots$ (SQG) and $0.4116738600\dots$ ($\alpha = 2$) of Section 5 the infima for four vortices, and what are the infima for more vortices when $\alpha \neq 0$?
5. For fixed circulations and $\alpha \neq 0$ only finitely many triangles collapse; for $\alpha = 2$ they are at most two and P is an explicit algebraic function of the circulations. Is there such a formula for SQG, where the collapsing triangles are the roots of a quartic?

8 Discussion

Self-similar collapse of three vortices in the α -models has been studied from several directions: Badin and Barry [1] derive the necessary conditions for collapse in Nambu form and show that in the SQG case collapse is self-similar when the energy vanishes and not self-similar otherwise; for circulations $(1, -\Gamma, 1)$ they find self-similar SQG collapse exactly for $0.387464\dots < \Gamma < 1/2$ [1, Lemma 1], an interval that Lemma 2 reproduces; Reinaud, Dritschel and Scott [14] give the conditions for self-similar collapse of three vortices and map the collapse time over the collapsing configurations; Reinaud [13] gives, for SQG, two necessary conditions and the collapse time in terms of the side lengths (as quoted in [2]), which has the form of the rate $\text{Re } \kappa$ of Lemma 2 at $\alpha = 1$; Chen and Liu [2] prove that the existence of self-similar collapse and expansion depends only on the circulations, give explicit expressions for all nontrivial self-similar solutions [2, Theorems 2.1, 2.2], and obtain the interval of Badin and Barry analytically [2, Theorem 4.1], a description from the side of the circulations that Lemma 2 complements from the side of the shape; Yasunaga, Ootobe and Iwayama [15] give necessary conditions and the orbits of self-similar motion, and Iwayama and Yajima [9] show that the collapsing motions are linearly unstable and the expanding ones stable; Iwayama, Yajima and Watanabe [10] prove that collapse that is not self-similar can occur for $0 < \alpha < 2$ and cannot for $-1 \leq \alpha < 0$; and Grotto and Pappalettera [7, Prop. 2.1] characterize the self-similar motions and write the collapsing spiral, as a step toward bursts out of arbitrary configurations. We have not found a bound on the winding, the spiral angle or the distance traveled in the literature. In the Euler case the closed forms of the rates go back to [3] and [11], and the bound $P > \sqrt{3}/2$ with its minimum for each ratio of circulations is in [8]; Theorem 1 gives, for $\alpha = 0$, a shorter proof of that bound through the side lengths.

The bound concerns self-similar collapse only; it says nothing about other singular motions of the α -models, such as the collapses at nonzero energy found in [1, 10]. Its value lies in the geometry: however the circulations are chosen, a collapsing triangle turns by more than $B(\alpha)$ radians per unit decrease of the logarithm of its squared size, and the infimum is realized only by a degenerate dipole beside a strong vortex.

Data availability. The programs `verify_alpha_winding.py` and `plot_alpha_winding.py` and their output are in the repository <https://github.com/SharpMeow/alpha-winding>, the programs in `code/` and their output in `data/`; the many-vortex configurations and their checks are in `research/generalizations-2026-09-24/`.

Funding. This research received no external funding.

This work was prepared with AI assistance. The author takes full responsibility for its content.

References

- [1] G. Badin and A. M. Barry, Collapse of generalized Euler and surface quasigeostrophic point vortices, *Phys. Rev. E* **98** (2018) 023110; arXiv:1805.10127.

- [2] J. Chen and Q. Liu, Sufficient and necessary conditions for self-similar motions of three point vortices in generalized fluid systems, *Physica D* **470** (2024) 134392.
- [3] R. Conte and L. de Seze, Exact solution of the planar motion of three arbitrary point vortices, report DPhG/PSRM/1697/80, CEN Saclay (1980); arXiv:1511.00069.
- [4] M. V. Demina and N. A. Kudryashov, Rotation, collapse, and scattering of point vortices, *Theor. Comput. Fluid Dyn.* **28** (2014) 357–368; doi:10.1007/s00162-014-0319-4.
- [5] M. Donati and L. Godard-Cadillac, Hölder regularity for collapses of point vortices, *Nonlinearity* **36** (2023) 5773; arXiv:2111.14230.
- [6] T. Gotoda, Self-similar motions and related relative equilibria in the N -point vortex system, *J. Dyn. Differ. Equ.* **33** (2021) 1759–1777; arXiv:2002.09624.
- [7] F. Grotto and U. Pappalettera, Collapse and burst of generalized surface quasi-geostrophic point vortices, preprint, arXiv:2505.19782 (2025).
- [8] C. Hendrick, Minimal winding in the self-similar collapse of three point vortices and of two concentric vortex polygons, preprint (2026), <https://github.com/SharpMeow/minimal-winding>.
- [9] T. Iwayama and T. Yajima, Linear stability of self-similar motions of three point vortices in a generalized two-dimensional fluid system, *J. Phys. Soc. Jpn.* **92** (2023) 084401.
- [10] T. Iwayama, S. Yajima and T. Watanabe, Non-self-similar collapse of three point vortices in a generalized two-dimensional fluid system, *J. Phys. A* **58** (2025) 075701.
- [11] Y. Kimura, Similarity solution of two-dimensional point vortices, *J. Phys. Soc. Jpn.* **56** (1987) 2024–2030.
- [12] K. A. O’Neil, Relative equilibrium and collapse configurations of four point vortices, *Regul. Chaotic Dyn.* **12** (2007) 117–126.
- [13] J. N. Reinaud, Self-similar collapse of three geophysical vortices, *Geophys. Astrophys. Fluid Dyn.* **115** (2021) 369–392.
- [14] J. N. Reinaud, D. G. Dritschel and R. K. Scott, Self-similar collapse of three vortices in the generalised Euler and quasi-geostrophic equations, *Physica D* **434** (2022) 133226.
- [15] T. Yasunaga, N. Otobe and T. Iwayama, *J. Phys. Soc. Jpn.* **90** (2021) 124401.