

Probability and Statistics: Lecture-16

Monsoon-2020

by Pawan Kumar (IIIT, Hyderabad)

on September 16, 2020

» Table of contents

1. Examples of Distributions

2. Expectations of Some Distributions

3. Variance and Standard Deviation: Understand Variability in Data

4. Higher Order Moments and Moment Generating Function

» Example of Poisson Distribution...

» Example of Poisson Distribution...

Example

We want to send a bit string of length $n = 10^4$.

» Example of Poisson Distribution...

Example

We want to send a bit string of length $n = 10^4$. The probability that a bit can be corrupted is $p = 10^{-6}$.

» Example of Poisson Distribution...

$n \gg 1$

Binomial (n, p)

Example

We want to send a bit string of length $n = 10^4$. The probability that a bit can be corrupted is $p = 10^{-6}$. What is the probability that the message will arrive **uncorrupted**?

$p \ll 1$

$\begin{cases} n \rightarrow \infty \\ p \rightarrow \text{small} \end{cases}$

» Example of Poisson Distribution...

Example

We want to send a bit string of length $n = 10^4$. The probability that a bit can be corrupted is $p = 10^{-6}$. What is the probability that the message will arrive **uncorrupted**?

- * We can solve this using a Poisson with $\lambda = np = 10^4 10^{-6}$

» Example of Poisson Distribution...

Example

We want to send a bit string of length $n = 10^4$. The probability that a bit can be corrupted is $p = 10^{-6}$. What is the probability that the message will arrive **uncorrupted**?

- * We can solve this using a Poisson with $\lambda = np = 10^4 10^{-6}$
- * With $X \sim \text{Poisson}(0.01)$ be the number of corrupted bits

» Example of Poisson Distribution...

Example

We want to send a bit string of length $n = 10^4$. The probability that a bit can be corrupted is $p = 10^{-6}$. What is the probability that the message will arrive **uncorrupted**?

- * We can solve this using a Poisson with $\lambda = np = 10^4 10^{-6}$
- * With $X \sim \text{Poisson}(0.01)$ be the number of corrupted bits
- * Using the PMF for Poisson

» Example of Poisson Distribution...

Example

We want to send a bit string of length $n = 10^4$. The probability that a bit can be corrupted is $p = 10^{-6}$. What is the probability that the message will arrive **uncorrupted**?

- * We can solve this using a Poisson with $\lambda = np = 10^4 10^{-6}$
- * With $X \sim \text{Poisson}(0.01)$ be the number of corrupted bits
- * Using the PMF for Poisson

$$\begin{aligned} P(X = 0) &= \frac{\lambda^i}{i!} e^{-\lambda} \\ &= \frac{0.01^0}{0!} e^{-0.01} \\ &\approx 0.9900498 \end{aligned}$$

- * We could have solved this using Binomial $X \sim \text{Binomial}(10^4, 10^{-6})$. This may have been harder to compute!
- * When n large, and p small: can use Poisson!

» Another Example of Poisson Distribution...

» Another Example of Poisson Distribution...

Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate.

» Another Example of Poisson Distribution...

Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this.

» Another Example of Poisson Distribution...

Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this. Suppose there are an average of 2.8 major earthquakes in the world each year.

» Another Example of Poisson Distribution...

Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this. Suppose there are an average of 2.8 major earthquakes in the world each year. What is the probability of getting more than one major earthquake next year?

$$\begin{aligned} X &\sim \text{Poisson}(2.8), \quad X = \# \text{ earthquakes} \\ P(X > 1) \\ &= 1 - P(X = 0) - P(X = 1) \end{aligned}$$

Handwritten notes and diagrams:

- A box around $X > 1$ with an arrow pointing to it from the text $X \sim \text{Poisson}(2.8)$.
- A diagram showing a horizontal line with tick marks representing values 0, 1, 2, The tick mark for 0 is circled, and the tick mark for 1 is marked with an 'X'. The text $= 1 - P(X=0) - P(X=1)$ is written below the line, with arrows pointing to the 0 and 1 tick marks.

» Another Example of Poisson Distribution...

Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this. Suppose there are an average of 2.8 major earthquakes in the world each year. What is the probability of getting more than one major earthquake next year?

- * Let $X \sim \text{Poisson}(2.8)$ be the number of major earthquakes next year. We want $P(X > 1)$.

» Another Example of Poisson Distribution...

Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this. Suppose there are an average of 2.8 major earthquakes in the world each year. What is the probability of getting more than one major earthquake next year?

- * Let $X \sim \text{Poisson}(2.8)$ be the number of major earthquakes next year. We want $P(X > 1)$.
- * We can use

$$\underline{P(X > 1)} = 1 - \underline{P(X = 0)} - \underline{P(X = 1)}.$$

Using PMF for Poisson

» Another Example of Poisson Distribution...

Example of Poisson Distribution

The Poisson distribution is often used to model the number of events that occur independently at any time in an interval of time or space, with a **constant** average rate. Earthquakes are a good example of this. Suppose there are an average of 2.8 major earthquakes in the world each year. What is the probability of getting more than one major earthquake next year?

- * Let $X \sim \text{Poisson}(2.8)$ be the number of major earthquakes next year. We want $P(X > 1)$.
- * We can use

Using PMF for Poisson

$$\begin{aligned} P(X > 1) &= 1 - P(X=0) - P(X=1). \\ P(X > 1) &= 1 - P(X=0) - P(X=1) \\ &= 1 - e^{-2.8} \frac{2.8^0}{0!} - e^{-2.8} \frac{2.8^1}{1!} \\ &= 1 - e^{-2.8} - 2.8e^{-2.8} \\ &\approx 1 - 0.06 - 0.17 = 0.77 \end{aligned}$$

77%

» Expectation of Poisson Distribution...

» Expectation of Poisson Distribution...

Expectation of Poisson

Let $X \sim \text{Poisson}(\lambda)$, with PMF given by

$$\underline{p_X(k)} = \begin{cases} \frac{e^{-\lambda} \lambda^k}{k!}, & k \in R_X \\ 0 & \text{otherwise} \end{cases}$$

$$E[X] = \sum x \underline{p(x)}$$

Show that the expectation $E[X] = \lambda$.

» Expectation of Poisson Distribution...

Expectation of Poisson

Let $X \sim \text{Poisson}(\lambda)$, with PMF given by

$$P_X(k) = \begin{cases} \frac{e^{-\lambda} \lambda^k}{k!}, & k \in R_X \\ 0 & \text{otherwise} \end{cases}$$

Show that the expectation $E[X] = \lambda$.

Proof

$$E[X] = \sum_{x=0}^{\infty} x \frac{e^{-\lambda} \lambda^x}{x!} = \sum_{x=1}^{\infty} x \frac{e^{-\lambda} \lambda^x}{x!} = e^{-\lambda} \sum_{x=1}^{\infty} \frac{\lambda^x}{(x-1)!}$$

$$= e^{-\lambda} \sum_{z=0}^{\infty} \frac{\lambda^{z+1}}{z!} = \lambda e^{-\lambda} \sum_{z=0}^{\infty} \frac{\lambda^z}{z!} = \lambda$$

Recall Calculus / Real Analysis

If $f: \mathbb{R} \rightarrow \mathbb{R}$ is infinitely diff., then f can be expanded around $x=a$

$$f(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots$$

$x=0$

$$e^x = e^0 + \frac{e^0}{1!}x + \frac{e^0}{2!}x^2 + \dots$$

Taylor's expansion

» Expectation of Binomial Distribution...

» Expectation of Binomial Distribution...

Expectation of a Binomial Distribution

Show that $E[X] = np$, where $X \sim \text{Binomial}(n, p)$, $0 < p < 1$ with PMF given as follows

$$P_X(k) = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k} & \text{for } k = 0, 1, 2, \dots, n \\ 0 & \text{otherwise,} \end{cases}$$

» Expectation of Binomial Distribution...

Expectation of a Binomial Distribution

Show that $E[X] = np$, where $X \sim \text{Binomial}(n, p)$, $0 < p < 1$ with PMF given as follows

$$P_X(k) = \begin{cases} \binom{n}{k} p^k (1-p)^{n-k} & \text{for } k = 0, 1, 2, \dots, n \\ 0 & \text{otherwise,} \end{cases}$$

Proof

$$E[X] = \sum_{x=0}^n x \binom{n}{x} p^x q^{n-x} = \sum_{x=1}^n x \binom{n}{x} p^x q^{n-x}.$$

Handwritten notes: A '0' is written above the sum. 'p x PMF' is written below the binomial coefficient.

$$\left[\text{For } 0 < x \leq n, \binom{n}{x} = \frac{n!}{(n-x)!x!} = \frac{n!}{(n-x)!(x-1)!} = n \binom{n-1}{x-1} \right]$$

Handwritten notes: The binomial coefficient and the final term are circled in orange. Arrows indicate the cancellation of $n!$ and the shift of indices.

$$\Rightarrow E[X] = \sum_{x=1}^n n \binom{n-1}{x-1} p^x q^{n-x} = \sum_{z=0}^{n-1} n \binom{n-1}{z} p^{z+1} q^{n-1-z} = np.$$

Handwritten notes: The binomial coefficient $\binom{n-1}{z}$ is circled in orange. The final result np is circled in orange. A note 'take n & p common' with an arrow points to the n and p in the sum. Another note 'z-1-->0' is written below the sum.

Motivation for Variance

- * The expectation value $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:

Motivation for Variance

- * The expectation value $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000

Motivation for Variance

- * The expectation value $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000
 - * A fair coin is tossed, and if it is head you get Rs 2000

Motivation for Variance

- * The expectation value $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000
 - * A fair coin is tossed, and if it is head you get Rs 2000
 - * A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million

Motivation for Variance

- * The expectation value $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000
 - * A fair coin is tossed, and if it is head you get Rs 2000
 - * A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million
- * Let X be the discrete random variable corresponding to your winnings. In all three cases, $E(X) = \text{Rs } 1000$, but you will agree that **your chances of actually getting any money are quite different in all three cases.**

Motivation for Variance

- * The expectation value $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000
 - * A fair coin is tossed, and if it is head you get Rs 2000
 - * A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million
- * Let X be the discrete random variable corresponding to your winnings. In all three cases, $E(X) = \text{Rs } 1000$, but you will agree that **your chances of actually getting any money are quite different in all three cases**.
- * The three cases differ in the spread of PMF. This can be measured by **Variance**.

» Variance and Standard Deviation...

Variance

Let $\mu = E[X]$. The **variance** $\text{Var}(X)$ of X is defined by

» Variance and Standard Deviation...


$$E[\underbrace{g(x)}] = \sum_x g(x) p_x(x)$$

Variance

Let $\mu = E[X]$. The **variance** $\text{Var}(X)$ of X is defined by

$$\text{Var}(X) = E(\underbrace{(X - \mu)^2}_{\rightarrow g(x)}) = \sum_x \underbrace{(x - \mu)^2}_{\leftarrow \text{sometimes use } p_x \text{ or } f_x} \underbrace{f_X(x)}$$

» Variance and Standard Deviation...



Variance

Let $\mu = E[X]$. The **variance** $\text{Var}(X)$ of X is defined by

$$\text{Var}(X) = E((X - \mu)^2) = \sum_x \underbrace{(x - \mu)^2}_{\leftarrow 0} \underbrace{f_X(x)}$$



The **positive** square root is called the **standard deviation** and it is usually denoted by σ , hence

» Variance and Standard Deviation...

Variance

Let $\mu = E[X]$. The **variance** $\text{Var}(X)$ of X is defined by

$$\text{Var}(X) = E((X - \mu)^2) = \sum_x (x - \mu)^2 f_X(x)$$

The **positive** square root is called the **standard deviation** and it is usually denoted by σ , hence

$$\sigma(X) = \sqrt{\sum_x (x - \mu)^2 f_X(x)} = \sqrt{\text{Var}(X)}.$$

» Example of Computing Variance...

» Example of Computing Variance...

$$R_X = \{1, 2, 3, 4, 5, 6\}$$

Example

Let X be the value on one roll of a 6-sided die. Recall that $E[X] = 7/2$. What is $\text{Var}(X)$?

$$\text{Var}(X) = E[(X - \mu)^2]$$

$$= \sum_x (x - 7/2)^2 \underbrace{P_X(x)}$$

$$= \sum_x (x - 7/2)^2 \cdot \frac{1}{6}$$

$$= \left(1 - 7/2\right)^2 \cdot \frac{1}{6} + \left(2 - 7/2\right)^2 \cdot \frac{1}{6} + \dots + \left(6 - 7/2\right)^2 \cdot \frac{1}{6}$$

» Variance

Motivation for Variance

- * The **expectation value** $E(X)$ (also called the mean) of a discrete random variable is a rather coarse measure of how X is distributed. For example, consider the following three situations:

Motivation for Variance

- * The **expectation value** $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000

Motivation for Variance

- * The **expectation value** $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000
 - * A fair coin is tossed, and if it is head you get Rs 2000

Motivation for Variance

- * The **expectation value** $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000
 - * A fair coin is tossed, and if it is head you get Rs 2000
 - * A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million

Motivation for Variance

- * The **expectation value** $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000
 - * A fair coin is tossed, and if it is head you get Rs 2000
 - * A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million

» Variance

Motivation for Variance

- * The **expectation value** $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000
 - * A fair coin is tossed, and if it is head you get Rs 2000
 - * A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million

Calculate Variance

- * **First case:** There is only one outcome and it is the mean, variance is 0

» Variance

$$R_T = \begin{cases} 0 \\ 2000 \end{cases}$$

Motivation for Variance

- * The **expectation value** $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000
 - * A fair coin is tossed, and if it is head you get Rs 2000
 - * A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million

Calculate Variance

- * **First case:** There is only one outcome and it is the mean, variance is 0
- * **Second case:** $\text{Var}(X) = \frac{1}{2}(\underbrace{2000 - 1000}_{n - \mu})^2 + \frac{1}{2}(\underbrace{0 - 1000}_{\mu - n})^2 = 10^6$

» Variance

Motivation for Variance

- * The **expectation value** $E(X)$ (also called the mean) of a discrete random variable is a rather **coarse** measure of how X is distributed. For example, consider the following three situations:
 - * You get Rs 1000
 - * A fair coin is tossed, and if it is head you get Rs 2000
 - * A number is chosen from 1 to 1000 and if you can guess this, you get Rs 1 Million

Calculate Variance

- * **First case:** There is only one outcome and it is the mean, variance is 0
- * **Second case:** $\text{Var}(X) = \frac{1}{2}(2000 - 1000)^2 + \frac{1}{2}(0 - 1000)^2 = 10^6$
- * **Third case:** $\text{Var}(X) = 10^{-3}(10^6 - 10^3)^2 + 999 \times 10^{-3}(0 - 10^3)^2 \approx 10^9$

» Remarks on Expectation, Variance, Standard Deviation...

» Remarks on Expectation, Variance, Standard Deviation...

Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, “Wow! That’s great.”

» Remarks on Expectation, Variance, Standard Deviation...

Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, “Wow! That’s great.” But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that’s a lot of variation in terms of how much money you can make,

» Remarks on Expectation, Variance, Standard Deviation...

Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, “Wow! That’s great.” But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that’s a lot of variation in terms of how much money you can make, so the average starting salary of Rs 70,000 isn’t as informative in the end, is it?

» Remarks on Expectation, Variance, Standard Deviation...

Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, “Wow! That’s great.” But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that’s a lot of variation in terms of how much money you can make, so the average starting salary of Rs 70,000 isn’t as informative in the end, is it?

- * If the standard deviation was only Rs 5,000, you would have a much better idea of what to expect for a starting salary at that company

» Remarks on Expectation, Variance, Standard Deviation...

Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, “Wow! That’s great.” But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that’s a lot of variation in terms of how much money you can make, so the average starting salary of Rs 70,000 isn’t as informative in the end, is it?

- * If the standard deviation was only Rs 5,000, you would have a much better idea of what to expect for a starting salary at that company
- * Without calculating standard deviation, you can’t get a handle on whether the data are close to the average

» Remarks on Expectation, Variance, Standard Deviation...

Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, “Wow! That’s great.” But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that’s a lot of variation in terms of how much money you can make, so the average starting salary of Rs 70,000 isn’t as informative in the end, is it?

- * If the standard deviation was only Rs 5,000, you would have a much better idea of what to expect for a starting salary at that company
- * Without calculating standard deviation, you can’t get a handle on whether the data are close to the average

~~Same Standard Deviation Does Not Imply Same Data set~~

For example, the data sets 199, 200, 201 and 0, 200, 400 both have the same average (200) yet they have very different standard deviations.

» Remarks on Expectation, Variance, Standard Deviation...

Example

If you are told that the average starting salary for someone working at Company Statistix is Rs 70,000, you may think, “Wow! That’s great.” But if the standard deviation for starting salaries at Company Statistix is Rs 20,000, that’s a lot of variation in terms of how much money you can make, so the average starting salary of Rs 70,000 isn’t as informative in the end, is it?

- * If the standard deviation was only Rs 5,000, you would have a much better idea of what to expect for a starting salary at that company
- * Without calculating standard deviation, you can’t get a handle on whether the data are close to the average

Same Standard Deviation Does Not Imply Same Dataset

For example, the data sets 199, 200, 201 and 0, 200, 400 both have the same average (200) yet they have very different standard deviations. The first data set has a very small standard deviation ($s=1$) compared to the second data set ($s=200$).

» Another expression for the variance

Theorem (Another Expression for Variance)

If X is a discrete random variable with mean μ , then

$$\text{Var}(X) = \underbrace{E[X^2]} - \underbrace{\mu^2}$$

» Another expression for the variance

Theorem (Another Expression for Variance)

If X is a discrete random variable with mean μ , then

$$\text{Var}(X) = E[X^2] - \mu^2$$

Proof

$$\begin{aligned}\text{Var}(X) &= \sum_x (x - \mu)^2 p_X(x) = \sum_x (x^2 - 2\mu x + \mu^2) p_X(x) \\ &= \sum_x x^2 p_X(x) - 2\mu \sum_x x p_X(x) + \mu^2 \sum_x p_X(x) \\ &= E[X^2] - 2\mu E[X] + \mu^2 = E[X^2] - \mu^2\end{aligned}$$

» Properties of Variance...

» Properties of Variance...

Theorem

Let X be a discrete random variable and α a constant. Then

$$\text{Var}(\alpha X) = \alpha^2 \text{Var}(X) \quad \text{and} \quad \text{Var}(X + \alpha) = \text{Var}(X)$$

$$E[X] = \mu$$

$$E[\alpha X] = \alpha \mu$$

$$\begin{aligned} \text{Var}(\alpha X) &= E[(\alpha X)^2] - \alpha^2 \mu^2 \\ &= \alpha^2 E[X^2] - \alpha^2 \mu^2 \\ &= \alpha^2 (E[X^2] - \mu^2) \\ &= \alpha^2 \text{Var}(X) \end{aligned}$$

$$\text{Var}(X + \alpha) = E[(X + \alpha)^2] - (\mu + \alpha)^2$$

$$= E[X^2 + 2\alpha X + \alpha^2] - (\mu + \alpha)^2$$

$$= E[X^2] + 2\alpha E[X] + \alpha^2 - (\mu + \alpha)^2$$

$$= E[X^2] + 2\alpha \mu + \alpha^2 - (\mu^2 + 2\alpha \mu + \alpha^2)$$

$$= E[X^2] - \mu^2$$

$$= \text{Var}(X)$$

» Computing Variance: Binomial

Binomial $\binom{n}{k}$

$$E[X] = np$$

Variance of Binomial Distribution

Let $X \sim \text{Binomial}(n, p)$. Then the variance $\text{Var}(X) = np(1 - p)$.

$$\text{Var}(X) = E(X^2) - \mu^2 = E[X(X-1)] + E[X] - E[X]^2$$

$$E[X(X-1)] = \sum_{x=0}^n x(x-1) \binom{n}{x} p^x (1-p)^{n-x}$$

$$\sum_{x=2}^n \frac{n!}{(x-2)! (n-x)!} p^x (1-p)^{n-x}$$

» Variance of Binomial Distribution...

$$= \sum_{x=2}^n \frac{n!}{(x-2)! (n-x)!} p^x (1-p)^{n-x}$$

$$= n(n-1)p^2 \sum_{x=2}^n \frac{(n-2)!}{(x-2)! (n-x)!} p^{x-2} (1-p)^{n-x}$$

$$\begin{cases} m = n-2 \\ y = x-2 \end{cases} = n(n-1)p^2 \sum \frac{m!}{y! (m-y)!} p^y (1-p)^{m-y}$$

$$= \underline{n(n-1)p^2} (p + (1-p))^m = n(n-1)p^2 = E[X(X-1)]$$

» Variance of Binomial Distribution...

$$\begin{aligned}\text{Var}(X) &= E[\underline{X(X-1)}] + E[X] - (E[X])^2 \\ &= n(n-1)p^2 + np - (np)^2 \\ &= np(1-p) \\ &= \end{aligned}$$

» Higher Order Moments...

Define n th moment

The n th moment about the mean or n th central moment of a real valued random variable X is defined as follows

$$\mu_n = E[(X - E[X])^n],$$

where E is the expectation operator.

$E[X]$

$\rightarrow E[X - \mu]$


$E[(X - \mu)^2]$

$E[(X - \mu)^3]$

» Higher Order Moments...

Define n th moment

The n th moment about the mean or n th central moment of a real valued random variable X is defined as follows

$$\mu_n = E[(X - E[X])^n],$$


where E is the expectation operator.

* The zeroth central moment μ_0 is 1

» Higher Order Moments...

Define n th moment

The n th moment about the mean or n th central moment of a real valued random variable X is defined as follows

$$\mu_n = E[(X - E[X])^n],$$

where E is the expectation operator.

$E[(X - \mu)^n]$
 $E[X] - \mu = 0$

- * The **zeroth** central moment μ_0 is 1
- * The **first** central moment μ_1 is 0 (not the same as expected value!)

» Higher Order Moments...

Define n th moment

The n th moment about the mean or n th central moment of a real valued random variable X is defined as follows

$$\mu_n = E[(X - E[X])^n],$$

where E is the expectation operator.

- * The zeroth central moment μ_0 is 1
- * The first central moment μ_1 is 0 (not the same as expected value!)
 - * First central moment about origin is indeed expectation!

» Higher Order Moments...

Define n th moment

The n th moment about the mean or n th central moment of a real valued random variable X is defined as follows

$$\mu_n = E[(X - E[X])^n],$$



Handwritten orange notation: $E[(X - \mu)^n]$. An arrow points from the μ in the original formula to the μ in this handwritten version.

where E is the expectation operator.

- * The zeroth central moment μ_0 is 1
- * The first central moment μ_1 is 0 (not the same as expected value!)
 - * First central moment about origin is indeed expectation!
- * The second central moment μ_2 is called variance, denoted by σ^2

» Higher Order Moments...

Define n th moment

The n th moment about the mean or n th central moment of a real valued random variable X is defined as follows

$$\mu_n = E[(X - E[X])^n],$$

where E is the expectation operator.

- * The zeroth central moment μ_0 is 1
- * The first central moment μ_1 is 0 (not the same as expected value!)
 - * First central moment about origin is indeed expectation!
- * The second central moment μ_2 is called variance, denoted by σ^2

Generating Moments...

Is there a quick way to generate moments?

» Moment Generating Function...

» Moment Generating Function...

$$e^{tX} = \underline{g(x)}$$

Moment Generating Function

The moment generating function $M_X(t)$ is the expectation value

$$M_X(t) = E[e^{tX}] = \sum_x e^{tx} p_X(x)$$

$$\sum p_X(x) = 1$$

Lemma

$$* M_X(0) = 1$$

$$* E[X] = M'_X(0), \text{ where ' is the derivative w.r.t. } t$$

$$\frac{d}{dt} M_X(t) = \sum_x \frac{d}{dt} e^{tx} p_X(x)$$

$$= \sum x p_X(x) = \underline{\mu} = E[X]$$