

# Probability and Statistics: Lecture-6

Monsoon-2020

by Pawan Kumar (IIIT, Hyderabad)

on August 21, 2020

## » Table of contents

### 1. Motivation for Probability

### 2. Digress: Game Theory

## » Computing probabilities...

**Experiment:** Throw two dice.

## » Computing probabilities...

**Experiment:** Throw two dice.

**Event:** Bigger number on red than blue.

## » Computing probabilities...

**Experiment:** Throw two dice.

**Event:** Bigger number on red than blue.

**Quiz:** What is the probability of this event?

## » Computing probabilities...

**Experiment:** Throw two dice.

**Event:** Bigger number on red than blue.

**Quiz:** What is the probability of this event?

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

## » Computing probabilities...

**Experiment:** Throw two dice.

**Event:** Bigger number on red than blue.

**Quiz:** What is the probability of this event?

\* **Probability space:** all outcomes

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

## » Computing probabilities...

**Experiment:** Throw two dice.

**Event:** Bigger number on red than blue.

**Quiz:** What is the probability of this event?

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

- \* **Probability space:** all outcomes
- \* **Event:** some favourable outcomes



## » Computing probabilities...

**Experiment:** Throw two dice.

**Event:** Bigger number on red than blue.

**Quiz:** What is the probability of this event?

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

- \* **Probability space:** all outcomes
- \* **Event:** some favourable outcomes
- \* **Probability:** (#favorable)/36

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

## » Computing probabilities...

**Experiment:** Throw two dice.

**Event:** Bigger number on red than blue.

**Quiz:** What is the probability of this event?

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

Hence, probability,  $p = \frac{15}{36}$

- \* **Probability space:** all outcomes
- \* **Event:** some favourable outcomes
- \* **Probability:** (#favorable)/36

11	12	13	14	15	16
21	22	23	24	25	26
31	32	33	34	35	36
41	42	43	44	45	46
51	52	53	54	55	56
61	62	63	64	65	66

## Quiz

- \* **Experiment:** Toss a coin  $n$  times

## Quiz

- \* **Experiment:** Toss a coin  $n$  times
- \* **Outcome:** sequence of  $n$  bits

## Quiz

- \* **Experiment:** Toss a coin  $n$  times
- \* **Outcome:** sequence of  $n$  bits
- \* **Total number of outcomes:**  $2^n$  outcomes

## Quiz

- \* **Experiment:** Toss a coin  $n$  times
- \* **Outcome:** sequence of  $n$  bits
- \* **Total number of outcomes:**  $2^n$  outcomes
- \* **Assumption:** equiprobable

## Quiz

- \* **Experiment:** Toss a coin  $n$  times
- \* **Outcome:** sequence of  $n$  bits
- \* **Total number of outcomes:**  $2^n$  outcomes
- \* **Assumption:** equiprobable
- \* **Probability of all heads:**  $1/2^n$

## Quiz

- \* **Experiment:** Toss a coin  $n$  times
- \* **Outcome:** sequence of  $n$  bits
- \* **Total number of outcomes:**  $2^n$  outcomes
- \* **Assumption:** equiprobable
- \* **Probability of all heads:**  $1/2^n$
- \* Consider an event of “first bit” = “last bit”



## Quiz

- \* **Experiment:** Toss a coin  $n$  times
- \* **Outcome:** sequence of  $n$  bits
- \* **Total number of outcomes:**  $2^n$  outcomes
- \* **Assumption:** equiprobable
- \* **Probability of all heads:**  $1/2^n$
- \* Consider an event of “first bit” = “last bit”
- \* What is the **probability** of the above event?

» **Answer to Quiz...**

## » Answer to Quiz...

Answer to Quiz:

## » Answer to Quiz...

Answer to Quiz:

\* Number of possible cases:

$$H * \dots * H$$
$$T * \dots * H$$
$$H * \dots * T$$
$$T * \dots * T$$

## » Answer to Quiz...

Answer to Quiz:

- \* Number of possible cases:

$$H * \dots * H$$
$$T * \dots * H$$
$$H * \dots * T$$
$$T * \dots * T$$

- \* **Observation:** In **half** of the cases, first bit does not equal last bit

## » Answer to Quiz...

Answer to Quiz:

- \* Number of possible cases:

$$H * \dots * H$$
$$T * \dots * H$$
$$H * \dots * T$$
$$T * \dots * T$$

- \* **Observation:** In **half** of the cases, first bit does not equal last bit
- \* Hence, probability of the event is  $1/2$

» Remark on Counting. Does probability boils down to counting always?

## » Remark on Counting. Does probability boils down to counting always?

- \* It is true that in many cases probability boils down to counting



## » Remark on Counting. Does probability boils down to counting always?

- \* It is true that in many cases probability boils down to counting
- \* However, there are **non-uniform** distributions

## » Remark on Counting. Does probability boils down to counting always?

- \* It is true that in many cases probability boils down to counting
- \* However, there are **non-uniform** distributions
- \* We also study **continuous** distributions

## » Remark on Counting. Does probability boils down to counting always?

- \* It is true that in many cases probability boils down to counting
- \* However, there are **non-uniform** distributions
- \* We also study **continuous** distributions
- \* In the continuous case, we need to consider some other measure

## » Remark on Counting. Does probability boils down to counting always?

- \* It is true that in many cases probability boils down to counting
- \* However, there are **non-uniform** distributions
- \* We also study **continuous** distributions
- \* In the continuous case, we need to consider some other measure
- \* What is the probability that a number 1.5 is picked in  $[1, 2]$ ?


## » Remark on Counting. Does probability boils down to counting always?

- \* It is true that in many cases probability boils down to counting
- \* However, there are **non-uniform** distributions
- \* We also study **continuous** distributions
- \* In the continuous case, we need to consider some other measure
- \* What is the probability that a number 1.5 is picked in  $[1, 2]$ ? Does this make sense?


## » Probability, Galton Board, and Pascal Triangle

Movie of Galton board here!

## » Probability, Galton Board, and Pascal Triangle




## » Probability, Galton Board, and Pascal Triangle



- \* This is called galton board, **Sir Francis Galton (1822-1911)**




## » Probability, Galton Board, and Pascal Triangle




- \* This is called galton board, **Sir Francis Galton (1822-1911)**
- \* The set of red marbles are allowed to fall

## » Probability, Galton Board, and Pascal Triangle




- \* This is called galton board, **Sir Francis Galton (1822-1911)**
- \* The set of red marbles are allowed to fall
- \* We observe the distribution of the marbles after the fall

## » Probability, Galton Board, and Pascal Triangle




- \* This is called galton board, **Sir Francis Galton (1822-1911)**
- \* The set of red marbles are allowed to fall
- \* We observe the distribution of the marbles after the fall
- \* **Quiz:** Why do we have more marbles in the middle?

## » Probability, Galton Board, and Pascal Triangle




- \* This is called galton board, **Sir Francis Galton (1822-1911)**
- \* The set of red marbles are allowed to fall
- \* We observe the distribution of the marbles after the fall
- \* **Quiz:** Why do we have more marbles in the middle?
- \* Does probability have any role in this distribution?

## » Probability, Galton Board, and Pascal Triangle




- \* This is called galton board, **Sir Francis Galton (1822-1911)**
- \* The set of red marbles are allowed to fall
- \* We observe the distribution of the marbles after the fall
- \* **Quiz:** Why do we have more marbles in the middle?
- \* Does probability have any role in this distribution?
- \* Can we **prove that the distribution will look like this mathematically?**

## » Probability, Galton Board, and Pascal Triangle




- \* This is called galton board, **Sir Francis Galton (1822-1911)**
- \* The set of red marbles are allowed to fall
- \* We observe the distribution of the marbles after the fall
- \* **Quiz:** Why do we have more marbles in the middle?
- \* Does probability have any role in this distribution?
- \* Can we **prove that the distribution will look like this mathematically?**
- \* Does **Pascal triangle** has anything to do with this?

## » Probability, Galton Board, and Pascal Triangle




- \* This is called galton board, **Sir Francis Galton (1822-1911)**
- \* The set of red marbles are allowed to fall
- \* We observe the distribution of the marbles after the fall
- \* **Quiz:** Why do we have more marbles in the middle?
- \* Does probability have any role in this distribution?
- \* Can we **prove that the distribution will look like this mathematically?**
- \* Does **Pascal triangle** has anything to do with this?
- \* Let us analyze this in detail ...

## » Probability, Galton Board, and Pascal Triangle






## » Probability, Galton Board, and Pascal Triangle




\* Assume that at each level the **beams are splitted evenly**

## » Probability, Galton Board, and Pascal Triangle




- \* Assume that at each level the **beams are splitted evenly**
- \* Probability that ball hits current beam is sum of probability of this ball hitting left and right beam above

## » Probability, Galton Board, and Pascal Triangle



- \* Assume that at each level the **beams are splitted evenly**
- \* Probability that ball hits current beam is sum of probability of this ball hitting left and right beam above
- \* If  $x$  is the top left probability, and  $y$  is the top right probability, then probability of hitting the current bin is  $(x + y)/2$


## » Probability, Galton Board, and Pascal Triangle



- \* Assume that at each level the **beams are splitted evenly**
- \* Probability that ball hits current beam is sum of probability of this ball hitting left and right beam above
- \* If  $x$  is the top left probability, and  $y$  is the top right probability, then probability of hitting the current bin is  $(x + y)/2$

		1				
		1/2		1/2		
	1/4		2/4		1/4	
1/8		3/8		3/8		1/8

## » Probability, Galton Board, and Pascal Triangle




- \* Assume that at each level the **beams are splitted evenly**
- \* Probability that ball hits current beam is sum of probability of this ball hitting left and right beam above
- \* If  $x$  is the top left probability, and  $y$  is the top right probability, then probability of hitting the current bin is  $(x + y)/2$

		1				
		1/2		1/2		
	1/4		2/4		1/4	
1/8		3/8		3/8		1/8

- \* The probability of current bin =  $\binom{n}{k}/2^n$


## » Non equiprobable outcomes!

What is wrong with these dice?



## » Non equiprobable outcomes!


What is wrong with these dice?



- \* So far we assumed **equiprobable** outcomes

## » Non equiprobable outcomes!

What is wrong with these dice?



- \* So far we assumed **equiprobable** outcomes
- \* On the left, we have **skewed dice**, i.e., **unfair dice**



## » Non equiprobable outcomes!


What is wrong with these dice?



- \* So far we assumed **equiprobable** outcomes
- \* On the left, we have **skewed dice, i.e., unfair dice**
- \* Will our previous definition of calculating probability work?


## » Non equiprobable outcomes!

What is wrong with these dice?



- \* So far we assumed **equiprobable** outcomes
- \* On the left, we have **skewed dice, i.e., unfair dice**
- \* Will our previous definition of calculating probability work?
- \* Unfortunately not, how to deal with skewed cases?

## » Non equiprobable outcomes!



## » Non equiprobable outcomes!




- \* Since the dice is skewed, frequency of outcomes are **different**

## » Non equiprobable outcomes!




- \* Since the dice is skewed, frequency of outcomes are **different**
  - \* That is, for example, number 4 may appear more frequently

## » Non equiprobable outcomes!




- \* Since the dice is skewed, frequency of outcomes are **different**
  - \* That is, for example, number 4 may appear more frequently
- \* Let  $p_i$  denote the probability of  $i$ th number

## » Non equiprobable outcomes!



- \* Since the dice is skewed, frequency of outcomes are **different**
  - \* That is, for example, number 4 may appear more frequently
- \* Let  $p_i$  denote the probability of  $i$ th number
- \* If we throw the dice long enough, the frequencies **stabilise**


## » Non equiprobable outcomes!



- \* Since the dice is skewed, frequency of outcomes are **different**
  - \* That is, for example, number 4 may appear more frequently
- \* Let  $p_i$  denote the probability of  $i$ th number
- \* If we throw the dice long enough, the frequencies **stabilise**
- \* For example, the probability of getting an even number  
 $= p_2 + p_4 + p_6$



## » Non equiprobable outcomes!



- \* Since the dice is skewed, frequency of outcomes are **different**
  - \* That is, for example, number 4 may appear more frequently
- \* Let  $p_i$  denote the probability of  $i$ th number
- \* If we throw the dice long enough, the frequencies **stabilise**
- \* For example, the probability of getting an even number =  $p_2 + p_4 + p_6$
- \* Also, we know that the sum of **all the probabilities sum to 1, even though the dice is skewed!**
  - \* That is:  $p_1 + p_2 + \dots + p_6 = 1$

## » Example: Dice Throw and Venn Diagram

**Experiment:** Throw a fair dice once

## » Example: Dice Throw and Venn Diagram


**Experiment:** Throw a fair dice once

**Sample Space:**  $S = \{1, 2, 3, 4, 5, 6\}$

## » Example: Dice Throw and Venn Diagram

**Experiment:** Throw a fair dice once

**Sample Space:**  $S = \{1, 2, 3, 4, 5, 6\}$




\* Consider **event**  $A = \{1, 2, 3\}$

## » Example: Dice Throw and Venn Diagram

**Experiment:** Throw a fair dice once

**Sample Space:**  $S = \{1, 2, 3, 4, 5, 6\}$




\* Consider **event**  $A = \{1, 2, 3\}$

\* Consider **event**  $B = \{5, 6\}$

## » Example: Dice Throw and Venn Diagram

**Experiment:** Throw a fair dice once

**Sample Space:**  $S = \{1, 2, 3, 4, 5, 6\}$



\* Since the dice is fair, we have  
 $p_1 + p_2 + \cdots + p_6 = 1$


\* Consider **event**  $A = \{1, 2, 3\}$

\* Consider **event**  $B = \{5, 6\}$

## » Example: Dice Throw and Venn Diagram

**Experiment:** Throw a fair dice once

**Sample Space:**  $S = \{1, 2, 3, 4, 5, 6\}$



\* Consider **event**  $A = \{1, 2, 3\}$

\* Consider **event**  $B = \{5, 6\}$

\* Since the dice is fair, we have


$$p_1 + p_2 + \cdots + p_6 = 1$$

\*  $Pr(A) = p_1 + p_2 + p_3$

## » Example: Dice Throw and Venn Diagram

**Experiment:** Throw a fair dice once

**Sample Space:**  $S = \{1, 2, 3, 4, 5, 6\}$



\* Consider **event**  $A = \{1, 2, 3\}$

\* Consider **event**  $B = \{5, 6\}$

\* Since the dice is fair, we have

$$p_1 + p_2 + \cdots + p_6 = 1$$

$$* Pr(A) = p_1 + p_2 + p_3$$


$$* Pr(B) = p_5 + p_6$$



## » Example: Dice Throw and Venn Diagram

**Experiment:** Throw a fair dice once

**Sample Space:**  $S = \{1, 2, 3, 4, 5, 6\}$



\* Consider **event**  $A = \{1, 2, 3\}$

\* Consider **event**  $B = \{5, 6\}$

\* Since the dice is fair, we have

$$p_1 + p_2 + \cdots + p_6 = 1$$

$$* Pr(A) = p_1 + p_2 + p_3$$


$$* Pr(B) = p_5 + p_6$$

\* Consider the **event**  $A$  or  $B = \{1, 2, 3, 5, 6\}$

## » Example: Dice Throw and Venn Diagram

**Experiment:** Throw a fair dice once

**Sample Space:**  $S = \{1, 2, 3, 4, 5, 6\}$



- \* Consider **event**  $A = \{1, 2, 3\}$
- \* Consider **event**  $B = \{5, 6\}$

- \* Since the dice is fair, we have

$$p_1 + p_2 + \cdots + p_6 = 1$$

- \*  $Pr(A) = p_1 + p_2 + p_3$

- \*  $Pr(B) = p_5 + p_6$

- \* Consider the **event**  $A$  or  $B = \{1, 2, 3, 5, 6\}$

- \*  $Pr(A \text{ or } B) =$


$$Pr(A) + Pr(B) = p_1 + p_2 + p_3 + p_5 + p_6$$

**Why  $Pr(A \text{ or } B) = Pr(A) + Pr(B)$ ?**

## » Example: Dice Throw and Venn Diagram

**Experiment:** Throw a fair dice once

**Sample Space:**  $S = \{1, 2, 3, 4, 5, 6\}$



- \* Consider **event**  $A = \{1, 2, 3\}$
- \* Consider **event**  $B = \{5, 6\}$

- \* Since the dice is fair, we have

$$p_1 + p_2 + \cdots + p_6 = 1$$

- \*  $Pr(A) = p_1 + p_2 + p_3$

- \*  $Pr(B) = p_5 + p_6$

- \* Consider the **event**  $A$  or  $B = \{1, 2, 3, 5, 6\}$

- \*  $Pr(A \text{ or } B) =$

$$Pr(A) + Pr(B) = p_1 + p_2 + p_3 + p_5 + p_6$$

**Why  $Pr(A \text{ or } B) = Pr(A) + Pr(B)$ ?**

**When are we allowed to add probabilities?**

## » Mutually Exclusive Events Events

### Mutually Exclusive Events

Two events are said to be **mutually exclusive** if they have disjoint set of outcomes.

## » Mutually Exclusive Events

### Mutually Exclusive Events

Two events are said to be **mutually exclusive** if they have disjoint set of outcomes. Let  $A$  and  $B$  be two **mutually exclusive events**, then we have

$$Pr(A \cup B) = Pr(A) + Pr(B)$$

## » Mutually Exclusive Events

### Mutually Exclusive Events

Two events are said to be **mutually exclusive** if they have disjoint set of outcomes. Let  $A$  and  $B$  be two **mutually exclusive events**, then we have

$$Pr(A \cup B) = Pr(A) + Pr(B)$$

### Probability of a complement

Let  $A$  be any event, and  $A^c$  be its **complement**.

## » Mutually Exclusive Events

### Mutually Exclusive Events

Two events are said to be **mutually exclusive** if they have disjoint set of outcomes. Let  $A$  and  $B$  be two **mutually exclusive events**, then we have

$$Pr(A \cup B) = Pr(A) + Pr(B)$$

### Probability of a complement

Let  $A$  be any event, and  $A^c$  be its **complement**. Then clearly  $A \cup A^c$  is the **complete set of outcomes**, i.e., it is a **sample space**.

## » Mutually Exclusive Events

### Mutually Exclusive Events

Two events are said to be **mutually exclusive** if they have disjoint set of outcomes. Let  $A$  and  $B$  be two **mutually exclusive events**, then we have

$$Pr(A \cup B) = Pr(A) + Pr(B)$$

### Probability of a complement

Let  $A$  be any event, and  $A^c$  be its **complement**. Then clearly  $A \cup A^c$  is the **complete set of outcomes**, i.e., it is a **sample space**. Hence, we have

$$Pr(A) + Pr(A^c) = 1 \quad \Rightarrow \quad Pr(A^c) = 1 - Pr(A)$$



## » Mutually Exclusive Events..

## » Mutually Exclusive Events..

### Mutually disjoint events

A set of  $n$  events denoted by  $A_1, A_2, \dots, A_n$  are called **mutually disjoint events** if the following holds:

## » Mutually Exclusive Events..

### Mutually disjoint events

A set of  $n$  events denoted by  $A_1, A_2, \dots, A_n$  are called **mutually disjoint events** if the following holds:

$$P(A_1 \cup \dots \cup A_n) = P(A_1) + \dots + P(A_n).$$

## » Mutually Exclusive Events..

### Mutually disjoint events

A set of  $n$  events denoted by  $A_1, A_2, \dots, A_n$  are called **mutually disjoint events** if the following holds:

$$P(A_1 \cup \dots \cup A_n) = P(A_1) + \dots + P(A_n).$$

From **Axiom-3**, it holds for **countably infinite unions**.

## » Quiz: Use concept of disjoint events

### Quiz

In a cricket tournament with four teams denoted by  $\{A, B, C, D\}$ , team  $A$  has 20% chance of winning, while team  $B$  has a 40% chance of winning. What is the probability that  $A$  or  $B$  win the tournament?

## » Quiz-Use probability axioms

### Quiz

Suppose we know the following:

- \* there is a 50% chance that it will be hot Today
- \* there is a 30% chance that it will be hot Tomorrow
- \* there is 10% chance that it wont be hot Today or Tomorrow

## » Quiz-Use probability axioms

### Quiz

Suppose we know the following:

- \* there is a 50% chance that it will be hot Today
- \* there is a 30% chance that it will be hot Tomorrow
- \* there is 10% chance that it wont be hot Today or Tomorrow

Answer the following:

## » Quiz-Use probability axioms

### Quiz

Suppose we know the following:

- \* there is a 50% chance that it will be hot Today
- \* there is a 30% chance that it will be hot Tomorrow
- \* there is 10% chance that it wont be hot Today or Tomorrow

Answer the following:

- \* probability that it will be hot today or tomorrow
- \* probability that it will be hot today and tomorrow
- \* probability that it will be hot today but not tomorrow
- \* probability that it either will be hot today or tomorrow, but not both



## » Non-Mutually Exclusive Events

- \* What if the two events  $A$  and  $B$  are **not mutually exclusive**?

## » Non-Mutually Exclusive Events

- \* What if the two events  $A$  and  $B$  are **not mutually exclusive**?
- \* What relation we have


$$Pr(A \cup B) \quad ? \quad Pr(A) + Pr(B) \quad (1)$$

## » Non-Mutually Exclusive Events

- \* What if the two events  $A$  and  $B$  are **not mutually exclusive**?
- \* What relation we have

$$Pr(A \cup B) \quad ? \quad Pr(A) + Pr(B) \quad (1)$$

- \* Let us consider an example below with two events  $A$  and  $B$  shaded




## » Non-Mutually Exclusive Events

- \* What if the two events  $A$  and  $B$  are **not mutually exclusive**?
- \* What relation we have


$$Pr(A \cup B) \quad ? \quad Pr(A) + Pr(B) \quad (1)$$

- \* Let us consider an example below with two events  $A$  and  $B$  shaded




- \* For the events  $A$  and  $B$ , which relation (1) holds?

\* Consider the example as before




- \* Consider the example as before



- \* Let us calculate the probabilities of the two events  $A$  and  $B$

$$Pr(A) = p_1 + p_2 + p_3, \quad Pr(B) = p_3 + p_4$$

- \* Consider the example as before




- \* Let us calculate the probabilities of the two events  $A$  and  $B$

$$Pr(A) = p_1 + p_2 + p_3, \quad Pr(B) = p_3 + p_4$$

- \* Now we calculate probability of  $A \cup B$

- \* Consider the example as before



- \* Let us calculate the probabilities of the two events  $A$  and  $B$


$$Pr(A) = p_1 + p_2 + p_3, \quad Pr(B) = p_3 + p_4$$

- \* Now we calculate probability of  $A \cup B$

$$Pr(A \cup B) = Pr(\{1, 2, 3, 4\}) = p_1 + p_2 + p_3 + p_4$$



- \* Consider the example as before



- \* Let us calculate the probabilities of the two events  $A$  and  $B$


$$Pr(A) = p_1 + p_2 + p_3, \quad Pr(B) = p_3 + p_4$$

- \* Now we calculate probability of  $A \cup B$

$$Pr(A \cup B) = Pr(\{1, 2, 3, 4\}) = p_1 + p_2 + p_3 + p_4$$

- \* Using above:  $Pr(A \cup B) = Pr(A) + Pr(B) - Pr(A \cap B)$

- \* Consider the example as before



- \* Let us calculate the probabilities of the two events  $A$  and  $B$

$$Pr(A) = p_1 + p_2 + p_3, \quad Pr(B) = p_3 + p_4$$

- \* Now we calculate probability of  $A \cup B$

$$Pr(A \cup B) = Pr(\{1, 2, 3, 4\}) = p_1 + p_2 + p_3 + p_4$$

- \* Using above:  $Pr(A \cup B) = Pr(A) + Pr(B) - Pr(A \cap B)$
- \* Recall discrete mathematics: **inclusion-exclusion principle**. (Proof?)

## » Generalized Inclusion-Exclusion Principle

## » Generalized Inclusion-Exclusion Principle

### Inclusion-Exclusion Principle

$$* P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

## » Generalized Inclusion-Exclusion Principle

### Inclusion-Exclusion Principle

- \*  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

- \*  $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$

## » Generalized Inclusion-Exclusion Principle

### Inclusion-Exclusion Principle

- \*  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
- \*  $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$
- \* In general for  $n$  events, we have

## » Generalized Inclusion-Exclusion Principle

### Inclusion-Exclusion Principle

- \*  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$
- \*  $P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(B \cap C) - P(A \cap C) + P(A \cap B \cap C)$
- \* In general for  $n$  events, we have

$$P(\cup_{i=1}^n A_i) = \sum_{i=1}^n P(A_i) - \sum_{i < j} P(A_i \cap A_j) + \sum_{i < j < k} P(A_i \cap A_j \cap A_k) \\ - \dots (-1)^{n-1} P(\cap_{i=1}^n A_i)$$





## » Discrete Probability Models

- \* Sample space can be of two types: **discrete or continuous**

## » Discrete Probability Models

- \* Sample space can be of two types: **discrete or continuous**
- \* in discrete probability models we can compute the probability of events by **adding** all the corresponding outcomes

## » Discrete Probability Models

- \* Sample space can be of two types: **discrete or continuous**
- \* in discrete probability models we can compute the probability of events by **adding** all the corresponding outcomes
- \* in continuous probability models we need to use **integration instead of summation**

## » Discrete Probability Models

- \* Sample space can be of two types: **discrete or continuous**
- \* in discrete probability models we can compute the probability of events by **adding** all the corresponding outcomes
- \* in continuous probability models we need to use **integration instead of summation**

### Definition of discrete probability model

If the sample space  $S$  is a countable set, then it refers to discrete probability model. Since  $S$  is countable:  $S = \{s_1, s_2, \dots, \}$

## » Discrete Probability Models

- \* Sample space can be of two types: **discrete or continuous**
- \* in discrete probability models we can compute the probability of events by **adding** all the corresponding outcomes
- \* in continuous probability models we need to use **integration instead of summation**

### Definition of discrete probability model

If the sample space  $S$  is a countable set, then it refers to discrete probability model. Since  $S$  is countable:  $S = \{s_1, s_2, \dots, \}$

- \* For an event  $A \subset S$ , by **3rd axiom**

$$P(A) = \sum_{s_j \in A} P(s_j)$$



## » Choices and Choice Tree

- \* Consider 6 balls labelled 1,2,3,4,5,6


## » Choices and Choice Tree

- \* Consider 6 balls labelled 1,2,3,4,5,6
- \* The balls are kept in two boxes. In the first box, we have 1, 2 balls, and in the second box, we have 3, 4, 5, 6 balls




## » Choices and Choice Tree

- \* Consider 6 balls labelled 1,2,3,4,5,6
- \* The balls are kept in two boxes. In the first box, we have 1, 2 balls, and in the second box, we have 3, 4, 5, 6 balls



## » Choices and Choice Tree


- \* Consider 6 balls labelled 1,2,3,4,5,6
- \* The balls are kept in two boxes. In the first box, we have 1, 2 balls, and in the second box, we have 3, 4, 5, 6 balls



- \* Assume the following:

## » Choices and Choice Tree


- \* Consider 6 balls labelled 1,2,3,4,5,6
- \* The balls are kept in two boxes. In the first box, we have 1, 2 balls, and in the second box, we have 3, 4, 5, 6 balls



- \* Assume the following:
  - \* The two boxes are **equiprobable**

## » Choices and Choice Tree


- \* Consider 6 balls labelled 1,2,3,4,5,6
- \* The balls are kept in two boxes. In the first box, we have 1, 2 balls, and in the second box, we have 3, 4, 5, 6 balls




- \* Assume the following:
  - \* The two boxes are **equiprobable**
  - \* All the balls in each box is **equiprobable**

## » Choices and Choice Tree


- \* Consider 6 balls labelled 1,2,3,4,5,6
- \* The balls are kept in two boxes. In the first box, we have 1, 2 balls, and in the second box, we have 3, 4, 5, 6 balls



- \* Assume the following:
  - \* The two boxes are **equiprobable**
  - \* All the balls in each box is **equiprobable**
- \* **Question:** What are the probabilities  $p_1, p_2, \dots, p_6$ ?




Consider a choice tree for the problem



\* We have the following

$$p_1 + p_2 = 1/2, \quad p_3 + p_4 + p_5 + p_6 = 1/2$$

Consider a choice tree for the problem




Consider a choice tree for the problem

\* We have the following

$$p_1 + p_2 = 1/2, \quad p_3 + p_4 + p_5 + p_6 = 1/2$$

\* Balls in box-1 are **equiprobable**:  $p_1 = p_2$






Consider a choice tree for the problem

\* We have the following

$$p_1 + p_2 = 1/2, \quad p_3 + p_4 + p_5 + p_6 = 1/2$$

- \* Balls in box-1 are **equiprobable**:  $p_1 = p_2$
- \* Balls in box-2 are **equiprobable**:  $p_3 = p_4 = p_5 = p_6$



Consider a choice tree for the problem

\* We have the following

$$p_1 + p_2 = 1/2, \quad p_3 + p_4 + p_5 + p_6 = 1/2$$

- \* Balls in box-1 are **equiprobable**:  $p_1 = p_2$
- \* Balls in box-2 are **equiprobable**:  $p_3 = p_4 = p_5 = p_6$
- \* From above, we have

$$p_1 = p_2 = 1/4, \quad p_3 = p_4 = p_5 = p_6 = 1/8$$

## » Shrewd Prisoner and Maximizing Chance!

### Shrewd Prisoner Problem

Long ago a prisoner was to be executed.

## » Shrewd Prisoner and Maximizing Chance!

### Shrewd Prisoner Problem

Long ago a prisoner was to be executed. In response to his supplications, he was promised that he would be released if he drew a white ball from one of two similar urns.

## » Shrewd Prisoner and Maximizing Chance!

### Shrewd Prisoner Problem

Long ago a prisoner was to be executed. In response to his supplications, he was promised that he would be released if he drew a white ball from one of two similar urns. The provisions were that he had to distribute 50 white and 50 black balls between the two urns,

## » Shrewd Prisoner and Maximizing Chance!

### Shrewd Prisoner Problem

Long ago a prisoner was to be executed. In response to his supplications, he was promised that he would be released if he drew a white ball from one of two similar urns. The provisions were that he had to distribute 50 white and 50 black balls between the two urns, in any way he liked,

## » Shrewd Prisoner and Maximizing Chance!

### Shrewd Prisoner Problem

Long ago a prisoner was to be executed. In response to his supplications, he was promised that he would be released if he drew a white ball from one of two similar urns. The provisions were that he had to distribute 50 white and 50 black balls between the two urns, in any way he liked, after which he had to draw a ball at random from one of these urns.

## » Shrewd Prisoner and Maximizing Chance!

### Shrewd Prisoner Problem

Long ago a prisoner was to be executed. In response to his supplications, he was promised that he would be released if he drew a white ball from one of two similar urns. The provisions were that he had to distribute 50 white and 50 black balls between the two urns, in any way he liked, after which he had to draw a ball at random from one of these urns.

**Question:** How should the prisoner put the balls such that the probability of his release is maximized?