

Baby Sleep Training Methods Compared Cheat Sheet

14 tables · 80 concepts · Updated 2026-07-28 · cheatgrid.com

Baby sleep training encompasses a spectrum of evidence-based and widely-used approaches that help infants learn to fall asleep independently and consolidate nighttime sleep. From graduated check-in methods to fully parent-present gentle strategies, each approach differs in how much crying is involved, parental presence, and the age at which it is appropriate. This cheat sheet covers all major sleep training methods, age-appropriate sleep needs, wake windows, environmental setup, sleep regressions, troubleshooting, and special scenarios — giving caregivers the complete picture to choose, start, and sustain a sleep training approach with confidence.

CONTENTS

Sleep Training Method Comparison: Core Approaches	1
When to Start Sleep Training — Readiness Criteria	3
Sleep Needs by Age (AASM/AAP Guidelines)	3
Wake Windows by Age	4
Overtired vs. Undertired Signs	5
The Drowsy-But-Awake Principle	6
Sleep Environment Setup (Room, Darkness, White Noise, Temperature)	6
Naps vs. Nights - Training Order and Strategy	7
Sleep Regressions and How to Handle Them	8
Night Weaning	9
Early Morning Waking Troubleshooting	10
Sleep Training Twins and Shared Rooms	10
When Sleep Training Is Not Working	11
Special Scenarios and Advanced Topics	12
References	13

TABLE 1

Sleep Training Method Comparison: Core Approaches

All major sleep training methods share the same goal, teaching a baby to fall asleep independently, but they differ dramatically in parental presence, tolerated crying level, and typical time to results. Methods range from full extinction, with no check-ins and the fastest results, to fully parent-present approaches, which are the most gradual. Choose based on your family's emotional tolerance, your baby's temperament, and your pediatrician's guidance.

Technique/Method	Example	Description
Ferber Method (Graduated Extinction)	Night 1: wait 3 min → comfort briefly → wait 5 min → comfort → wait 10 min; Night 2: 5/10/12 min	• Timed check-ins with progressively longer wait intervals • Parent enters briefly (1-2 min) to reassure verbally, no picking up • Intervals increase each night up to Day 7: 20/25/30 min • Well studied graduated-extinction protocol; results typically in 3-7 nights
Weissbluth Method (Full Extinction / Cry It Out)	Put baby down drowsy-but-awake at 6-8 pm; leave room; do not return until morning wake time	• No check-ins after initial bedtime routine; baby is left to fuss or cry without a time limit until morning • Often works fastest (2-4 nights) but requires high parental tolerance • Dr. Marc Weissbluth recommends starting around 4-6 months • The strongest randomized-trial evidence covers graduated extinction and bedtime fading specifically; these same trials found no long-term harm to attachment or child behavior, and that evidence is generally taken to support full extinction as well

Technique/Method	Example	Description
Sleep Lady Shuffle (Kim West)	Night 1-3: chair beside crib; Night 4-6: chair halfway to door; Night 7-9: chair at doorway; Night 10-12: outside door	• Parent stays in room and moves chair further away every few nights • Verbal and brief physical reassurance permitted • Gentle approach; some crying expected but hysterical crying avoided • Works 6 months to 6 years; Kim West's own site reports many families see a real change within 7-10 days, though it can take longer if the routine needs adjusting
Pick Up Put Down / PUPD (Tracy Hogg)	Baby cries → pick up, soothe until calm (not asleep) → put down; repeat until asleep in crib	• Parent responds immediately but does not allow baby to fall asleep in arms • Works best for 4-8 months; can over-stimulate younger babies • Labor-intensive method; may take 45-90 min early nights • Teaches self-soothing without prolonged crying
No-Cry Sleep Solution (Elizabeth Pantley)	Dream feed at 11 pm; Pantley Pull-Off: remove breast/bottle at drowsy moment each time; gradual weaning over 4+ weeks	• Zero crying philosophy; works through gradual habit replacement • Pantley Pull-Off breaks feed-to-sleep association incrementally • Slowest method (4-8 weeks); most compatible with attachment parenting • Works from birth; no minimum age
Sleepeasy Solution (Waldburger & Spivack)	Check at 5 min intervals first night; speak calmly to reassure; increase intervals each night	• Graduated extinction with emotional support emphasis, grouped with Ferber as a gradual, check-in based approach rather than a no-check-in one • Includes detailed nap and schedule coaching • Results in as little as 3-7 nights; designed for infants around 5 months and up
Babywise (Ezzo & Bucknam)	Feed every 2.5-3 hr; Eat-Wake-Sleep cycle throughout day; independent sleep from nap 1	• Scheduled feed-wake-sleep cycle used to prevent feed-to-sleep association • Pediatric groups, including an AAP district chapter, have raised cautions that a rigid version of this schedule has been associated in some reported cases with poor weight gain and reduced milk supply • Works best when feeding cues are still met within the schedule • Independent sleep often achieved by 8-12 weeks
Chair Method / Gradual Withdrawal	Parent sits in chair next to crib; provides minimal verbal reassurance; moves chair 1-2 feet further each night	• Slower graduated-extinction variant; parent presence is the key comfort, though the parent does not pick the baby up • Similar to Sleep Lady Shuffle but without formal 3-night intervals • Low-cry for most babies; takes 2-4 weeks • Well-suited for highly parent-attached babies, though the parent's visible presence can itself confuse or distract some babies

Topic edits

- Description: rewritten (was: "share the same goal, teaching a baby to fall asleep independently, joined by em-dashes rather than commas"), kept the same content and meaning but replaced em-dashes with commas per house style, since content otherwise already specific and learning-focused
- Rows added: none
- Rows removed: none
- Rows reordered: no
- Column 1 link updates: Weissbluth Method (Full Extinction / Cry It Out): <https://www.healthysleepassociation.org/articles/cry-it-out-sleep-training> → <https://marcweissbluth.com/25-let-cry-sleep-solutions-graduated-extinction-extinction-sleep-solutions-3/> (Weissbluth's own official site, verified with the actual Extinction/Graduated Extinction protocol text); Pick Up Put Down / PUPD (Tracy Hogg): https://www.babycenter.com/baby/sleep/pick-up-put-down-sleep-training_10334349 → <https://health.clevelandclinic.org/pick-up-put-down-method> (babycenter.com is on this domain's avoid list; replaced with a verified Tier 2 clinical source); No-Cry Sleep Solution (Elizabeth Pantley): <https://www.pantley.com/elizabeth/books/the-no-cry-sleep-solution/> → <https://www.pantley.com/> (old link returns a 404, confirmed dead; replaced with the author's live official site); Sleepeasy Solution (Waldburger & Spivack): <https://www.sleepeasybaby.com> → <https://www.sleepyplanet.com/> (old domain no longer resolves, confirmed dead; replaced with the authors' current live official site); Chair Method / Gradual Withdrawal: <https://www.healthline.com/health/baby/chair-method-sleep-training> → <https://health.clevelandclinic.org/chair-method-sleep-training> (healthline.com is on this domain's avoid list absent a named MD/RD byline; replaced with a verified Tier 2 clinical source)
- Row description rewrites: Weissbluth Method (Full Extinction / Cry It Out) (softened the "strong evidence base" claim; the cited RCTs directly test graduated extinction and bedtime fading, not literally unmodified full extinction), Sleep Lady Shuffle (Kim West) (corrected the "takes 2-3 weeks" claim; the method's own official site states most families see results within 7-10 days), Sleepeasy Solution (Waldburger & Spivack) (removed the unverified "check-ins from doorway, not bedside" detail, which no reputable source confirmed, and replaced it with the verified "gradual, check-in based approach" framing), Babywise (Ezzo & Bucknam) (tightened the AAP caution language to precisely

match what AAP News actually reported, an association raised by pediatricians and a district-level resolution, rather than implying a single blanket AAP policy statement)

- Duplicates merged: none
- Deprecated demotions: none

TABLE 2

When to Start Sleep Training — Readiness Criteria

Starting sleep training before a baby is developmentally and physically ready can be ineffective and stressful. Pediatricians and sleep specialists agree on a core set of readiness milestones before any method is introduced. Always consult your pediatrician before beginning, especially for premature infants or babies with medical conditions.

Technique/Method	Example	Description
Minimum Age for Sleep Training	4-6 months corrected age for term infants; premature babies calculated from due date	• No structured method recommended before about 4 months corrected age • 6 months is a commonly cited ideal starting point for many families • Earlier than 4 months: circadian rhythm and sleep cycle maturity are not developed enough
Weight Threshold Guideline	Around 12-14 lbs (5.4-6.3 kg), along with steady weight gain on the baby's own growth curve	• Suggests the baby can usually go without an overnight feeding • Pediatricians cite slightly different numbers within the 12-14 lb range • Premature or low-birthweight babies need individualized guidance from their pediatrician
Pediatrician Sign-Off	Discuss at the 4-month well-child visit; rule out reflux, sleep apnea, or feeding concerns as causes of night waking	• Rules out medical causes of night waking, like reflux or apnea, before assuming it's just a habit • Confirms the baby is growing, feeding, and gaining weight well • Special situations (reflux, colic, prematurity) may need a modified approach
Developmental Signs of Readiness	Some predictable wake and nap pattern, plus early signs of day and night sleep distinction (typically 3-6 months)	• Circadian rhythm (the body's internal sleep clock) begins maturing around 3-4 months • Baby shows recognizable tired cues rather than only overstimulated crying • Day and night sleep starts to separate, usually settling in by 4-6 months
Parental Readiness	Both caregivers agree on one approach and can stay consistent for at least the next one to two weeks	• Consistency across every caregiver is the factor most linked to sleep training success • Better to start during a stable stretch, not right before travel, illness, or another big change • Every caregiver on duty overnight needs to follow the same plan

TABLE 3

Sleep Needs by Age (AASM/AAP Guidelines)

Understanding how much sleep a baby actually needs at each age prevents both over-tiredness (putting down too late) and under-tiredness (putting down too early). The American Academy of Sleep Medicine and AAP recommend total 24-hour sleep including naps; these are norms, not prescriptions — individual variation of 30-60 minutes either side is normal.

Technique/Method	Example	Description
Newborn Sleep Needs (0-3 months)	16-18 total hours in 24h; naps every 45-90 min; no structured schedule possible	• Polyphasic sleep — no day/night differentiation until ~6-8 weeks • Sleep cycles 45-50 min (short active sleep dominates) • Night stretches of 3-5 hr emerge around 6-8 weeks
Infant Sleep Needs (4-6 months)	14-16 total hours; 3-4 naps; first longer night stretch 5-7 hr possible	• Circadian rhythm forming; melatonin production starting • Sleep architecture shifting toward adult pattern (more REM at end of night) • 4-month sleep regression peaks in this window
Infant Sleep Needs (6-9 months)	12-15 total hours; 2-3 naps; 10-12 hr night typical	• Most babies biologically capable of sleeping 11-12 hr with no feeds • 3→2 nap transition often happens 7-9 months • Night waking at this age is typically habit, not hunger

Technique/Method	Example	Description
Infant Sleep Needs (9-12 months)	12-14 total hours; 2 naps; 11-12 hr night	• 8-month and 9-12 month sleep regressions common • Object permanence development drives separation anxiety and night calling • Most babies can consolidate to 2 naps by 9 months
Toddler Sleep Needs (12-24 months)	11-14 total hours; 1 nap (90-120 min); 11-12 hr night	• 2→1 nap transition typically 12-18 months • 18-month and 2-year sleep regressions driven by developmental leaps • Total sleep below 10 hours associated with behavioral and developmental impacts
Toddler Sleep Needs (2-3 years)	11-13 total hours; 1 nap (fading after age 3); 11-12 hr night	• Some 2-year-olds drop nap; most still benefit until age 3 • Quiet time (30-60 min) even without sleep supports learning and mood • Under-sleep at this age strongly linked to increased behavioral dysregulation

TABLE 4

Wake Windows by Age

Wake windows are the amount of time a baby can comfortably stay awake between sleep periods, and they are one of the most useful tools for spotting the line between overtired and undertired. Following wake windows instead of a fixed clock schedule lets parents respond to their own baby's pace instead of someone else's average. Getting them right tends to shorten the time it takes to fall asleep and can noticeably improve how well a baby naps and sleeps at night.

Technique/Method	Example	Description
Wake Windows 4-10 Weeks	60-90 minutes max awake time; watch for 45-min mark tired signs	• Newborns tire extremely quickly; overstimulation leads to harder settling • Wake window includes feeding, diaper change, brief awake time • First morning wake window is often shortest
Wake Windows 10-14 Weeks	75-90 minutes; last wake window before bed may stretch to 100-110 min	• Wakeful periods lengthening as visual and social development accelerates • 3-4 naps typical • Sleep consolidation beginning overnight
Wake Windows 3-5 Months	90-120 minutes; 3 naps needed; last wake window often 120 min	• 4-month regression hits in this window • Wake windows extending rapidly; daily adjustment common • Circadian rhythm strengthening, making a 7-8 pm bedtime appropriate now
Wake Windows 5-7 Months	2-2.5 hours; 3 naps to 2 naps transition begins	• Third nap often short (30-45 min catnap) in late afternoon • Two-nap schedule (AM and PM) emerging • Last wake window before bed 2.5-3 hours
Wake Windows 7-14 Months	3-4 hours between sleep; firmly on 2 naps	• 8-month, 9-month, 10-month mini-regressions overlap with this window • Last wake window before bed 3.5-4 hours typical • Early transition to 1 nap before 12 months usually backfires
Wake Windows 14-24 Months	5-6 hours between sleep; 1 nap at midday after 2 to 1 transition	• 2 to 1 nap transition 14-18 months for most toddlers • Nap at 12-1 pm; bedtime 7-8 pm • Overtiredness from premature nap drop causes night waking and early rising

Topic edits

- Description: rewritten (was: "Wake windows — the amount of time a baby can comfortably...")
- Rows added: none
- Rows removed: none
- Rows reordered: no
- Column 1 link updates: Wake Windows 4-10 Weeks: <https://www.taking-cara-babies.com/wake-windows/> (dead domain, does not resolve) → <https://huckleberrycare.com/blog/first-year-of-sleep-expectations>, Wake Windows 10-14 Weeks: <https://www.taking-cara-babies.com/wake-windows/> (dead domain) → <https://huckleberrycare.com/blog/first-year-of-sleep-expectations>, Wake Windows 3-5 Months: <https://www.babysleepscience.com/single-post/2015/03/31/baby-wake-windows> (404) → <https://huckleberrycare.com/blog/first-year-of-sleep-expectations>, Wake Windows 5-7 Months: <https://www.huckleberrycare.com/blog/baby-wake-windows> (404) → <https://huckleberrycare.com/blog/3-to-2-nap-transition>, Wake Windows 7-14 Months: <https://www.huckleberrycare.com/blog/baby-wake-windows> (404) → <https://huckleberrycare.com/blog/nap-transitions-when-they-occur-and-how-to-handle-them>, Wake Windows 14-24 Months: <https://www.huckleberrycare.com/blog/baby-wake-windows> (404) → <https://huckleberrycare.com/blog/2-to-1-nap-transition>

- Row description rewrites: Wake Windows 3-5 Months (removed em dash from the circadian rhythm bullet)
- Row example rewrites: none
- Duplicates merged: none
- Deprecated demotions: none

TABLE 5

Overtired vs. Undertired Signs

One of the most common errors in baby sleep is misreading tired signs. Overtired babies have elevated cortisol and are harder to settle; undertired babies don't have sufficient sleep pressure. Identifying which state a baby is in allows caregivers to time sleep correctly and avoid the "second wind" trap.

Technique/Method	Example	Description
Early Tired Signs (Optimal Sleep Window)	Staring/zoning out, decreased activity, yawning 1-2 times, pulling at ears	• This is the ideal window to start sleep routine • Eyes may go slightly glassy; social engagement decreasing • Catching this window = fastest sleep onset and calmer settling
Overtired Signs	Arching back, second burst of energy/"wired" behavior, rubbing face intensely, inconsolable crying, yawning repeatedly	• Cortisol and adrenaline spike when overtired, causing paradoxical energy • Settling takes significantly longer; sleep is lighter and more fragmented • Overtired babies resist comfort and cry instead of settling, unlike a simply sleepy baby
Undertired Signs	Bouncy/playful at bedtime, talking/babbling in crib, awake for 60+ min at bedtime without distress	• Sleep pressure insufficient; wake window was too short • Solution: extend wake window for next sleep, not an earlier bedtime • Undertired babies also wake early from naps and nights (partial arousal without returning to sleep)
Cortisol Second Wind	Baby's crying gets more intense the longer you try to settle them; each soothing attempt seems to make it harder, not easier	• Stress hormones (cortisol/adrenaline) create hyperarousal despite fatigue • Overtired babies often wake more at night, not less • Prevention: protecting a steady, consistent bedtime rather than pushing it later breaks the cycle
Age-Specific Tired Sign Differences	Newborn: pulling at ears, closing fists, or arching backwards; 6 to 12 months: clumsiness, grizzling, or increased activity; 12+ months: clinginess or demands for attention	• Tired sign "vocabulary" changes with developmental stage • Older babies mask tired signs with stimulation seeking behavior like increased activity, not obvious slowing down • Consistent nap/bedtime routine helps baby's body clock predict sleep need

Topic edits

- Description: kept-as-is (already specific to this table's scope and learning-focused, no rewrite needed)
- Rows added: none
- Rows removed: none
- Rows reordered: no
- Column 1 link updates: Early Tired Signs (Optimal Sleep Window): <https://www.healthychildren.org/English/ages-stages/baby/sleep/Pages/Signs-of-Tired-Babies.aspx> (dead, 404) → <https://health.clevelandclinic.org/wake-windows-by-age>, Overtired Signs: <https://www.happiestbaby.com/blogs/baby/overtired-baby> (avoid-list domain) → <https://pathways.org/sleep-cues-to-tell-if-baby-is-sleepy>, Undertired Signs: <https://www.babysleepsite.com/sleep-training/is-baby-not-tired-enough/> (avoid-list domain) → <https://littlesleepcoach.com/is-baby-overtired-or-under-tired/>, Cortisol Second Wind: <https://www.preciouslittlesleep.com/are-you-accidentally-keeping-your-baby-overtired/> (avoid-list domain) → <https://www.earlylearningsociety.org/blog/Why-Wont-My-Toddler-Sleep-Understanding-and-Navigating-Bedtime-Resistance-in-Two-Year-Olds.htm>, Age-Specific Tired Sign Differences: <https://www.babysleepsite.com/baby-naps-2/all-about-tired-signs/> (avoid-list domain) → <https://www.firstfiveyears.org.au/child-development/parenting-techniques/tired-signs-in-babies-and-toddlers>
- Row description rewrites: Overtired Signs (removed an em dash and tightened the cortisol bullet, added a third bullet naming the behavioral contrast with simply sleepy), Cortisol Second Wind (removed an unverifiable specific clock-time recommendation, replaced with the verified "consistent bedtime, not later" fix)

- Row example rewrites: Cortisol Second Wind (removed an unverifiable specific 30 minute figure, reworded to the verified general pattern), Age-Specific Tired Sign Differences (replaced an unsupported "frantic rooting" claim with the source's actual newborn and older-baby sign lists)
- Duplicates merged: none
- Deprecated demotions: none

TABLE 6

The Drowsy-But-Awake Principle

Drowsy-but-awake is the foundational skill underlying every sleep training method. When a baby is placed in their sleep environment while still conscious (even slightly), they develop the ability to transition from light sleep back to deep sleep independently, the core mechanism behind "sleeping through the night."

Technique/Method	Example	Description
Drowsy-But-Awake Definition	Eyes at half-mast, body relaxed, head drooping but eyes still partially open when placed in crib	• Baby is sleepy but not fully asleep at the time of crib placement • This lets the baby complete the last stretch of falling asleep independently • Essential prerequisite for self soothing development
Sleep Associations (Negative)	Feeding to sleep, rocking to sleep, pacifier replacement required, being held in arms	• Any external prop present at sleep onset becomes something the baby needs recreated at every sleep cycle transition • Baby stirs at the end of each short sleep cycle and calls out for the prop to fall back asleep • A common cause of frequent night waking in infants past the newborn stage
Sleep Associations (Positive)	White noise machine (plays all night), dark room, sleep sack, consistent bedtime routine	• Environmental cues present at sleep onset stay in place all night without anyone doing anything • Do not require caregiver intervention to maintain, unlike feeding or rocking • Running a sound machine at high volume close to the crib can risk hearing damage, so keep the volume low and the device well away from the crib
Breaking the Feed-to-Sleep Association	Feed 20 to 30 minutes before sleep rather than immediately before it, then use a calm activity like a short book before the crib	• Move feeding earlier in the bedtime routine so it is not the last step before sleep • Falling asleep at the breast or bottle each night can turn into something the baby needs recreated at every waking • Sleep problems that start in infancy often persist for years rather than resolving on their own without any change
Transferring a Sleeping Baby	Wait for signs of deeper sleep, such as looser limbs and slower breathing, before transferring to the crib	• The sudden change in support and movement during a transfer can trigger the Moro (startle) reflex common in young babies • Relying on being rocked or held to fall asleep and then transferred does not build the skill of falling asleep independently • Aim to replace transfers with drowsy but awake crib placement as that skill develops
Pacifier as Sleep Association	Pacifier falls out during the night; baby is already asleep and does not need it replaced	• Offering a pacifier at sleep onset is recommended to help reduce the risk of SIDS • If it falls out after the baby is already asleep, there is no need to put it back in • A pacifier repeatedly falling out overnight is not a safety problem and is not a reason to stop offering one at bedtime

TABLE 7

Sleep Environment Setup (Room, Darkness, White Noise, Temperature)

The sleep environment is the most controllable variable in infant sleep and provides the foundation all methods build on. AAP safe sleep guidance and pediatric sleep research point to specific conditions that support both safety and sleep quality. Getting the environment right often resolves a meaningful share of sleep issues before any method is applied.

Technique/Method	Example	Description
Darkness for Infant Sleep	Blackout curtains or blinds that block outside light; cover device LEDs; aim for a room dark enough that you can't see your hand in front of your face	• A clear switch between bright days and dark nights, not constant dim lighting, helps entrain a baby's internal clock to the 24 hour day and night cycle • Research on infants found a softly lit room around the clock lines up worse with that clock than a true light and dark cycle does • Keeping the room dark also helps prevent early morning waking, since sunrise light is a strong wake up cue
White Noise for Infant Sleep	A sound machine set to a low, steady volume and placed well away from the crib rather than right beside it	• Sound level drops off quickly with distance, so placement matters as much as the volume dial • Testing on infant sleep machines found that close to the crib, every device measured exceeded the noise limit used for hospital nurseries, and some produced levels loud enough to risk hearing damage with hours of nightly use • Louder or closer is not simply "better masking with no downside"
Room Temperature	A room that feels comfortably cool, roughly 68 to 72°F (20 to 22°C) for many households; dress baby in about one layer more than an adult would wear, using a sleep sack instead of loose blankets	• Overheating is a recognized SIDS risk factor, so a room on the cool side is safer than a warm one • Babies take time to develop the ability to self regulate temperature and sweat less than adults, so matching adult clothing exactly can leave them too warm • A wearable sleep sack replaces loose blankets and can be used safely from birth
AAP Safe Sleep Surface (2022)	A firm, flat mattress in a safety approved crib, bassinet, or play yard, with no bumpers, pillows, positioners, or loose blankets; baby placed on their back for every sleep	• Back sleeping for every sleep until age 1 lowers the risk of sudden infant death syndrome (SIDS) compared with stomach or side sleeping • Room sharing (baby's own sleep surface in the parents' room) for at least the first 6 months is linked to roughly half the SIDS risk and is far safer than bed sharing • Bed sharing is not recommended, and carries added risk when a caregiver is fatigued, has used alcohol or drugs, or the baby is very young or was born preterm
Sleep Sack / Swaddle Transition	Stopping swaddling at the first sign of rolling attempts, typically around 3 to 4 months but sometimes earlier, and switching to an arms out swaddle or a sleep sack	• A swaddled baby who manages to roll onto their stomach cannot free their arms to push up or turn their head, which raises the risk of suffocation • This is a hard stop at the first sign of rolling, not a "keep going while watching closely" judgment call • Wearable sleep sacks can be used from birth onward as a swaddle free alternative, with warmth chosen to suit room temperature
Noise Environment Optimization	A sound machine kept at a low, steady volume rather than run at maximum output for hours every night	• Some sound machines can exceed 91 decibels at maximum volume, above the noise exposure limits set for an 8 hour adult work shift • A 2024 review of the research found continuous, moderate to loud white noise exposure raises early development concerns in the study data, while lower intensity noise appeared safe or even helpful • Limiting both volume and how long a machine runs at higher output is the more cautious approach while research continues

TABLE 8

Naps vs. Nights - Training Order and Strategy

Many parents wonder whether to sleep train naps, nights, or both at the same time. Clinical sources agree that nighttime sleep is easier to train first, since the drive to sleep is highest at bedtime, and that naps predictably take longer to settle no matter when training starts.

Technique/Method	Example	Description
Nights First, Naps Second (Nap Training Protocol)	Begin sleep training at bedtime; once nighttime sleep is consistent, apply the same approach to naps	• Sleep drive (the pressure to sleep that builds the longer a baby is awake) is highest at bedtime, so nighttime training usually clicks faster than daytime naps • The same method that works at night can be applied to naps too, but naps predictably take longer to improve, so patience matters once training moves to naps

Technique/Method	Example	Description
Contact Napping to Crib Transition	Start with one crib nap per day (usually the morning nap, when sleep drive is highest); keep other naps as contact naps for now	- The morning nap carries the strongest sleep drive of the day, which makes it the easiest nap to move to independent sleep first - Moving one nap at a time, rather than overhauling every nap at once, is the recommended approach - Even a short or partial crib nap during the transition counts as real progress
Split Approach (Different Methods for Naps/Nights)	A gentler method like Pick Up Put Down for naps alongside a firmer method like Ferber at night	- There is no single required sleep training method; picking what fits the family and staying consistent within that choice is what matters - Consistency within each context (the same approach every night, the same approach every nap) is the emphasis, not using an identical method for naps and nights
Bedtime Timing for Optimal Training	A consistent, held bedtime, with the last nap ending well before bedtime	- Being overtired does not make a baby sleepier; it makes them more wired and harder to console, which is why a too-late bedtime is a common reason training stalls - Setting a bedtime and holding it every night is one of the central pieces every sleep training method depends on

Topic edits

- Description: rewritten (was: "Many parents wonder whether to sleep train naps, nights, or both simu...")
- Rows added: none
- Rows removed: none
- Rows reordered: no
- Column 1 link updates: Nights First, Naps Second (Nap Training Protocol): <https://www.babysleepsite.com/sleep-training/should-you-sleep-train-naps-and-night-sleep-at-the-same-time/> -> <https://health.clevelandclinic.org/ferber-method>, Contact Napping to Crib Transition: <https://www.preciouslittlesleep.com/contact-naps/> -> <https://cradlewise.com/blog/contact-naps-babies-sleep-safety/>, Split Approach (Different Methods for Naps/Nights): <https://www.happiestbaby.com/blogs/baby/how-to-sleep-train-2> -> <https://www.sleepfoundation.org/baby-sleep/sleep-training>, Bedtime Timing for Optimal Training: <https://www.sleepfoundation.org/baby-sleep/bedtime-for-babies> (404, dead) -> <https://healthier.stanfordchildrens.org/en/pediatrician-insights-on-childhood-sleep-problems/>
- Row description rewrites: Nights First, Naps Second (Nap Training Protocol), Contact Napping to Crib Transition, Split Approach (Different Methods for Naps/Nights), Bedtime Timing for Optimal Training
- Row example rewrites: Nights First, Naps Second (Nap Training Protocol) (removed unconfirmed "7-14 days" / "1-2 weeks" figures), Bedtime Timing for Optimal Training (removed unconfirmed "6:30-7:30 pm" and "2-3.5 hours" figures that could not be verified on any Tier 1-3 source)
- Duplicates merged: Nights First, Naps Second + Nap Training Protocol (the original "Nap Training Protocol" row's distinguishing claim, a hard 45-60 minute nap-attempt cutoff, could not be confirmed on any Tier 1-3 source despite extensive search across Cleveland Clinic, Sleep Foundation, Yale Medicine, Stanford Children's Health, AASM-adjacent material, and PubMed; it only appeared on unnamed content-aggregator sites below the sourcing bar for a graded fact. The row's verifiable substance, that naps take longer to train than nights even with the same method, duplicated the sequencing row, so the two are merged into one Column 1 entry)
- Deprecated demotions: none

Justification: n/a (edits made; see above)

TABLE 9

Sleep Regressions and How to Handle Them

Sleep regressions are temporary stretches of disrupted sleep tied to developmental leaps, neurological changes, or physical milestones. The 4-month regression stands apart from the rest, since it is a permanent shift in how sleep itself is organized rather than a phase that resolves on its own. Every regression after it is temporary, so keeping a consistent response through each one is what actually shortens the disruption.

Technique/Method	Example	Description
4-Month Sleep Regression	Was sleeping 5-6 hr stretches; suddenly waking every 45-90 min at night; naps falling apart	- Permanent shift in sleep organization as cycling matures toward an adult-like pattern - Does not simply pass; baby needs to learn to settle between cycles independently - Best time to begin sleep training if not already started

Technique/Method	Example	Description
8-Month Sleep Regression	Baby who was sleeping well begins waking 2-4x/night; separation anxiety at bedtime; may be crawling or pulling to stand	• Driven by object permanence taking hold, fresh separation anxiety, and new mobility like crawling or standing • Temporary, typically 2-6 weeks • Maintain the existing sleep training approach consistently through it
12-Month Sleep Regression	New walker waking at night; pushing back on naps; bedtime battles increasing	• Linked to the walking milestone, early word attempts, and growing independence • Nap transition to 1 nap is often confused with this regression; distinguish by watching wake windows • Typical duration 2-4 weeks
18-Month Sleep Regression	Toddler who slept well begins bedtime resistance, night calling, early waking; vocabulary growing fast	• A major developmental leap where language growth, autonomy, and still-immature emotional regulation combine • Separation anxiety is at its peak during this window; bedtime battles intensify • A firm, consistent routine plus a brief goodbye ritual works best
2-Year Sleep Regression	Stalling tactics at bedtime, nighttime waking calling for parent, early morning waking, refusing nap	• Driven by language growth, early imagination (nighttime fears beginning), and autonomy assertion • Switching from crib to bed during the regression tends to make things harder; delaying it if possible usually goes smoother • Consistent limit setting plus brief positive reinforcement (like a sticker chart) works best
Maintaining Sleep Training During Regression	Continue the same method during a regression; offer extra daytime comfort but hold the line at night	• Introducing a new soothing method mid regression creates a fresh sleep association that has to be unlearned later, stretching out the disruption • Extra daytime physical contact and attention can help buffer separation anxiety • Most regressions resolve within 2-6 weeks if the existing routine is kept steady

TABLE 10

Night Weaning

Night weaning, cutting back or dropping nighttime feeds, is a separate decision from sleep training, though many families tackle both around the same time. How early it makes sense depends heavily on how a baby is fed: guides generally treat formula fed babies as reasonable candidates from about 6 months, while breastfed babies are usually not considered ready until closer to 12 months, since breast milk digests faster and cutting back too early can affect milk supply. Whichever method is used, protecting that supply for a breastfeeding parent is the main reason gradual approaches are favored over abrupt ones.

Technique/Method	Example	Description
When Night Weaning Is Appropriate	Formula fed baby 6+ months; breastfed baby closer to 12 months; pediatrician confirms steady weight gain and no medical concerns	• Formula digests more slowly than breast milk, so formula fed babies are commonly ready to consider night weaning from around 6 months • Breastfed babies typically need night feeds longer since breast milk digests faster, guides generally suggest waiting until closer to 12 months • A steady weight gain and normal feeding pattern is worth confirming with a pediatrician before dropping feeds, regardless of feeding type
Dream Feed (Scheduled Night Feed)	Feed a still mostly asleep baby about two to three hours after bedtime, close to the parent's own bedtime, before the baby would otherwise wake to feed	• Proactively feeds the baby before an unpredictable night waking, instead of waiting for the baby to wake and cry for food • Most useful in the earliest months, effectiveness commonly fades within the first 12 weeks or so as overnight sleep naturally consolidates • Best dropped last, after other night feeds are already eliminated, rather than kept as a permanent nightly ritual
Gradual Night Weaning (Reduce-Duration Method)	Shorten a breastfeed by 2 to 5 minutes, or a bottle by 20 to 30 ml, every second night over about 5 to 7 nights until the feed is gone	• Slow enough to let milk supply adjust instead of dropping suddenly • A feed that is already short (under 5 minutes at the breast, or 60 ml or less from a bottle) can often just be stopped outright and resettled • A single feed is typically eliminated within about a week using this pace

Technique/Method	Example	Description
Cold Turkey Night Weaning (Older Babies)	9+ months, well established daytime solids and feeds; stop offering the night feed and resettle with whatever sleep approach the family is using	• Works faster for older babies whose daytime solids and feeds already meet their calorie needs • May take some adjustment and extra fussing at first, which usually settles within a handful of nights • If breastfeeding, still worth checking in with a lactation consultant, since stopping a regular feed abruptly can affect supply at any age, though less so than in a younger baby
Night Weaning Twins / Multiple Babies	Feed both twins around the same time, gently waking the second right after finishing the first, rather than letting feeds spread across the night	• Feeding together, or waking the second baby right after the first, cuts the total number of separate wake ups rather than adding to them • White noise or soft lighting can help both babies settle back down after a feed • Twins still follow the same room sharing safe sleep guidance as a single baby, so moving them to separate rooms to dodge a shared wake up is not a safety trade worth making

TABLE 11

Early Morning Waking Troubleshooting

Early morning waking (before 6 am) is one of the most common and frustrating sleep challenges after sleep training is established. It is caused by a combination of factors: circadian biology (cortisol peaks at dawn), light, insufficient sleep pressure, and overtiredness. Most fixes require environmental or schedule changes, not method changes.

Technique/Method	Example	Description
Light as Early Wake Cause	Baby wakes at 5:30-6 am as dawn light enters room; adding blackout curtains resolves problem	• First and most effective fix: complete darkness • Summer months especially problematic (sunrise 5-5:30 am) • Tape over device LEDs; block door gaps with draft stopper
Overtiredness Causing Early Rising	Baby going to bed at 8:30 pm waking at 5 am; shifting bedtime to 7 pm extends morning wake to 6:30 am	• Paradox: later bedtime = earlier rising (overtiredness elevates cortisol) • Earlier bedtime (6:30-7:30 pm) is often the counter-intuitive solution • Most impactful schedule adjustment available
Insufficient Night Sleep Pressure	Baby taking 3 naps + bedtime 7 pm; transition to 2 naps and cap last nap at 3:30 pm	• Ending the last nap earlier shifts more sleep toward the night • Cap total daytime sleep per age (see Table 3) and cut off last nap by 4 pm at latest • Often helps resolve early morning waking within about a week
Treating 5 am as Night (Not Morning)	Baby wakes at 5:15 am; parent treats as night waking (minimal interaction, dark room) and uses brief check-in	• Do not feed or start day until minimum desired wake time (6-6:30 am) • Reinforces that pre-6 am is nighttime, so the early waking isn't rewarded with attention • Consistency over 7-14 days typically shifts wake time forward
Hunger as Early Wake Cause	Night-weaned 8-month-old waking at 5 am; adding solid food dinner resolves it	• Ensure adequate calories during day especially in early stages of night weaning • More daytime and bedtime calories can reduce night feeding, but often doesn't stop the waking itself once a baby is past early infancy • Growth spurts increase caloric need and can cause temporary early waking

TABLE 12

Sleep Training Twins and Shared Rooms

Twins and siblings sharing a room present unique logistical challenges for sleep training. The main concerns are one baby waking the other, and whether to start training simultaneously. Research and clinical experience support training twins together in the same room as generally preferable to separating them for training.

Technique/Method	Example	Description
Training Twins Together vs. Separately	Keep twins in same room; begin training at same time with same method for both	• Separation for training is rarely necessary; twins typically habituate to each other's crying within the first few nights • Twins get used to each other's sounds faster than parents expect • Same schedule and method for both twins simplifies implementation
White Noise for Twins	Place white noise machine between cribs or one machine per crib; continuous operation	• Placing the machine between the two cribs buffers each twin's sound from the other • Volume sufficient to mask crying without exceeding AAP limits (under 50 dB at baby's ear); the limit does not rise just because two babies share the room • Cribs placed at opposite ends of room if space permits
Responding to Only One Twin Waking	Twin A cries; respond per chosen method; Twin B not yet awake, do not preemptively wake	• Do not wake the second twin unless a scheduled nighttime feed requires tandem feeds • Allow the waking twin to settle independently before assuming the second will wake too • Middle of the night wakings rarely involve both twins waking; this is less common than parents expect
Sibling in Shared Room	Move older sibling to a different room (or parent's room) for about the first week of sleep training a new baby	• Temporarily separating during the initial training phase reduces disruption to both children, since most crying happens in the first few nights • Older sibling returns to the shared room once the baby settles into the method • A sound machine helps the older sibling sleep through the baby's adjustment cries
Nap Training Twins	Train naps on the same schedule; natural synchronization builds gradually over the early months as both babies' internal sleep clocks mature	• A synchronized nap schedule is genuinely valuable for caregiver rest and a simpler daily routine • Twin sleep studies show naps and night sleep become more aligned gradually over the first several months, not within days of starting a shared schedule • Synchronized naps give caregivers real rest and simplify the daily routine

TABLE 13

When Sleep Training Is Not Working

When sleep training extends beyond 2-3 weeks without improvement, there is usually an identifiable cause. The most common culprits are: a hidden medical issue, inconsistent application, incorrect schedule/wake windows, or an environmental factor. Troubleshooting systematically before abandoning a method is recommended.

Technique/Method	Example	Description
Medical Causes of Sleep Failure	GERD (acid reflux), obstructive sleep apnea, ear infections, food protein intolerance causing discomfort	• Rule out medical causes before any method • Sleep training over unmanaged pain rarely works • Signs: back-arching during feeds, spitting up during/after sleep, snoring/mouth breathing • Pediatric evaluation is step zero if training not progressing
Inconsistency as Primary Cause	Method applied one night, abandoned second night when crying continues; baby learns crying long enough works	• Most common cause of training failure • Partial reinforcement (sometimes responding, sometimes not) is harder to extinguish than consistent responding • Consistency across all caregivers (both parents, grandparents, overnight sitters) is essential
Incorrect Wake Windows	Training attempted right after a much shorter wake time than usual; 60 minutes of crying with no signs of drowsiness at all	• Wake windows too short means insufficient sleep pressure, which means training attempts fail • Audit the schedule first when a method isn't working • Extend the wake window by 15-30 min and re-assess over a few days
Extinction Burst (Normal, Not Failure)	Nights 1-3: 45 min crying; Night 4: 90 min crying, parents assume it's not working	• Extinction burst: temporary spike in crying intensity/duration, typically around night 4-7 • Sign the method IS working, behavior escalates before it fades • Parents who push through the burst see dramatic improvement by night 8+

Technique/Method	Example	Description
Split Night Problem	Baby sleeps 7 pm to 12 am, then awake 12 to 3 am playfully (not crying), then sleeps 3 am to 6 am	• Split night caused by an underdeveloped body clock, circadian misalignment, or too much daytime sleep • Solution: earlier nap cutoff, reduce total daytime sleep, keep the middle of the night calm and quiet • Not a sleep training failure, a schedule adjustment issue
Developmental Leaps During Training	Baby learning to pull to stand; wakes and stands in crib; cannot get back down	• Teach the new skill (lowering from stand) during waking hours • Motor development temporarily disrupts sleep training progress • Continue the method; most disruptions resolve within a few weeks (commonly two to four) as the skill consolidates

TABLE 14

Special Scenarios and Advanced Topics

Beyond the core methods and schedules, caregivers encounter specific situations that require adapted strategies. This table covers scenarios including swaddle transitions, travel disruption recovery, sleep training after illness, toddler sleep in a bed (not crib), and caregiver mental health considerations.

Technique/Method	Example	Description
Swaddle-to-Sleep-Sack Transition	Baby is around 2-3 months old; the one-arm-out step should already be finished, with both arms free, before any rolling attempt shows up	• Stop swaddling completely as soon as baby shows any sign of trying to roll, per AAP guidance • The one-arm-out step works best done proactively, before rolling begins, not as a week-long response after it starts • No evidence that one arm in versus out changes safety, so a partial swaddle is not a safe holding pattern once rolling begins
Post-Illness Sleep Reset	Baby sick for a week; caregiver comforted through every wake-up; now healthy again but waking often	• Normal: increased waking during and right after illness is a common, temporary regression, not training failure • Reset: return to the same consistent routine used before, without adding new sleep supports; usually settles within a couple weeks • New accommodations added during the sick stretch (co-sleeping, extra rocking), not the illness itself, are what tend to prolong the disruption
Travel / Vacation Sleep Disruption	Pack a portable crib or playard, a familiar crib sheet, and keep the usual bedtime routine at the hotel	• Keep the same safe sleep setup and routine cues, not just a good location; AAP guidance says not to relax safe sleep habits while traveling • Use a portable crib or playard rather than bed-sharing, even for just a few nights • Bringing something familiar, like the usual crib sheet, helps recreate home cues in a new place
Crib-to-Bed Transition	Most toddlers switch between 18 months and 3 years old; if crib escape is the reason, try a crib tent first	• No official minimum age; timing follows safety cues like climbing attempts or the crib rail reaching chest height • Switching before a child shows readiness signs (sleeping through the night, following house rules) tends to increase night wandering and disruption • Bed rails, a visual okay-to-get-up cue (like a color-changing night light), and a consistent return-to-bed routine all help
Caregiver Mental Health and Sleep Training	Caregivers take turns overnight so no one carries every high-cry night alone	• Parental burnout and postpartum mood disorders are real risk factors that affect training consistency • Alternating overnight duties between caregivers protects each person's own sleep, which is itself linked to mood • Postpartum Support International: 1-800-944-4773
Positive Reinforcement for Toddlers	Sticker chart: every morning the child stayed in bed all night, they get a sticker; a set number of stickers earns a small reward	• Works once a toddler can cognitively connect tonight's behavior with tomorrow's reward, not at one fixed universal age • Reward must be immediate and consistent to land • Combine with a visual cue (like a clock or light that shows when it's okay to get up) for best results

Technique/Method	Example	Description
High-Needs Babies and Sleep Training	Baby with an intense, easily overwhelmed temperament; full extinction feels like too much, so a gentler approach like the Sleep Lady Shuffle or Pick Up Put Down fits better	• A gentler, more parent-present approach still leads to independent sleep, just typically on a slower timeline • No single method fits every child; temperament shapes the right approach • Real-world data show gentler approaches produce no difference in parent-child bonding compared with faster methods

References

Infant & toddler sleep (behavioral pediatrics)

- 12-Month Sleep Regression - <https://www.healthychildren.org/English/ages-stages/baby/Pages/Developmental-Milestones-12-Months.aspx>
- 4-Month Sleep Regression - <https://health.clevelandclinic.org/the-4-month-sleep-regression-what-parents-need-to-know>
- AAP Safe Sleep Surface (2022) - <https://publications.aap.org/pediatrics/article/150/1/e2022057990/188304/Sleep-Related-Infant-Deaths-Updated-2022>
- Early Tired Signs (Optimal Sleep Window) - <https://health.clevelandclinic.org/wake-windows-by-age>
- Hunger as Early Wake Cause - <https://pubmed.ncbi.nlm.nih.gov/25973527/>
- Maintaining Sleep Training During Regression - <https://www.healthychildren.org/English/ages-stages/baby/sleep/Pages/Getting-Your-Baby-to-Sleep.aspx>
- Minimum Age for Sleep Training - <https://health.clevelandclinic.org/when-and-how-to-sleep-train-your-baby>
- Nights First, Naps Second (Nap Training Protocol) - <https://health.clevelandclinic.org/ferber-method>
- Pacifier as Sleep Association - <https://safetosleep.nichd.nih.gov/reduce-risk/reduce>
- Pick Up Put Down / PUPD (Tracy Hogg) - <https://health.clevelandclinic.org/pick-up-put-down-method>
- Sleep Associations (Negative) - https://aasm.org/resources/practiceparameters/review_nightwakingschildren.pdf
- Swaddle-to-Sleep-Sack Transition - <https://www.healthychildren.org/English/ages-stages/baby/diapers-clothing/Pages/Swaddling-Is-it-Safe.aspx>
- Toddler Sleep Needs (2-3 years) - <https://www.sleepfoundation.org/children-and-sleep/how-much-sleep-do-kids-need>
- White Noise for Infant Sleep - <https://publications.aap.org/pediatrics/article/133/4/677/32749/Infant-Sleep-Machines-and-Hazardous-Sound-Pressure>
- 2-Year Sleep Regression - <https://health.clevelandclinic.org/from-crib-to-toddler-bed>
- 8-Month Sleep Regression - <https://www.healthychildren.org/English/ages-stages/toddler/Pages/Soothing-Your-Childs-Separation-Anxiety.aspx>
- Chair Method / Gradual Withdrawal - <https://health.clevelandclinic.org/chair-method-sleep-training>
- High-Needs Babies and Sleep Training - <https://www.healthychildren.org/English/ages-stages/baby/Pages/Babys-Temperament.aspx>
- Infant Sleep Needs (9-12 months) - <https://www.healthychildren.org/English/healthy-living/sleep/Pages/healthy-sleep-habits-how-many-hours-do-es-your-child-need.aspx>
- Medical Causes of Sleep Failure - <https://www.healthychildren.org/English/health-issues/conditions/abdominal/Pages/GERD-Reflux.aspx>
- Newborn Sleep Needs (0-3 months) - <https://www.sleepfoundation.org/baby-sleep/how-much-sleep-do-babies-need>
- Noise Environment Optimization - <https://pubmed.ncbi.nlm.nih.gov/38663282/>
- Pediatrician Sign-Off - <https://www.healthychildren.org/English/ages-stages/baby/sleep/Pages/Sleeping-Through-the-Night.aspx>
- Room Temperature - <https://www.healthychildren.org/English/ages-stages/baby/sleep/Pages/A-Parents-Guide-to-Safe-Sleep.aspx>
- Split Approach (Different Methods for Naps/Nights) - <https://www.sleepfoundation.org/baby-sleep/sleep-training>
- Toddler Sleep Needs (12-24 months) - <https://www.aasm.org/resources/clinicalguidelines/pediatric-sleep-duration.pdf>
- Travel / Vacation Sleep Disruption - <https://www.healthychildren.org/English/safety-prevention/on-the-go/Pages/Hotel-Dangers-that-Put-Baby-Safety-at-Risk.aspx>

Leadership & health authorities

- Nap Training Twins - <https://pmc.ncbi.nlm.nih.gov/articles/PMC6347043/>
- Overtiredness Causing Early Rising - <https://pmc.ncbi.nlm.nih.gov/articles/PMC8813037/>

Behavioral science & psychology

- Dream Feed (Scheduled Night Feed) - https://www.apa.org/pubs/books/supplemental/pediatric-sleep-problems/Dream_Feed.pdf

Academic papers & preprints

- Insufficient Night Sleep Pressure - <https://www.nature.com/articles/srep27246>

Articles, guides, & named-author resources

- 18-Month Sleep Regression - <https://www.msmanuals.com/home/children-s-health-issues/symptoms-in-infants-and-children/separation-anxiety-and-stranger-anxiety>
- Babywise (Ezzo & Bucknam) - <https://www.babywise.life>
- Age-Specific Tired Sign Differences - <https://www.firstfiveyears.org.au/child-development/parenting-techniques/tired-signs-in-babies-and-toddler-s>
- Bedtime Timing for Optimal Training - <https://healthier.stanfordchildrens.org/en/pediatrician-insights-on-childhood-sleep-problems/>

36. Breaking the Feed-to-Sleep Association - <https://www.stanfordchildrens.org/en/topic/default?id=infant-sleep-90-P02237>
38. Contact Napping to Crib Transition - <https://cradlewise.com/blog/contact-naps-babies-sleep-safety/>
40. Darkness for Infant Sleep - <https://link.springer.com/article/10.1007/s00431-024-05951-3>
42. Developmental Signs of Readiness - <https://www.merckmanuals.com/professional/pediatrics/care-of-newborns-and-infants/sleeping-behaviors-in-infants-and-children>
44. Extinction Burst (Normal, Not Failure) - https://www.psywww.com/intropsych/ch05_conditioning/extinction_burst.html
46. Inconsistency as Primary Cause - https://www.psywww.com/intropsych/ch05_conditioning/intermittent_reinforcement.html
48. No-Cry Sleep Solution (Elizabeth Pantley) - <https://www.pantley.com/>
50. Parental Readiness - <https://childmind.org/article/choosing-a-sleep-training-method-that-works-for-your-family/>
52. Responding to Only One Twin Waking - <https://drcraigcanapari.com/agonny-ecstasy-room-sharing-siblings/>
54. Sleep Lady Shuffle (Kim West) - <https://sleeplady.com/baby-sleep/sleep-lady-shuffle/>
56. Split Night Problem - <https://my.clevelandclinic.org/health/articles/14300-sleep-in-your-babys-first-year>
58. Transferring a Sleeping Baby - <https://my.clevelandclinic.org/health/articles/moro-reflex>
60. Wake Windows 14-24 Months - <https://huckleberrycare.com/blog/2-to-1-nap-transition>
62. Wake Windows 5-7 Months - <https://huckleberrycare.com/blog/3-to-2-nap-transition>
64. Weight Threshold Guideline - <https://www.uchicagomedicine.org/forefront/pediatrics-articles/sleep-training-in-infants-and-toddlers>
66. When Night Weaning Is Appropriate - <https://raisingchildren.net.au/babies/sleep/settling-routines/night-weaning>
37. Caregiver Mental Health and Sleep Training - <https://www.postpartum.net/get-help/>
39. Cortisol Second Wind - <https://www.earlylearningsociety.org/blog/Why-Wont-My-Toddler-Sleep-Understanding-and-Navigating-Bedtime-Resistance-in-Two-Year-Olds.htm>
41. Developmental Leaps During Training - <https://www.childrens.com/health-wellness/sleep-regression-in-infants-and-toddlers>
43. Drowsy-But-Awake Definition - <https://parenting.mountsinai.org/tip-of-the-week/drowsy-but-awake-2024>
45. Ferber Method (Graduated Extinction) - <https://www.ferbermethod.com>
47. Night Weaning Twins / Multiple Babies - <https://twinstrust.org/advice-and-support/first-year/sleep-tips-for-parents-of-twins/>
49. Overtired Signs - <https://pathways.org/sleep-cues-to-tell-if-baby-is-sleepy>
51. Post-Illness Sleep Reset - <https://www.phoenixchildrens.org/articles-faqs/blog/managing-sleep-regression-infants-and-toddlers>
53. Sibling in Shared Room - <https://www.babywisemom.com/managing-cry-it-out-while-room-sharing/>
55. Sleepeasy Solution (Waldburger & Spivack) - <https://www.sleepyplanet.com/>
57. Training Twins Together vs. Separately - <https://baby.chparenting.com/blog/sleep-training-twins/>
59. Undertired Signs - <https://littlesleepcoach.com/is-baby-overtired-or-under-tired/>
61. Wake Windows 4-10 Weeks - <https://huckleberrycare.com/blog/first-year-of-sleep-expectations>
63. Wake Windows 7-14 Months - <https://huckleberrycare.com/blog/nap-transitions-when-they-occur-and-how-to-handle-them>
65. Weissbluth Method (Full Extinction / Cry It Out) - <https://marcweissbluth.com/25-let-cry-sleep-solutions-graduated-extinction-extinction-sleep-solutions-3/>