

ReefSafe decision dashboard



Observed reef evidence + cautious next-observation screening | Malaysia, 2012–2025

Selected island Labuan Change the yellow cell to inspect any monitored island

Current LCC	Observed change	Predicted next change	Priority
24.2%	-7.3 pp/yr	-14.4 pp/yr	High screening priority
Uncertainty interval -30.7 to -0.8 pp/yr		Intervals are empirical screening bands, not causal or site-specific certainty.	

Observed evidence
NOAA thermal stress observed

Recommended next step
Coordinate bleaching survey; do not attribute thermal loss to visitors



Island	State	Current LCC	Predicted change	Evidence
Labuan	W.P. Labuan	24.2	-14.4	NOAA thermal stress observed
Kapas	Terengganu	52.6	-5.8	Modelled decline; no single dominant observed stressor
Seri Buat	Pahang	54.2	-5.2	Anchor impact mentioned in Reef Check report
Mertang	Johor	58.3	-5.0	Anchor impact mentioned in Reef Check report
Mensirip	Johor	54.2	-4.8	Waste or pollution indicator is elevated
Tenggol	Terengganu	54.2	-4.6	Waste or pollution indicator is elevated
Tinggi	Johor	52.5	-4.6	Anchor impact mentioned in Reef Check report
Lima	Johor	51.9	-4.4	Anchor impact mentioned in Reef Check report
Sibu	Johor	52.8	-4.3	Anchor impact mentioned in Reef Check report
Lankayan	Sabah	51.0	-4.1	NOAA thermal stress observed

Model	MAE	RMSE	R²
Baseline mean	5.991	8.100	-0.019
Ridge regression	5.990	7.939	0.021
Gradient boosting	5.859	7.739	0.070
Random forest	6.074	8.017	0.002

Model check Gradient boosting MAE 5.86 pp/yr vs baseline 5.99 pp/yr (+2.2%). R² 0.070. NOAA heat is context only. Use the model to rank field checks, not as proof of tourism causality.