

随机模拟方法与应用导论作业五

陈稼霖 45875852

2019-10-03

5.5 (Exploring percentages of full-time faculty)

The variable `Full.time` in the college dataset (see Example 5.3) contains the percentage of faculty who are hired full-time in the group of National Universities.


- Using the `hist` function, construct a histogram of the full-time percentages and comment on the shape of the distribution.
- Use the `froot` and `flog` transformations to reexpress the full-time percentages. Construct histograms of the collection of `froots` and the collection of `flogs`. Is either transformation successful in making the full-time percentages approximately symmetric?
- For data that is approximately normally distributed, about 68% of the data fall within one standard deviation of the mean. Assuming you have found a transformation in part (b) that makes the full-time percentages approximately normal, find an interval that contains roughly 68% of the data on the new scale.

- 首先读取文件`college.txt`

```
dat = read.table('college.txt', head = TRUE, sep = '\t')
```

然后依据变量`Full.time`绘制全职教职工百分比的直方图


```
hist(dat$Full.time, xlab = 'Full-time Faculty Percentage',  
     main = 'Full-time Faculty Percentage')
```



评价：由图可见，绝大多数大学的全职教职工百分比在80%以上，直方图形状右偏(right-skewed)。

b. 用froot转换重新表示全职教职工百分比并绘制其直方图


```
froot = sqrt(dat$Full.time) - sqrt(100 - dat$Full.time)
hist(froot,xlab = 'Full-time Faculty Percentage(froot)',
     main = 'Full-time Faculty Percentage (froot) histogram')
```



用flog转换重新表示全职教职工百分比并绘制其分布直方图

```
flog = log(dat$Full.time) - log(100 - dat$Full.time)
hist(flog,xlab = 'Full-time Faculty Percentage (flog)',
     main = 'Full-time Faculty Percentage (flog) histogram')
```

Full-time Faculty Percentage (flog) histogram



两种转换都使得全职教职工百分比的直方图更加对称。

- c. 对于froot转换后得到的直方图，约68%的数据集中的区间为(4, 8]；对于flog转换后得到的直方图，约68%的数据集中的区间为(1, 3]。