

Name: 陈稼霖  
StudentID: 45875852

**Problem 1.** Let  $X$  be the total from rolling 6 fair dice, and let  $X_1, \dots, X_6$  be the individual rolls. What is  $P(X = 18)$ ?

*Solution:* The generating function of  $X_i$  ( $i = 1, 2, \dots, 6$ ) is

$$g_{X_i}(z) = \frac{1}{6}z + \frac{1}{6}z^2 + \frac{1}{6}z^3 + \frac{1}{6}z^4 + \frac{1}{6}z^5 + \frac{1}{6}z^6 = \frac{z}{6}(1 + z + z^2 + z^3 + z^4 + z^5 + z^6) \quad (i = 1, 2, \dots, 6)$$

Then the generating function of  $X = \sum_{i=1}^6 X_i$  is

$$g_X(z) = \prod_{i=1}^6 g_{X_i}(z) = \frac{z^6}{6^6}(1 + z + z^2 + z^3 + z^4 + z^5)^6 = \frac{z^6}{6^6}\left(\frac{1 - z^6}{1 - z}\right)^6$$

where

$$(1 - t^6)^6 = \sum_{j=0}^6 \binom{6}{j} (-1)^j z^{6j}$$

and

$$\frac{1}{(1 - z)^6} = (1 + z + z^2 + \dots)^6 = \sum_{k=0}^{\infty} a_k z^k$$

Here,  $a_k$  is the number of lists (“list” means order matters) consisting of six independent non-negative integer which add to  $k$ . We can also regard  $a_k$  as the number of cases in which  $k$  balls in a line are separated by five boards, so

$$a_k = \binom{k + 5}{5}$$

So

$$\frac{1}{(1 - z)^6} = (1 + z + z^2 + \dots)^6 = \sum_{k=0}^{\infty} \binom{k + 5}{5} z^k$$

and

$$g_X(z) = \frac{z^6}{6^6} \left[ \sum_{j=0}^6 \binom{6}{j} (-1)^j z^{6j} \right] \left[ \sum_{k=0}^{\infty} \binom{k + 5}{5} z^k \right]$$

In the generating function above, the coefficient of  $z^{18}$  item is

$$\frac{1}{6^6} \left[ \binom{6}{0} \binom{17}{5} - \binom{6}{1} \binom{11}{5} + \binom{6}{2} \binom{5}{5} \right] = \frac{3431}{6^6} \approx 0.07354$$

Reference: Blitzstein, Joseph K., and Jessica Hwang. *Introduction to probability*. Chapman and Hall/CRC, 2014: p264-265. □

**Problem 2.** Find the MGF of  $X \sim Unif(a, b)$  and  $Y \sim Expo(\lambda)$ .

Name: 陈稼霖  
StudentID: 45875852

*Solution:* The PDF of  $X$  is

$$f_X(x) = \frac{1}{b-a}, \quad x \in (a, b)$$

The MGF of  $X$  is

$$M_X(t) = \int_a^b e^{-tx} f_X(x) dx = \frac{e^{-at} - e^{-bt}}{(b-a)t}$$

The PDF of  $Y$  is

$$f_Y(y) = \lambda e^{-\lambda y}, \quad y \geq 0$$

The MGF of  $Y$  is

$$M_Y(t) = \int_0^\infty e^{-tx} f_Y(y) dy = \frac{\lambda}{t + \lambda}$$

□

**Problem 3.** Find the MGF of  $X \sim \text{Bern}(p)$  and  $Y \sim \text{Bin}(n, p)$ .

*Solution:* The PMF of  $X$  is

$$P_X(x) = \begin{cases} p, & \text{if } x = 1 \\ 1 - p, & \text{if } x = 0 \end{cases}$$

The MGF of  $X$  is

$$M_X(t) = (1 - p) + pe^{-t}$$

The MGF of  $Y$  is

$$M_Y(t) = [M_X(t)]^n = [(1 - p) + pe^{-t}]^n$$

□

**Problem 4.** Consider a setting where a Poisson approximation should work well: let  $A_1, \dots, A_n$  be independent, rare events, with  $n$  large and  $p_j = P(A_j)$  small for all  $j$ . Let  $X = I(A_1) + \dots + I(A_n)$  count how many of the rare events occur, and Let  $\lambda = E(X)$ .

(a). Find the MGF of  $X$ .

(b). If the approximation  $1 + x \approx e^x$  (this is a good approximation when  $x$  is very close to 0 but terrible when  $x$  is not close to 0) is used to write each factor in the MGF of  $X$  as  $e$  to a power. What happens to the MGF? (Hint: if  $Y \sim \text{Pois}(\lambda)$ , then  $M_Y = e^{(e^t - 1)\lambda}$ )

Name: 陈稼霖  
StudentID: 45875852

*Solution:*

(a) The MGF of  $X$  is

$$\begin{aligned} M_X(t) &= \sum_{k=0}^{\infty} \frac{\lambda^k}{k!} e^{-\lambda} e^{kt} \\ &= e^{-\lambda} \sum_{k=0}^{\infty} \frac{(\lambda e^t)^k}{k!} \\ &= e^{-\lambda} e^{\lambda e^t} = e^{\lambda(e^t - 1)} \end{aligned}$$

(b) If the approximation  $1 + x \approx e^x$  works, the MGF of  $x$  is

$$M_X(t) = e^{\lambda t}$$

□

**Problem 5.** Suppose  $X \sim \text{Bin}(n, p)$ . By using Binomial PGF, find the expectation  $E(X)$  and variance  $\text{Var}(X)$ .

*Solution:* According to the definition, the PGF of a Bernoulli distributed random variable is

$$G_{X_i}(z) = (1 - p) + pz$$

The PGF of  $X$  is

$$G_X(z) = G_{\sum_{i=1}^n X_i}(z) = \prod_{i=1}^n G_{X_i}(z) = [(1 - p) + pz]^n$$

Then

$$\begin{aligned} G'_X(z) &= np[(1 - p) + pz]^{n-1} \\ G''_X(z) &= n(n - 1)p^2[(1 - p) + pz]^{n-2} \end{aligned}$$

The expectation is

$$E(X) = G'(1) = np$$

The variance is

$$\text{Var}(X) = G''(1) + G'(1) - [G'(1)]^2 = n(n - 1)p^2 + np - (np)^2 = np(1 - p)$$

□

Name: 陈稼霖  
StudentID: 45875852

**Problem 6.** If a random variable  $X$  has the following moment-generating function:

$$M(t) = \frac{1}{10}e^t + \frac{2}{10}e^{2t} + \frac{3}{10}e^{3t} + \frac{4}{10}e^{4t}$$

for all  $t$ , then what is the *PMF* of  $X$ ?

*Solution:* Let  $t = -\ln z$ , the PGF of  $X$  is

$$G_X(z) = M_X(-\ln z) = \frac{1}{10}z^{-1} + \frac{2}{10}z^{-2} + \frac{3}{10}z^{-3} + \frac{4}{10}z^{-4}$$

The PMF of  $X$  is

$$P_X(x) = \begin{cases} \frac{1}{10}, & x = -1 \\ \frac{2}{10}, & x = -2 \\ \frac{3}{10}, & x = -3 \\ \frac{4}{10}, & x = -4 \end{cases}$$

□

**Problem 7.** Suppose that  $Y$  has the following moment-generating function:

$$M_Y(t) = \frac{e^t}{4 - 3e^t}, t < -\ln(0.75)$$

I).Find  $E(Y)$

II).Find  $\text{Var}(Y)$

*Solution:*

I)

$$E(Y) = (-1)^1 E'(0) = \frac{4e^t}{(4 - 3e^t)^2} \Big|_{t=0} = 4$$

II)

$$E(Y^2) = (-1)^2 E''(0) = \frac{4e^t(16 - 9e^{2t})}{(4 - 3e^t)^4} \Big|_{t=0} = 28$$

$$\text{Var}(Y) = E(Y^2) - [E(Y)]^2 = 12$$

□

**Problem 8.** If a random variable  $X$  has  $E[X^k] = 0.2$   $k=1,2,3,\dots$ , then what is the *PMF* of  $X$ ?

Name: 陈稼霖  
StudentID: 45875852

*Solution:* The MGF of  $X$

$$\begin{aligned}M_X(t) &= 1 + \frac{(-1)^1}{1!} E(X^1)t + \frac{(-1)^2}{2!} E(X^2)t^2 + \frac{(-1)^3}{3!} E(X^3)t^3 + \dots \\&= 0.8 + 0.2\left(1 + \frac{(-1)^1}{1!}t + \frac{(-1)^2}{2!}t^2 + \frac{(-1)^3}{3!}t^3 + \dots\right) \\&= 0.8 + 0.2e^{-t}\end{aligned}$$

Let  $t = -\ln z$ , the PGF of  $X$  is

$$G_X(z) = M_X(-\ln z) = 0.8 + 0.2z$$

The PMF of  $X$  is

$$P_X(x) = \begin{cases} 0.2, & \text{if } x = 1 \\ 0.8, & \text{if } x = 0 \end{cases}$$

□