

A summary of some mainstream spatial statistical methods for large datasets

Yewen Chen

October 13, 2020

CONTENTS

1	Gaussian random field and its challenge	3
2	Several approaches to overcome the computational problem	4
3	Likelihood approximations in the spectral domain	5
3.1	Whittle's likelihood	5
4	Likelihood approximations in the spatial domain	5
4.1	Low rank methods	5
4.1.1	Fixed rank Kriging	6
4.1.2	Gaussian predictive processes (GPP)	6
4.1.3	Multi-resolution approximations	6
4.2	Sparse covariance methods	7
4.2.1	Tapering	7
4.2.2	Spatial partitioning	7
4.3	Sparse precision methods	8
4.3.1	LatticeKrig	8
4.3.2	Gaussian Markov random field approximations by SPDE	8
4.3.3	The nearest neighbor Gaussian process	9
5	Two exploratory limitations of the NNGP approach	10
5.1	Exploratory analysis	10
5.2	Improvement project	11

LIST OF FIGURES

Figure 1	The K-L distance, the Bhattacharyya distance and the Frobenius norm from the estimated covariance matrix to the true covariance matrix for the Matérn family with smoothness parameters $\nu = \frac{1}{2}$ and different spatial range parameters ϕ . And this true covariance matrix at 500 random locations from $[0, 100] \times [0, 100]$ was estimated by the NNGP with $m = 20$	10
----------	---	----

LIST OF TABLES

Table 1	Scope of application	5
---------	--------------------------------	---

1 GAUSSIAN RANDOM FIELD AND ITS CHALLENGE

Gaussian spatial processes has been popular for decades in spatial data contexts like geostatistics where they are known as kriging, and in computer experiments where they are deployed as surrogate models or emulators. More recently, they have become a popular prediction engine in the machine learning literature. The reasons are many, but the most important are probably that: the Gaussian structure affords a large degree of analytic capability not enjoyed by other general-purpose approaches to nonparametric nonlinear modeling; and because they perform well in out-of-sample tests.

Assume there is a response or dependent variable $Y(s)$ at a generic location $s \in \mathbf{D} \subset \mathbb{R}^2$ along with a $p \times 1$ vector of spatially referenced predictors $\mathbf{X}(s)$. A spatial regression model has the form

$$y(s) = \beta \mathbf{X}(s) + w(s) + \epsilon(s) \quad (1)$$

where β is the vector of regression coefficients. The residual from the regression is decomposed into two independent parts: a spatial process, $w(s)$, modelling spatial association, and an independent process, $\epsilon(s)$, also known as the nugget effect, modelling measurement error. The spatial process $w(s)$ in (1) is often referred to as spatial random effects, capturing the effect of unmeasured or unobserved covariates with spatial pattern.

The nugget effect $\epsilon(s)$ is often assumed to follow a normal distribution with variance τ^2 for every location s . The most common specification for $w(s)$ is $w(s) \sim \text{GP}(0, \mathbf{C}(\cdot, \cdot))$, a zero-mean Gaussian process with a valid covariance function $\mathbf{C}(s, s')$. It is often reasonable to assume a constant process variance and thus we specify $\mathbf{C}(s, s') = \sigma^2 \rho(s, s'; \theta)$, where $\rho(s, s'; \theta)$ is a correlation function and θ is a vector of correlation parameters which needs to be estimated from a finite number of observations, $\mathbf{Y} = (y(s_1), \dots, y(s_n))'$.

According to the above assumptions, $y(s)$ follows a spatial Gaussian process, and thus we have the log-likelihood function for $(\beta, \tau^2, \sigma^2, \theta)$:

$$l(\beta, \tau^2, \sigma^2, \theta) \propto -\frac{1}{2} \log |\Sigma| - \frac{1}{2} (\mathbf{Y} - \beta \mathbf{X})' \Sigma^{-1} (\mathbf{Y} - \beta \mathbf{X}) \quad (2)$$

where $\Sigma = \mathbf{C} + \tau^2 \mathbf{I}$. Then best linear unbiased prediction (BLUP) at an unobserved location s_0 can be obtained by the **Kriging equation**:

$$\hat{y}(s_0) = \beta \mathbf{X}(s_0) + \mathbf{c}'_{s_0} \Sigma^{-1} (\mathbf{Y} - \beta \mathbf{X}) \quad (3)$$

where $\mathbf{c}_{s_0} = (\mathbf{C}(s_0, s_1), \dots, \mathbf{C}(s_0, s_n))'$.

However, with large or massive data, direct implementation of these statistical process, including parameter estimation by (2) and interpolation by (3), becomes computationally prohibitive, since evaluating the log-likelihood in (2) or solving the Kriging equation (3) involve the Cholesky factorization of an $n \times n$ covariance matrix for data of size n , which requires $O(n^3)$ operations and $O(n^2)$ memory in general (see, e.g., Porcu et al. [2012], Sang and Huang [2012], Heaton et al. [2019], Ma and Kang [2020]).

2 SEVERAL APPROACHES TO OVERCOME THE COMPUTATIONAL PROBLEM

The approximation of the likelihood in either the spatial or spectral domain and the lower-dimensional representation are fundamental solutions to overcome computational obstacles. These solutions focus on model development, the design of efficient and parallel algorithms, and the improvement and efficient use of modern computing platforms (e.g., Using [TensorFlow](#) and [GPU](#)).

Developed models, official websites of scholars:

- 1) Conditional distributions: [Michael L. Stein](#) (The restricted likelihood, [2004](#));
- 2) Sparse covariance by tapering method: [Reinhard Furrer](#)([2006](#)); [Douglas Nychka](#) (bias correction, [2008](#)); [Michael L. Stein](#) (Statistical Properties, [2013](#));
- 3) Low-Rank methods: [Noel Cressie](#) (FRK, [2008](#)), [Sudipto Banerjee](#) (Predictive process, [2008](#)); [Matthias Katzfuss](#) (multiresolution approximation where the basis functions are automatically chosen, [2017](#));
- 4) Sparse precision: [Douglas Nychka](#) (lattice kriging, [2015](#)) by basis-function; [Finn Lindgren](#) (GRMF Approximations, [2011](#)) by SPDE; [Abhi Datta](#) (NNGP, [2016a](#), [2016b](#), [2016c](#), [2019](#)) by conditional distributions;
- 5) Spectral method: [Montserrat Fuentes](#) and [Joe Guinness](#) (Circulant embedding, [2017](#) and Periodic Embeddings, [2007](#), [2019](#)).
- 6) Discrete process convolutions. [Dave Higdon](#) ([2002](#)); [Bruno Sansó](#) ([2009](#)); [Franky Fouedjio](#) ([2016](#)).
- 7) Local approximate Gaussian processes: [Robert B. Gramacy](#), [[Gramacy and Apley](#), [2015](#)];
- 8) Vecchia's approximation: [Zilber and Katzfuss](#) [[2020](#)], [Katzfuss et al.](#) [[2020](#)] and [Jurek and Katzfuss](#) [[2020](#)].

Developed algorithms and platforms, official websites of scholars:

- 1) INLA: [Haavard Rue](#).
- 2) Parallel algorithm: [Matthias Katzfuss](#).
- 3) TensorFlow: [Andrew Zammit-Mangion](#).

Table 1: Scope of application

Model	non-stationary	space-time	multivariate data
FRK	✓	✓	✓
Predictive process	✓		✓
MRA	✓	✓	
Tapering		✓	✓
Spatial partitioning			
LatticeKrig	✓	×	×
SPDE	✓	✓	✓
NNGP	✓	✓	✓
Whittle's approximation	×	×	×

The websites of other scholars: 1) [Christopher K. Wikle](#), 2) [Bradley P. Carlin](#), 4) [David Bolin](#), 5) [Jonathan R. Stroud](#), 6) [YongTao Guan](#), 7) [Alan E. Gelfand](#), 8) [Peter J. Diggle](#), 9) [Huiyan Sang](#), 10) [Andrw Finley](#), 11) [Matthew J. Heaton](#), 12) [Ying Sun](#), 13) [Furong Sun](#), 14) [Jonathan R. Bradley](#).

3 LIKELIHOOD APPROXIMATIONS IN THE SPECTRAL DOMAIN

3.1 Whittle's likelihood

The spectral methods are computationally efficient by avoiding the calculation of determinants and can be easily adapted to model nonstationary processes as a mixture of independent stationary processes.

[Fuentes \[2007\]](#), [\[Guinness, 2019\]](#) and [\[Guinness and Fuentes, 2017\]](#) propose the use of small domain expansions and imputing data in a periodic fashion on the expanded lattice and presented a version of Whittle's approximation to the Gaussian negative log-likelihood by introducing a lattice process which can be used to deal with irregularly spaced data. Additional computational savings were obtained by truncating the spectral representation of the lattice process. The calculation requires $O(m \log_2 m + n)$ operations where m is lattice size.

Comment: The flexibility of the spectral methods mentioned above are disputable for prediction problem respecting out of sample data, since the expanded lattice is pre-defined.

4 LIKELIHOOD APPROXIMATIONS IN THE SPATIAL DOMAIN

4.1 Low rank methods

Comment: The reduced rank based methods usually fail to accurately capture the local, small scale dependence structure (see [\[Sang and Huang, 2012\]](#), [\[Stein, 2008\]](#) and [\[Finley et al., 2009\]](#)).

4.1.1 Fixed rank Kriging

FRK ([Cressie and Johannesson, 2006], [Cressie and Johannesson, 2008]) aims to approximate the spatial process $w(s)$ in (1) by a linear combination of K ($\ll n$) **multi-resolution bisquare** basis functions:

$$w(s) = \sum_{r=1}^R \sum_{k=1}^{K_r} h_{rk}(s) w_{rk}^* \quad (4)$$

where $K = \sum_{r=1}^R K_r$ and the coefficients w^* is an K -dimensional Gaussian vector with mean zero and exponential covariance defined by some small basic areal units, which ensures that all estimation and prediction equations only contain inverses of matrices of size $K \times K$.

Comment: Since covariance matrix for every resolutions is dense, that implies FRK can be challenging for large K_r .

4.1.2 Gaussian predictive processes (GPP)

With regard to the challenge of computational cost on covariance matrices, Banerjee et al. [2008] proposed a class of models based on the idea of a spatial predictive process which is motivated by Kriging equation in (3). The predictive process projects the original process, $w(s)$ in (1), onto a subspace generated by realizations of the original process at a specified set of locations (or knots), e.g., s_1^*, \dots, s_K^* , $K \ll n$. The approach is in the same spirit as process modeling approaches using basis functions and kernel convolutions, that is, specifications which attempt to facilitate computations through lower dimensional process representations, i.e.

$$w(s) = c'_{s_0} \Sigma_{w^*}^{-1} w_{rk}^* \quad (5)$$

where $c_{s_0} = (C(s_0, s_1^*), \dots, C(s_0, s_K^*))'$, and then $c'_{s_0} \Sigma_{w^*}^{-1}$ plays the same role as h in (4), and thus can be regarded as some basis functions.

Comment:

- 1) One advertised advantage of using the GPP approach as opposed to FRK or LatticeKrig is that the GPP basis functions are completely determined by the choice of covariance function $C(\cdot, \cdot)$, and note that the subsequent Multiresolution approximations is in step with GPP in this regard.
- 2) At the same time, however, when $C(\cdot, \cdot)$ is governed by unknown parameters (which is nearly always the case) the GPP basis functions need to be calculated iteratively rather than once as in FRK or LatticeKrig which will subsequently increase computation time.

4.1.3 Multi-resolution approximations

In contrast to FRK or LatticeKrig, the multi-resolution approximation (MRA) basis functions and the prior distribution of the corresponding weights w_r^* are chosen using the predictive-process approach to automatically adapt to any given covariance function C_r , and so the MRA can adjust flexibly to a desired spatial smoothness and dependence structure. The MRA allows the number of basis functions to be

approximately the same as the data by the two ways to do. The one is by increasing sparsity of the covariance matrices of the corresponding weights (achieved by tapering method, see [Katzfuss and Hammerling, 2017]), the other is by recursively partitioning the spatial domain (see [Katzfuss, 2017]).

Comment: Two obvious advantages are parallel computing and capturing spatial dependencies at different scales.

4.2 Sparse covariance methods

4.2.1 Tapering

Including **tapering for estimation** (e.g., approximation log-likelihood function by a tapered covariance. See Kaufman et al. [2008]) and **tapering for Kriging** (or for interpolation or for prediction, e.g., approximation Kriging equation by replacing the original covariance by a tapered version. See Furrer et al. [2006] and some R packages: `spam`, 2010; `KriSp`, 2006; `fields`, 2017).

Comment: The covariance tapering has shown great computational gains, but it also has its own drawbacks.

- 1) The covariance tapering may not be effective in accounting for **spatial dependence with long range**.
- 2) The accuracy of the tapering approximation for **nonstationary problems** remains an open question, and the application of tapering techniques to **multivariate random fields** (e.g., [Bevilacqua et al., 2016]) remains to be explored due to the lack of flexible compactly supported cross-covariance functions, see Porcu et al. [2012].

4.2.2 Spatial partitioning

By dividing region \mathbf{D} into m disjoint subregions ($d = 1, 2, \dots, m$), and then the modeling approach based on spatial partitioning is to again assume the model in (6) but take on the assumption of independence between observations across subregions, and then

$$\mathbf{y}_d = \beta \mathbf{X}_d + \mathbf{H}_d \mathbf{w} + \xi_d + \epsilon_d \quad (6)$$

where \mathbf{H}_d is a matrix of spatial basis function for subregions d .

Comment:

- 1) Notice that, in (6) each subregion shares common β and \mathbf{w} parameters which allows smoothing across subregions in spite of the independence assumption. Further, the assumption of independence across subregions effectively creates a block-diagonal structure for Σ and allows the likelihood to be computed in parallel (with one node per subregion), thereby facilitating computation.
- 2) The key to implementing the spatial partitioning approach is the choice of partition and the literature is replete with various options. A priori methods to define the spatial partitioning include partitioning the region into equal areas ([Sang et al., 2011]), partitioning based on centroid clustering ([Kim et al., 2005]), hierarchical clustering based on spatial gradients ([Heaton et al.,

2017]). Alternatively, model-based approaches to spatial partitioning include treed regression([Konomi et al., 2014]) and mixture modeling (Neelon et al. [2014]), but these approaches typically require more computation.

4.3 Sparse precision methods

The sparse precision methods focus on basis function and conditional likelihood.

4.3.1 *LatticeKrig*

LatticeKrig (LK, [Nychka et al., 2015]) uses nearly the same setup as is employed by FRK. Here is just a list of the differences:

- 1) FRK uses families of **bisquare** basis functions allowing different resolutions that are organized on **irregular** basic areal units (BAUs), and LK uses families of **radial** basis functions that are organized on **regular** grids of increasing resolution.
- 2) Covariance matrix, Σ_r , of the coefficients w_r^* in (4) is constrained by a exponential function, the exponential function is defined on the centroids of the BAUs. However, precision matrix, Q_r , of the coefficients w_r^* in LK is constrained to a spatial autoregressive (SAR, defined on the regular lattice) model and forced to admit some sparse structures by following the ideas of Lindgren et al. [2011].
- 3) Furthermore, because Q_r is sparse, **LK can set K to be very large (e.g., $K > n$) without much additional computational cost.** However, the FRK does not achieve this effect.

4.3.2 *Gaussian Markov random field approximations by SPDE*

Rue and Held [2005] rigorously suggest to approximate Gaussian random fields through Markov fields with a huge increase in speed for the simulations and demonstrated empirically that GMRFs could closely approximate some commonly used covariance functions in geostatistics, and whereafter Lindgren et al. [2011] show the construction of the corresponding GMRFs can be used to represent the Matérn field on a triangulated lattice (i.e. the finite element construction).

Similar to FRK or LatticeKrig, the spatial process $w(s)$ in model (1) is represented by the basis function, this method use piecewise linear functions on a triangulation of the domain as basis functions $h(\cdot)$ in (4), and then this yields sparse matrices C and G such that the appropriate precision matrix for the weights w^* is given by (see [Blangiardo and Cameletti, 2015])

$$\Omega = \tau^2(\kappa^4 C + 2\kappa^2 G + GC^{-1}G).$$

One particular advantage of this approach is that assigning the Gaussian distribution $w^* \sim N(0, \Omega^{-1})$ can generate continuously defined functions $\mathcal{X}(s) = \sum_{i=1}^m h_i(s)w_k^*$ that are approximative solutions to the following SPDE (in a stochastically weak sense):

$$(\kappa^2 - \Delta)^{\alpha/2} (\tau(\mathcal{X}(s))) = \mathcal{W}, \quad s \in D,$$

where Δ is the Laplacian, κ is the spatial scale parameter, α controls the smoothness of the realisations, τ controls the variance. The right-hand side of the equation, $\mathcal{W}(s)$, is a Gaussian spatial white noise process. Note that the stationary solutions on \mathcal{R}^d is the stationary Gaussian field $\mathcal{X}(s)$ with Matérn covariances function given by

$$\text{Cov}(\mathcal{X}(s_i), \mathcal{X}(s_j)) = \frac{\sigma^2}{\Gamma(\lambda)2^{\lambda-1}} (\kappa \|s_i - s_j\|)^\lambda K_\lambda(\kappa \|s_i - s_j\|) \quad (7)$$

The drawback of [Lindgren et al. \[2011\]](#) is that we can only find the explicit form of GMRFs for those Gaussian random fields that have a Matérn covariance structure at certain integer smoothnesses. Subsequently, [Bolin and Kirchner \[2020\]](#) extend this method to fractional order by the rational approximation.

Comment:

- 1) Unlike FRK or LatticeKrig where we need pre-define basis functions, the basis functions $h(\cdot)$ of GMRFs approximation are **automatically control or chosen by triangulation of the domain**, and thus constrained to be **sparse** generally.
- 2) The numerical factorization of the precision matrix using [INLA](#) can be done at a typical cost of $O(n^{3/2})$ for two-dimensional GMRFs.
- 3) SPDE can be extended to model Matérn covariances on the sphere, nonstationary locally isotropic Gaussian random fields (e.g., by using spatially varying parameters), Gaussian random fields with oscillating correlation functions, and non-isotropic fields.
- 4) The SPDE approach also facilitates extensions to multivariate random fields, see [Hu et al. \[2013\]](#), [Hu and Steinsland \[2016\]](#), [Bolin and Wallin \[2020\]](#).

4.3.3 *The nearest neighbor Gaussian process*

Similar to the pre-defined knots for Gaussian predictive processes, the starting point of nearest neighbor Gaussian process (NNGP) approach is to choose a fixed collection (e.g., \mathcal{S}) of distinct locations in \mathcal{D} , where \mathcal{S} need not coincide with or be apart of the observed locations, so its size k need not equal the size of the dataset n (or even larger than the size n). The set \mathcal{S} is called as reference set by [Datta et al. \[2016a\]](#).

A directed acyclic graph is defined on the reference set \mathcal{S} , and then the joint distribution of spatial process $w(s)$ from the reference set is represented by the product of conditional densities which is motivated by Vecchia's approximation [[Vecchia, 1988](#)] ideas, where a careful choice of suitable conditional sets is required, and this conditional set is constrained to be the m nearest neighbors by the NNGP approach and thereby facilitating computation in estimation and prediction problem (for more details on those problems, see [Appendix](#) of [Datta et al. \[2016a\]](#) and [[Finley et al., 2017](#)] and [Finley et al. \[2019\]](#)). Based on these results, the NNGP can be well-defined.

- 1) Total operations is $O((n+k)m^3)$ where $m(\approx 20)$ is the size of conditioning set or neighbor set, and several processes can run in parallel (e.g, computations of weights in Kriging equation (3)).
- 2) The other major advantage is that the precision matrix of well-defined NNGP is sparse with at most $km(m+1)/2$ nonzero entries.

- 3) NNGP can also be extended to large spatio-temporal data ([Datta et al. \[2016b\]](#)) and non-stationary process([Konomi et al. \[2019\]](#)), [\[Risser and Turek, 2020\]](#).

5 TWO EXPLORATORY LIMITATIONS OF THE NNGP APPROACH

5.1 Exploratory analysis

- 1) Problem 1: Since the joint distribution of spatial process $w(s)$ from the reference set is represented by using the product of conditional densities and those conditional sets are constrained to be the m nearest neighbors, some **large scale dependence** information maybe lost.
- 2) Problem 2: From an application point of view, the reference set S cannot be infinite because we need to solve neighbor set for every location, so the performance of the NNGP approximation depends on the size of the spatial dependence range relative to the spacing of the reference set, that maybe lead to the quality of the NNGP approximation gets worse when **the spatial dependence range gets shorter**.

We considered model (1) with $\beta = 0, \sigma^2 = 5, \tau^2 = 1$, and the Matérn correlation function for the spatial random effects, w , with a constant smoothness parameter $\nu = 0.5$ and varying spatial range parameters.

Three distance criteria:

$$D_{KL}(\hat{\Sigma}, \Sigma) = \frac{1}{2} \left[\text{trace} \left(\hat{\Sigma}^{-1} \Sigma \right) - p + \log |\hat{\Sigma}| - \log |\Sigma| \right] \quad (8)$$

$$D_B(\hat{\Sigma}, \Sigma) = \frac{1}{2} \log |\tilde{\Sigma}| - \frac{1}{4} \log |\hat{\Sigma}| - \frac{1}{4} \log |\Sigma|, \quad \tilde{\Sigma} = [\Sigma + \hat{\Sigma}]/2 \quad (9)$$

$$D_F(\hat{\Sigma}, \Sigma) = \sqrt{\text{trace} \left[(\hat{\Sigma} - \Sigma)(\hat{\Sigma} - \Sigma)' \right]} \quad (10)$$




Figure 1: The K-L distance, the Bhattacharyya distance and the Frobenius norm from the estimated covariance matrix to the true covariance matrix for the Matérn family with smoothness parameters $\nu = \frac{1}{2}$ and different spatial range parameters ϕ . And this true covariance matrix at 500 random locations from $[0, 100] \times [0, 100]$ was estimated by the NNGP with $m = 20$.

Figure 1 shows that **the large-scale spatial dependencies can not be well-captured** by the NNGP.

5.2 Improvement project

- 1) Ideas for Problem 1: A NNGP $w(s)$ used to capture the local, small scale dependence structure plus a smooth NNGP $f(s)$ used to capture the large scale dependence structure.

$$y(s) = w(s) + f(s) + \epsilon(s) \quad (11)$$

where $w = (w(s_1), \dots, w(s_n))' \sim \text{NNGP}(\mathbf{0}, \tilde{\mathbf{C}}_1(\cdot; \theta_1))$; $f(s) = \sum_{k=1}^K h_k(s)w_k^*$, and let $w^* = (w_1^*, \dots, w_K^*)' \sim \text{NNGP}(\mathbf{0}, \tilde{\mathbf{C}}_2(\cdot; \theta_2))$ and h_k is **automatically chosen by a given covariance function**, i.e. $\tilde{c}'(k, \mathcal{N}(k))\tilde{\mathbf{C}}_2^{-1}(\mathcal{N}(k), \mathcal{N}(k))$; $\epsilon = (\epsilon(s_1), \dots, \epsilon(s_n))' \sim N(\mathbf{0}, \tau^2 \mathbf{I})$. And thereby, we have

$$y \sim \text{GP}(\mathbf{0}, \Sigma(\cdot; \theta)) \quad (12)$$

where $\Sigma(\cdot; \theta) = \tilde{\mathbf{C}}_1(\cdot; \theta_1) + \mathbf{H}\tilde{\mathbf{C}}_2(\cdot; \theta_2)\mathbf{H}' + \tau^2 \mathbf{I}$.

- 2) Ideas for Problem 2: Similar to Sang and Huang [2012],

$$y(s) = w_l(s) + w_s(s) + \epsilon(s) \quad (13)$$

where w_l is a NNGP and w_s is a residual process with covariance function constrained by tapered function.

- 3) A more thorough solution for both problems: Similar to LatticeKrig, a multiresolution NNGP approximation is

$$y(s) = \sum_{r=1}^R \sum_{k=1}^{K_r} h_{rk}(s)w_{rk}^* + \epsilon(s) \quad (14)$$

where $w_r^* = (w_{r,1}^*, \dots, w_{r,K_r}^*)' \sim \text{NNGP}(\mathbf{0}, \tilde{\mathbf{C}}_r(\cdot; \theta_r))$.

REFERENCES

- S. Banerjee, A. E. Gelfand, A. O. Finley, and H. Sang. Gaussian predictive process models for large spatial data sets. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 70(4):825–848, 2008. URL [https://chenyw68.github.io/Literature/\[2008\]Gaussianpredictiveprocessmodelsforlargespatialdatasets.pdf](https://chenyw68.github.io/Literature/[2008]Gaussianpredictiveprocessmodelsforlargespatialdatasets.pdf).
- M. Bevilacqua, A. Fassò, C. Gaetan, E. Porcu, and D. Velandia. Covariance tapering for multivariate gaussian random fields estimation. *Statistical Methods & Applications*, 25(1):21–37, 2016. URL [https://chenyw68.github.io/Literature/\[2015\]CovariancetaperingformultivariateGaussianrandomfieldsestimation.pdf](https://chenyw68.github.io/Literature/[2015]CovariancetaperingformultivariateGaussianrandomfieldsestimation.pdf).
- M. Blangiardo and M. Cameletti. *Spatial and spatio-temporal Bayesian models with R-INLA*. John Wiley & Sons, 2015. URL [https://chenyw68.github.io/Literature/\[2015\]SpatialandSpatiotemporalBayesianModelswithRINLA.pdf](https://chenyw68.github.io/Literature/[2015]SpatialandSpatiotemporalBayesianModelswithRINLA.pdf).

- D. Bolin and K. Kirchner. The rational spde approach for gaussian random fields with general smoothness. *Journal of Computational and Graphical Statistics*, 29(2):274–285, 2020. URL [https://chenyw68.github.io/Literature/\[2020\]TheRationalSPDEApproachforGaussianRandomFieldsWithGeneralSmoothness.pdf](https://chenyw68.github.io/Literature/[2020]TheRationalSPDEApproachforGaussianRandomFieldsWithGeneralSmoothness.pdf).
- D. Bolin and J. Wallin. Multivariate type g matérn stochastic partial differential equation random fields. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 82(1):215–239, 2020. URL [https://chenyw68.github.io/Literature/\[2020\]MultivariatetypeGMaternstochasticpartialdifferentialsolutionrandomfields.pdf](https://chenyw68.github.io/Literature/[2020]MultivariatetypeGMaternstochasticpartialdifferentialsolutionrandomfields.pdf).
- N. Cressie and G. Johannesson. Fixed rank kriging for very large spatial data sets. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 70(1):209–226, 2008. URL [https://chenyw68.github.io/Literature/\[2008\]Fixedrankkrigingforverylargespatialdatasets.pdf](https://chenyw68.github.io/Literature/[2008]Fixedrankkrigingforverylargespatialdatasets.pdf).
- N. A. Cressie and G. Johannesson. Spatial prediction for massive datasets. *Proceedings of the Australian Academy of Science Elizabeth and Frederick White Conference*, 2006. URL [https://chenyw68.github.io/Literature/\[2006\]Spatialpredictionformassivedatasets.pdf](https://chenyw68.github.io/Literature/[2006]Spatialpredictionformassivedatasets.pdf).
- A. Datta, S. Banerjee, A. O. Finley, and A. E. Gelfand. Hierarchical nearest-neighbor gaussian process models for large geostatistical datasets. *Journal of the American Statistical Association*, 111(514):800–812, 2016a. URL [https://chenyw68.github.io/Literature/\[2016\]Hierarchicalnearest-neighborGaussianprocessmodelsforlargegeostatisticaldatasets.pdf](https://chenyw68.github.io/Literature/[2016]Hierarchicalnearest-neighborGaussianprocessmodelsforlargegeostatisticaldatasets.pdf).
- A. Datta, S. Banerjee, A. O. Finley, N. A. Hamm, and M. Schaap. Nonseparable dynamic nearest neighbor gaussian process models for large spatio-temporal data with an application to particulate matter analysis. *The annals of applied statistics*, 10(3):1286, 2016b. URL [https://chenyw68.github.io/Literature/\[2016\]NonseparabledynamicNNGPmodelsforlargespatio-temporaldata.pdf](https://chenyw68.github.io/Literature/[2016]NonseparabledynamicNNGPmodelsforlargespatio-temporaldata.pdf).
- A. O. Finley, H. Sang, S. Banerjee, and A. E. Gelfand. Improving the performance of predictive process modeling for large datasets. *Computational statistics & data analysis*, 53(8):2873–2884, 2009. URL [https://chenyw68.github.io/Literature/\[2009\]Improvingtheperformanceofpredictiveprocessmodelingforlargedatasets.pdf](https://chenyw68.github.io/Literature/[2009]Improvingtheperformanceofpredictiveprocessmodelingforlargedatasets.pdf).
- A. O. Finley, A. Datta, B. C. Cook, D. C. Morton, H. E. Andersen, and S. Banerjee. Applying nearest neighbor gaussian processes to massive spatial data sets forest canopy height prediction across tanana valley alaska.”. *arXiv preprint arXiv:1702.00434*, 2017. URL [https://chenyw68.github.io/Literature/\[2017\]ApplyingNearestNeighborGaussianProcessestoMassiveSpatialDataSets.pdf](https://chenyw68.github.io/Literature/[2017]ApplyingNearestNeighborGaussianProcessestoMassiveSpatialDataSets.pdf).
- A. O. Finley, A. Datta, B. D. Cook, D. C. Morton, H. E. Andersen, and S. Banerjee. Efficient algorithms for bayesian nearest neighbor gaussian processes. *Journal of Computational and Graphical Statistics*, 28(2):401–414, 2019. URL [https://chenyw68.github.io/Literature/\[2019\]Efficientalgorithmsforbayesiannearestneighborgaussianprocesses.pdf](https://chenyw68.github.io/Literature/[2019]Efficientalgorithmsforbayesiannearestneighborgaussianprocesses.pdf).
- M. Fuentes. Approximate likelihood for large irregularly spaced spatial data. *Journal of the American Statistical Association*, 102

- (477):321–331, 2007. URL [https://chenyw68.github.io/Literature/\[2007\]ApproximateLikelihoodforLargeIrregularlySpacedSpatialData.pdf](https://chenyw68.github.io/Literature/[2007]ApproximateLikelihoodforLargeIrregularlySpacedSpatialData.pdf).
- R. Furrer, M. G. Genton, and D. Nychka. Covariance tapering for interpolation of large spatial datasets. *Journal of Computational and Graphical Statistics*, 15(3):502–523, 2006. URL [https://chenyw68.github.io/Literature/\[2006\]Covariantetaperingforinterpolationoflargespatialdatasets.pdf](https://chenyw68.github.io/Literature/[2006]Covariantetaperingforinterpolationoflargespatialdatasets.pdf).
- R. B. Gramacy and D. W. Apley. Local gaussian process approximation for large computer experiments. *Journal of Computational and Graphical Statistics*, 24(2):561–578, 2015. URL [https://chenyw68.github.io/Literature/\[2014\]LocalGaussianProcessApproximationforLargeComputerExperiments.pdf](https://chenyw68.github.io/Literature/[2014]LocalGaussianProcessApproximationforLargeComputerExperiments.pdf).
- J. Guinness. Spectral density estimation for random fields via periodic embeddings. *Biometrika*, 106(2):267–286, 2019. URL [https://chenyw68.github.io/Literature/\[2020\]SpectralDensityEstimationforRandomFieldsviaPeriodicEmbeddings.pdf](https://chenyw68.github.io/Literature/[2020]SpectralDensityEstimationforRandomFieldsviaPeriodicEmbeddings.pdf).
- J. Guinness and M. Fuentes. Circulant embedding of approximate covariances for inference from gaussian data on large lattices. *Journal of computational and Graphical Statistics*, 26(1):88–97, 2017. URL [https://chenyw68.github.io/Literature/\[2017\]CirculantEmbeddingofApproximateCovariancesforInferenceFromGaussianData.pdf](https://chenyw68.github.io/Literature/[2017]CirculantEmbeddingofApproximateCovariancesforInferenceFromGaussianData.pdf).
- M. J. Heaton, W. F. Christensen, and M. A. Terres. Nonstationary gaussian process models using spatial hierarchical clustering from finite differences. *Technometrics*, 59(1):93–101, 2017. URL [https://chenyw68.github.io/Literature/\[2017\]NonstationaryGPModelsUsingSpatialHierarchicalClusteringfromFiniteDifferences.pdf](https://chenyw68.github.io/Literature/[2017]NonstationaryGPModelsUsingSpatialHierarchicalClusteringfromFiniteDifferences.pdf).
- M. J. Heaton, A. Datta, A. O. Finley, R. Furrer, J. Guinness, R. Guhaniyogi, F. Gerber, R. B. Gramacy, D. Hammerling, M. Katzfuss, et al. A case study competition among methods for analyzing large spatial data. *Journal of Agricultural, Biological and Environmental Statistics*, 24(3):398–425, 2019. URL [https://chenyw68.github.io/Literature/\[2018\]Acasestudycompetitionamongmethodsforanalyzinglargespatialdata.pdf](https://chenyw68.github.io/Literature/[2018]Acasestudycompetitionamongmethodsforanalyzinglargespatialdata.pdf).
- X. Hu and I. Steinsland. Spatial modeling with system of stochastic partial differential equations. *Wiley Interdisciplinary Reviews: Computational Statistics*, 8(2):112–125, 2016. URL [https://chenyw68.github.io/Literature/\[2016\]Spatialmodelingwithsystemofstochasticpartialdifferentialequations.pdf](https://chenyw68.github.io/Literature/[2016]Spatialmodelingwithsystemofstochasticpartialdifferentialequations.pdf).
- X. Hu, D. Simpson, F. Lindgren, and H. Rue. Multivariate gaussian random fields using systems of stochastic partial differential equations. *arXiv preprint arXiv:1307.1379*, 2013. URL [https://chenyw68.github.io/Literature/\[2013\]MultivariateGaussianrandomfieldsusingsystemsofstochasticpartialdifferentialequations.pdf](https://chenyw68.github.io/Literature/[2013]MultivariateGaussianrandomfieldsusingsystemsofstochasticpartialdifferentialequations.pdf).
- M. Jurek and M. Katzfuss. Hierarchical sparse cholesky decomposition with applications to high-dimensional spatio-temporal filtering. *arXiv preprint arXiv:2006.16901*, 2020. URL [https://chenyw68.github.io/Literature/\[2020\]HierarchicalsparseCholeskydecompositionwithapplicationstospatio-temporalfiltering.pdf](https://chenyw68.github.io/Literature/[2020]HierarchicalsparseCholeskydecompositionwithapplicationstospatio-temporalfiltering.pdf).

- M. Katzfuss. A multi-resolution approximation for massive spatial datasets. *Journal of the American Statistical Association*, 112(517):201–214, 2017. URL [https://chenyw68.github.io/Literature/\[2017\]AMulti-ResolutionApproximationforMassiveSpatialDatasets.pdf](https://chenyw68.github.io/Literature/[2017]AMulti-ResolutionApproximationforMassiveSpatialDatasets.pdf).
- M. Katzfuss and D. Hammerling. Parallel inference for massive distributed spatial data using low-rank models. *Statistics and Computing*, 27(2):363–375, 2017. URL [https://chenyw68.github.io/Literature/\[2017-formal\]Parallelinferenceformassivedistributedspatialdatausinglow-rankmodels.pdf](https://chenyw68.github.io/Literature/[2017-formal]Parallelinferenceformassivedistributedspatialdatausinglow-rankmodels.pdf).
- M. Katzfuss, J. Guinness, W. Gong, and D. Zilber. Vecchia approximations of gaussian-process predictions. *Journal of Agricultural, Biological and Environmental Statistics*, 25(3):383–414, 2020. URL [https://chenyw68.github.io/Literature/\[2020\]VecchiaApproximationsofGaussian-ProcessPredictions.pdf](https://chenyw68.github.io/Literature/[2020]VecchiaApproximationsofGaussian-ProcessPredictions.pdf).
- C. G. Kaufman, M. J. Schervish, and D. W. Nychka. Covariance tapering for likelihood-based estimation in large spatial data sets. *Journal of the American Statistical Association*, 103(484):1545–1555, 2008. URL [https://chenyw68.github.io/Literature/\[2008\]CovarianceTaperingforLikelihoodBasedEstimationinLargeSpatialDataSets.pdf](https://chenyw68.github.io/Literature/[2008]CovarianceTaperingforLikelihoodBasedEstimationinLargeSpatialDataSets.pdf).
- H.-M. Kim, B. K. Mallick, and C. Holmes. Analyzing nonstationary spatial data using piecewise gaussian processes. *Journal of the American Statistical Association*, 100(470):653–668, 2005. URL [https://chenyw68.github.io/Literature/\[2005\]AnalyzingnonstationaryspatialdatausingpiecewiseGaussianprocesses.pdf](https://chenyw68.github.io/Literature/[2005]AnalyzingnonstationaryspatialdatausingpiecewiseGaussianprocesses.pdf).
- B. Konomi, A. Hanandeh, P. Ma, and E. Kang. Computationally efficient nonstationary nearest-neighbor gaussian process models using data-driven techniques. *Environmetrics*, 30(8):e2571, 2019. URL [https://chenyw68.github.io/Literature/\[2020\]ComputationallyefficientnonstationaryNNGPmodelsusingdata-driventechiniques.pdf](https://chenyw68.github.io/Literature/[2020]ComputationallyefficientnonstationaryNNGPmodelsusingdata-driventechiniques.pdf).
- B. A. Konomi, H. Sang, and B. K. Mallick. Adaptive bayesian nonstationary modeling for large spatial datasets using covariance approximations. *Journal of Computational and Graphical Statistics*, 23(3):802–829, 2014. URL [https://chenyw68.github.io/Literature/\[2014\]AdaptiveBayesianNonstationaryModelingforLargeSpatialDatasets.pdf](https://chenyw68.github.io/Literature/[2014]AdaptiveBayesianNonstationaryModelingforLargeSpatialDatasets.pdf).
- F. Lindgren, H. Rue, and J. Lindström. An explicit link between gaussian fields and gaussian markov random fields: the stochastic partial differential equation approach. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(4):423–498, 2011. URL [https://chenyw68.github.io/Literature/\[2011\]AnexplicitlinkbetweenGFandGMRFstheSPDEapproach.pdf](https://chenyw68.github.io/Literature/[2011]AnexplicitlinkbetweenGFandGMRFstheSPDEapproach.pdf).
- P. Ma and E. L. Kang. A fused gaussian process model for very large spatial data. *Journal of Computational and Graphical Statistics*, pages 1–11, 2020. URL [https://chenyw68.github.io/Literature/\[2020\]AFusedGaussianProcessModelforVeryLargeSpatialData.pdf](https://chenyw68.github.io/Literature/[2020]AFusedGaussianProcessModelforVeryLargeSpatialData.pdf).
- B. Neelon, A. E. Gelfand, and M. L. Miranda. A multivariate spatial mixture model for areal data: examining regional differences in standardized

- test scores. *Journal of the Royal Statistical Society. Series C, Applied statistics*, 63(5):737, 2014. URL [https://chenyw68.github.io/Literature/\[2014\]Amultivariatespatialmixturemodelforarealdata.pdf](https://chenyw68.github.io/Literature/[2014]Amultivariatespatialmixturemodelforarealdata.pdf).
- D. Nychka, S. Bandyopadhyay, D. Hammerling, F. Lindgren, and S. Sain. A multiresolution gaussian process model for the analysis of large spatial datasets. *Journal of Computational and Graphical Statistics*, 24(2):579–599, 2015. URL [https://chenyw68.github.io/Literature/\[2015\]AMultiresolutionGaussianProcessModelfortheAnalysisofLargeSpatialDatasets.pdf](https://chenyw68.github.io/Literature/[2015]AMultiresolutionGaussianProcessModelfortheAnalysisofLargeSpatialDatasets.pdf).
- E. Porcu, J.-M. Montero, and M. Schlather. *Advances and challenges in space-time modelling of natural events*, volume 207. Springer Science & Business Media, 2012. URL [https://chenyw68.github.io/Literature/\[2012\]AdvancesandChallengesinSpace-timeModellingofNaturalEvents.pdf](https://chenyw68.github.io/Literature/[2012]AdvancesandChallengesinSpace-timeModellingofNaturalEvents.pdf).
- M. D. Risser and D. Turek. Bayesian inference for high-dimensional nonstationary gaussian processes. *Journal of Statistical Computation and Simulation*, pages 1–27, 2020. URL [https://chenyw68.github.io/Literature/\[2020-formal\]BayesianinferenceforhighdimensionalnonstationaryGaussianprocesses.pdf](https://chenyw68.github.io/Literature/[2020-formal]BayesianinferenceforhighdimensionalnonstationaryGaussianprocesses.pdf).
- H. Rue and L. Held. *Gaussian Markov random fields: theory and applications*. CRC press, 2005. URL [https://chenyw68.github.io/Literature/\[2005\]Gaussianmarkovrandomfieldstheoryandapplications.pdf](https://chenyw68.github.io/Literature/[2005]Gaussianmarkovrandomfieldstheoryandapplications.pdf).
- H. Sang and J. Z. Huang. A full scale approximation of covariance functions for large spatial data sets. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 74(1):111–132, 2012. URL [https://chenyw68.github.io/Literature/\[2012\]Afullscaleapproximationofcovariancefunctionsforlargespatialdatasets.pdf](https://chenyw68.github.io/Literature/[2012]Afullscaleapproximationofcovariancefunctionsforlargespatialdatasets.pdf).
- H. Sang, M. Jun, and J. Z. Huang. Covariance approximation for large multivariate spatial data sets with an application to multiple climate model errors. *The Annals of Applied Statistics*, pages 2519–2548, 2011. URL [https://chenyw68.github.io/Literature/\[2011\]Covarianceapproximationforlargemultivariatespatialdatasets.pdf](https://chenyw68.github.io/Literature/[2011]Covarianceapproximationforlargemultivariatespatialdatasets.pdf).
- M. L. Stein. A modeling approach for large spatial datasets. *Journal of the Korean Statistical Society*, 37:3–10, 2008. URL [https://chenyw68.github.io/Literature/\[2008\]Amodelingapproachforlargespatialdatasets.pdf](https://chenyw68.github.io/Literature/[2008]Amodelingapproachforlargespatialdatasets.pdf).
- A. V. Vecchia. Estimation and model identification for continuous spatial processes. *Journal of the Royal Statistical Society: Series B (Methodological)*, 50(2):297–312, 1988.
- D. Zilber and M. Katzfuss. Vecchia-laplace approximations of generalized gaussian processes for big non-gaussian spatial data. *Computational Statistics and Data Analysis*, 153:107081, 2020. URL [https://chenyw68.github.io/Literature/\[2020\]Vecchialaplaceapproximationsofgeneralizedgaussianprocesses.pdf](https://chenyw68.github.io/Literature/[2020]Vecchialaplaceapproximationsofgeneralizedgaussianprocesses.pdf).