

Systems of Ordinary Differential Equations > Nonlinear Systems of Two Equations

12.
$$x_{tt}'' + a(t)x = x^{-3}f(y/x)$$
, $y_{tt}'' + a(t)y = y^{-3}g(y/x)$.

Generalized Ermakov (Yermakov) system.

1°. First integral:

$$\frac{1}{2}(xy_t' - yx_t')^2 + \int_{-\infty}^{y/x} \left[uf(u) - u^{-3}g(u) \right] du = C,$$

where C is an arbitrary constant.

 2° . Let $\varphi = \varphi(t)$ is a nontrivial solution of the second-order linear differential equation

$$\varphi_{tt}^{"} + a(t)\varphi = 0. \tag{1}$$

Then, the transformation

$$\tau = \int \frac{dt}{\varphi^2(t)}, \quad u = \frac{x}{\varphi(t)}, \quad v = \frac{y}{\varphi(t)}$$
(2)

leads to the autonomous system of equations

$$u_{\tau\tau}^{"} = u^{-3} f(v/u), \quad v_{\tau\tau}^{"} = v^{-3} g(v/u).$$
 (3)

3°. Particular solution of system (3) is

$$u = A\sqrt{C_2\tau^2 + C_1\tau + C_0}, \quad v = Ak\sqrt{C_2\tau^2 + C_1\tau + C_0}, \quad A = \left[\frac{f(k)}{C_0C_2 - \frac{1}{4}C_1^2}\right]^{1/4},$$

where C_0 , C_1 , and C_2 are arbitrary constants, and k is a root of the algebraic (transcendental) equation

$$k^4 f(k) = q(k).$$

References

Ray, J. R. and Reid, J. L., More exact invariants for the time dependent harmonic oscillator, *Phys. Lett., Ser. A*, Vol. 71, pp. 317–319, 1979.

Athorne, C., Rational Ermakov systems of Fuchsian type, Dept. of Math. Gardens, Glasgow Univ., Preprint N91/41, 1991.

Govinger, K. S., Athorne, C., and Leach, P. J. L., The algebraic structure of generalized Ermakov systems of three dimensions, *J. Phys. A: Math. Gen.*, Vol. 26, pp. 4035–4046, 1993.

Berkovich, L. M., Factorization and Transformations of Ordinary Differential Equations [in Russian], Regulyarnaya i Khaoticheskaya Dinamika, Moscow, 2002.

Copyright © 2004 Andrei D. Polyanin

http://eqworld.ipmnet.ru/en/solutions/sysode/sode0312.pdf