

STAT 423/523 Statistical Methods for Engineers and Scientists

Lecture 4: Hypothesis Testing I

Chencheng Cai

Washington State University

Hypothesis Testing

A **statistical hypothesis** is a statement or assumption about one or more population parameters.

Hypothesis Testing

A **statistical hypothesis** is a statement or assumption about one or more population parameters.

- ▶ The average height of WSU students is 68 inches.
- ▶ The proportion of defective items produced by a machine is greater than 0.10.
- ▶ The average commute time of WSU students is the same as the UI students.

Hypothesis Testing

A **statistical hypothesis** is a statement or assumption about one or more population parameters.

- ▶ The average height of WSU students is 68 inches.
- ▶ The proportion of defective items produced by a machine is greater than 0.10.
- ▶ The average commute time of WSU students is the same as the UI students.

A **hypothesis-testing procedure** usually involves two contradictory hypotheses.

Hypothesis Testing

A **statistical hypothesis** is a statement or assumption about one or more population parameters.

- ▶ The average height of WSU students is 68 inches.
- ▶ The proportion of defective items produced by a machine is greater than 0.10.
- ▶ The average commute time of WSU students is the same as the UI students.

A **hypothesis-testing procedure** usually involves two contradictory hypotheses.

- ▶ The average height of WSU students is 68 inches.
- ▶ v.s. The average height of WSU students is NOT 68 inches.

Hypothesis Testing

The **null hypothesis** H_0 is the claim that is initially assumed to be true.

The **alternative hypothesis** H_a is the claim that we are trying to find evidence for.

Hypothesis Testing

The **null hypothesis** H_0 is the claim that is initially assumed to be true.

The **alternative hypothesis** H_a is the claim that we are trying to find evidence for.

Example:

Let μ be the average height of WSU students.

To test whether μ is exactly 68 inches:

Hypothesis Testing

The **null hypothesis** H_0 is the claim that is initially assumed to be true.

The **alternative hypothesis** H_a is the claim that we are trying to find evidence for.

Example:

Let μ be the average height of WSU students.

To test whether μ is exactly 68 inches:

- ▶ $H_0: \mu = 68$ inches.
- ▶ $H_a: \mu \neq 68$ inches.

Hypothesis Testing

The **null hypothesis** H_0 is the claim that is initially assumed to be true.

The **alternative hypothesis** H_a is the claim that we are trying to find evidence for.

Example:

Let μ be the average height of WSU students.

To test whether μ is exactly 68 inches:

- ▶ $H_0: \mu = 68$ inches.
- ▶ $H_a: \mu \neq 68$ inches.

To test whether μ is greater than or equal to 68 inches:

Hypothesis Testing

The **null hypothesis** H_0 is the claim that is initially assumed to be true.

The **alternative hypothesis** H_a is the claim that we are trying to find evidence for.

Example:

Let μ be the average height of WSU students.

To test whether μ is exactly 68 inches:

- ▶ $H_0: \mu = 68$ inches.
- ▶ $H_a: \mu \neq 68$ inches.

To test whether μ is greater than or equal to 68 inches:

- ▶ $H_0: \mu \geq 68$ inches.
- ▶ $H_a: \mu < 68$ inches.

Hypothesis Testing

A **test of hypothesis** is a procedure for deciding between two contradictory claims.

Hypothesis Testing

A **test of hypothesis** is a procedure for deciding between two contradictory claims.

Two possible outcomes of a hypothesis test:

- ▶ **Reject** H_0 : There is enough evidence against H_0 to support H_a .
- ▶ **Fail to reject** H_0 : There is not enough evidence to support H_a .

Hypothesis Testing

A **test of hypothesis** is a procedure for deciding between two contradictory claims.

Two possible outcomes of a hypothesis test:

- ▶ **Reject** H_0 : There is enough evidence against H_0 to support H_a .
- ▶ **Fail to reject** H_0 : There is not enough evidence to support H_a .

Note: We never “accept” the null hypothesis. We either reject it or fail to reject it.

Hypothesis Testing

A **test of hypothesis** is a procedure for deciding between two contradictory claims.

Two possible outcomes of a hypothesis test:

- ▶ **Reject** H_0 : There is enough evidence against H_0 to support H_a .
- ▶ **Fail to reject** H_0 : There is not enough evidence to support H_a .

Note: We never “accept” the null hypothesis. We either reject it or fail to reject it.

The decision is based on the sample data. (So the decision is a random variable.)

Hypothesis Testing

	H_0 is true	H_0 is false
Reject H_0	Type I error	Correct decision
Fail to reject H_0	Correct decision	Type II error

- ▶ A **Type I error** occurs when we reject H_0 when H_0 is true.
- ▶ A **Type II error** occurs when we fail to reject H_0 when H_0 is false.

Hypothesis Testing

	H_0 is true	H_0 is false
Reject H_0	Type I error	Correct decision
Fail to reject H_0	Correct decision	Type II error

- ▶ A **Type I error** occurs when we reject H_0 when H_0 is true.
- ▶ A **Type II error** occurs when we fail to reject H_0 when H_0 is false.
- ▶ Usually, Type I error and Type II error are inversely related.

Hypothesis Testing

	H_0 is true	H_0 is false
Reject H_0	Type I error	Correct decision
Fail to reject H_0	Correct decision	Type II error

- ▶ A **Type I error** occurs when we reject H_0 when H_0 is true.
- ▶ A **Type II error** occurs when we fail to reject H_0 when H_0 is false.
- ▶ Usually, Type I error and Type II error are inversely related.
- ▶ Type I error can be controlled by the **significance level** α .

$$P(\text{reject} \mid H_0) = \alpha$$

- ▶ α is also called the **size** of the test.

Example

Suppose the heights of WSU students are normally distributed with mean μ and a standard deviation of 2 inches.

Let X_1, \dots, X_{100} be a random sample from the population.

We want to test whether the average height of WSU students is **greater than** 68 inches.

Example

Suppose the heights of WSU students are normally distributed with mean μ and a standard deviation of 2 inches.

Let X_1, \dots, X_{100} be a random sample from the population.

We want to test whether the average height of WSU students is **greater than** 68 inches.

Step 1: Set up the null and alternative hypotheses.

- ▶ $H_0: \mu = 68$ inches.
- ▶ $H_a: \mu > 68$ inches.

Example

Suppose the heights of WSU students are normally distributed with mean μ and a standard deviation of 2 inches.

Let X_1, \dots, X_{100} be a random sample from the population.

We want to test whether the average height of WSU students is **greater than** 68 inches.

Step 1: Set up the null and alternative hypotheses.

▶ $H_0: \mu = 68$ inches.

▶ $H_a: \mu > 68$ inches.

Step 2: Formulate a decision based on the sample.

reject null if $\bar{X} > 68.6$

Example

Suppose the heights of WSU students are normally distributed with mean μ and a standard deviation of 2 inches.

Let X_1, \dots, X_{100} be a random sample from the population.

We want to test whether the average height of WSU students is **greater than** 68 inches.

Step 1: Set up the null and alternative hypotheses.

▶ $H_0: \mu = 68$ inches.

▶ $H_a: \mu > 68$ inches.

Step 2: Formulate a decision based on the sample.

reject null if $\bar{X} > 68.6$

Step 3: Collect the sample data and make a decision.

Example

Suppose the heights of WSU students are normally distributed with mean μ and a standard deviation of 2 inches.

Let X_1, \dots, X_{100} be a random sample from the population.

We want to test whether the average height of WSU students is **greater than** 68 inches.

Step 1: Set up the null and alternative hypotheses.

▶ $H_0: \mu = 68$ inches.

▶ $H_a: \mu > 68$ inches.

Step 2: Formulate a decision based on the sample.

reject null if $\bar{X} > 68.6$

Step 3: Collect the sample data and make a decision.

The significance level of this test is

$$\alpha = P(\text{reject} | H_0) = P(N(68, 0.04) > 68.5) = 1 - \Phi(3).$$

Example

Now consider another test: whether the average height of WSU students is **exactly** 68 inches.

Example

Now consider another test: whether the average height of WSU students is **exactly** 68 inches.

Step 1: Set up the null and alternative hypotheses.

- ▶ $H_0: \mu = 68$ inches.
- ▶ $H_a: \mu \neq 68$ inches.

Example

Now consider another test: whether the average height of WSU students is **exactly** 68 inches.

Step 1: Set up the null and alternative hypotheses.

- ▶ $H_0: \mu = 68$ inches.
- ▶ $H_a: \mu \neq 68$ inches.

Step 2: Formulate a decision based on the sample.

reject null if $\bar{X} > 68.4$ or $\bar{X} < 67.6$

Example

Now consider another test: whether the average height of WSU students is **exactly** 68 inches.

Step 1: Set up the null and alternative hypotheses.

- ▶ $H_0: \mu = 68$ inches.
- ▶ $H_a: \mu \neq 68$ inches.

Step 2: Formulate a decision based on the sample.

reject null if $\bar{X} > 68.4$ or $\bar{X} < 67.6$

Step 3: Collect the sample data and make a decision.

Example

Now consider another test: whether the average height of WSU students is **exactly** 68 inches.

Step 1: Set up the null and alternative hypotheses.

- ▶ $H_0: \mu = 68$ inches.
- ▶ $H_a: \mu \neq 68$ inches.

Step 2: Formulate a decision based on the sample.

reject null if $\bar{X} > 68.4$ or $\bar{X} < 67.6$

Step 3: Collect the sample data and make a decision.

The significance level of this test is

$$\alpha = P(\text{reject} | H_0) = P(N(68, 0.04) > 68.4 \text{ or } N(68, 0.04) < 67.6) = 0.95.$$

Test Statistics

In our previous sample, our decision is based on the sample mean \bar{X} , which is called the test statistic.

A **test statistic** is a function of the sample data whose value is used to make a decision in a hypothesis test.

Test Statistics

In our previous sample, our decision is based on the sample mean \bar{X} , which is called the test statistic.

A **test statistic** is a function of the sample data whose value is used to make a decision in a hypothesis test.

The set of values of the test statistic for which the null hypothesis is rejected is called the **rejection region**.

Test Statistics

In our previous sample, our decision is based on the sample mean \bar{X} , which is called the test statistic.

A **test statistic** is a function of the sample data whose value is used to make a decision in a hypothesis test.

The set of values of the test statistic for which the null hypothesis is rejected is called the **rejection region**.

Example:

- ▶ The rejection region for the test of whether the average height of WSU students is greater than 68 inches is $(68.6, \infty)$.

Test Statistics

In our previous sample, our decision is based on the sample mean \bar{X} , which is called the test statistic.

A **test statistic** is a function of the sample data whose value is used to make a decision in a hypothesis test.

The set of values of the test statistic for which the null hypothesis is rejected is called the **rejection region**.

Example:

- ▶ The rejection region for the test of whether the average height of WSU students is greater than 68 inches is $(68.6, \infty)$.
- ▶ The rejection region for the test of whether the average height of WSU students is exactly 68 inches is $(-\infty, 67.6) \cup (68.4, \infty)$.

Example

A company producing Brand D yogurt would like to increase its market share.

The company sends out a survey to 100 Brand C consumers asking whether they would like to switch brand.

Let p be the proportion of Brand C consumers who would like to switch to Brand D.

Example

A company producing Brand D yogurt would like to increase its market share. The company sends out a survey to 100 Brand C consumers asking whether they would like to switch brand.

Let p be the proportion of Brand C consumers who would like to switch to Brand D.

Step 1: Set up the null and alternative hypotheses.

- ▶ $H_0: p = 0.5.$
- ▶ $H_a: p < 0.5.$

Example

A company producing Brand D yogurt would like to increase its market share. The company sends out a survey to 100 Brand C consumers asking whether they would like to switch brand.

Let p be the proportion of Brand C consumers who would like to switch to Brand D.

Step 1: Set up the null and alternative hypotheses.

► $H_0: p = 0.5.$

► $H_a: p < 0.5.$

Step 2: Formulate a decision based on the sample. Let X be the number of consumers who would like to switch. Rejection region:

$$X \in [0, 37]$$

Example

A company producing Brand D yogurt would like to increase its market share. The company sends out a survey to 100 Brand C consumers asking whether they would like to switch brand.

Let p be the proportion of Brand C consumers who would like to switch to Brand D.

Step 1: Set up the null and alternative hypotheses.

► $H_0: p = 0.5.$

► $H_a: p < 0.5.$

Step 2: Formulate a decision based on the sample. Let X be the number of consumers who would like to switch. Rejection region:

$$X \in [0, 37]$$

Step 3: Collect the sample data and make a decision.

Example

A company producing Brand D yogurt would like to increase its market share. The company sends out a survey to 100 Brand C consumers asking whether they would like to switch brand.

Let p be the proportion of Brand C consumers who would like to switch to Brand D.

Step 1: Set up the null and alternative hypotheses.

► $H_0: p = 0.5.$

► $H_a: p < 0.5.$

Step 2: Formulate a decision based on the sample. Let X be the number of consumers who would like to switch. Rejection region:

$$X \in [0, 37]$$

Step 3: Collect the sample data and make a decision.

The significance level of this test is

$$\alpha = P(\text{reject} | H_0) = P(\text{Binom}(100, 0.5) \leq 37) = 0.006$$