

STAT 423/523 Statistical Methods for Engineers and Scientists

Lecture 2: Point Estimation

Chencheng Cai

Washington State University

Sample Mean

Proposition

Let X_1, X_2, \dots, X_n be a random sample from a population with mean μ and standard deviation σ . Then

- ▶ $E(\bar{X}) = \mu$
- ▶ $Var(\bar{X}) = \frac{\sigma^2}{n}$

In addition, with $T = X_1 + \dots + X_n$, we have

- ▶ $E(T) = n\mu$
- ▶ $Var(T) = n\sigma^2$

Sample Mean

Proposition

Let X_1, X_2, \dots, X_n be a random sample from a population with mean μ and standard deviation σ . Then

- ▶ $E(\bar{X}) = \mu$
- ▶ $Var(\bar{X}) = \frac{\sigma^2}{n}$

In addition, with $T = X_1 + \dots + X_n$, we have

- ▶ $E(T) = n\mu$
- ▶ $Var(T) = n\sigma^2$

Interpretation:

The sample mean's expectation is the population mean, and its variance is the population variance divided by the sample size.

Sample Mean — Concepts

- ▶ **Population:** In statistics, a population is the entire pool from which a statistical sample is drawn. It is the complete set of individuals or objects that we are interested in.
- ▶ **Sample:** A sample is a subset of the population. It is the group of individuals or objects that we actually collect data from.

Sample Mean — Concepts

- ▶ **Population:** In statistics, a population is the entire pool from which a statistical sample is drawn. It is the complete set of individuals or objects that we are interested in.
- ▶ **Sample:** A sample is a subset of the population. It is the group of individuals or objects that we actually collect data from.
- ▶ **Random Sample:** A random sample is a sample in which each individual or object in the population has an equal chance of being selected.

Sample Mean — Concepts

- ▶ **Population:** In statistics, a population is the entire pool from which a statistical sample is drawn. It is the complete set of individuals or objects that we are interested in.
- ▶ **Sample:** A sample is a subset of the population. It is the group of individuals or objects that we actually collect data from.
- ▶ **Random Sample:** A random sample is a sample in which each individual or object in the population has an equal chance of being selected.
- ▶ An alternative expression is
 X_1, X_2, \dots, X_n are independent and identically distributed (i.i.d.) random variables with mean μ and variance σ^2 .

Sample Mean — Justification

- By linearity of expectation, we have

$$E(T) = E(X_1 + X_2 + \cdots + X_n) = E(X_1) + E(X_2) + \cdots + E(X_n) = n\mu.$$

Sample Mean — Justification

- By linearity of expectation, we have

$$E(T) = E(X_1 + X_2 + \cdots + X_n) = E(X_1) + E(X_2) + \cdots + E(X_n) = n\mu.$$

- By independence, we have

$$\text{Var}(T) = \text{Var}(X_1 + X_2 + \cdots + X_n) = \text{Var}(X_1) + \text{Var}(X_2) + \cdots + \text{Var}(X_n) = n\sigma^2$$

Sample Mean — Justification

- By linearity of expectation, we have

$$E(T) = E(X_1 + X_2 + \cdots + X_n) = E(X_1) + E(X_2) + \cdots + E(X_n) = n\mu.$$

- By independence, we have

$$\text{Var}(T) = \text{Var}(X_1 + X_2 + \cdots + X_n) = \text{Var}(X_1) + \text{Var}(X_2) + \cdots + \text{Var}(X_n) = n\sigma^2$$

- Since $\bar{X} = T/n$, we have

$$E(\bar{X}) = E(T/n) = E(T)/n = \mu$$

and

$$\text{Var}(\bar{X}) = \text{Var}(T/n) = \text{Var}(T)/n^2 = \sigma^2/n.$$

Example: Bernoulli and Binomial

Suppose we have an unfair coin whose probability of landing heads is p . We toss the coin n times and let X_i be the indicator of the i -th toss.

- ▶ $X_i = 1$ if the i -th toss is head
- ▶ $X_i = 0$ if the i -th toss is tail

Example: Bernoulli and Binomial

Suppose we have an unfair coin whose probability of landing heads is p . We toss the coin n times and let X_i be the indicator of the i -th toss.

- ▶ $X_i = 1$ if the i -th toss is head
- ▶ $X_i = 0$ if the i -th toss is tail

X_i follows a **Bernoulli distribution** with $P(X_i = 1) = p$ and $P(X_i = 0) = 1 - p$.

- ▶ $E(X_i) = 1 \cdot P(X_i = 1) + 0 \cdot P(X_i = 0) = p$
- ▶ $Var(X_i) = E(X_i^2) - [E(X_i)]^2 = E(X_i) - [E(X_i)]^2 = p - p^2 = p(1 - p)$

Example: Bernoulli and Binomial

Let $T = X_1 + X_2 + \cdots + X_n$ be the number of heads from n tosses.

Example: Bernoulli and Binomial

Let $T = X_1 + X_2 + \cdots + X_n$ be the number of heads from n tosses.

By definition, T follows a **binomial distribution** with parameters n and p , denoted as $T \sim \text{Bin}(n, p)$. (number of successes in n independent Bernoulli trials)

Example: Bernoulli and Binomial

Let $T = X_1 + X_2 + \cdots + X_n$ be the number of heads from n tosses.

By definition, T follows a **binomial distribution** with parameters n and p , denoted as $T \sim \text{Bin}(n, p)$. (number of successes in n independent Bernoulli trials)

From our previous statement:

- ▶ $E(T) = n \cdot E(X_i) = np$
- ▶ $\text{Var}(T) = n \cdot \text{Var}(X_i) = np(1 - p)$

Example: Bernoulli and Binomial

Let $T = X_1 + X_2 + \cdots + X_n$ be the number of heads from n tosses.

By definition, T follows a **binomial distribution** with parameters n and p , denoted as $T \sim \text{Bin}(n, p)$. (number of successes in n independent Bernoulli trials)

From our previous statement:

- ▶ $E(T) = n \cdot E(X_i) = np$
- ▶ $\text{Var}(T) = n \cdot \text{Var}(X_i) = np(1 - p)$

Similarly, let $\bar{X} = T/n$ be the proportion of heads from n tosses. Then

- ▶ $E(\bar{X}) = E(X_i) = p$
- ▶ $\text{Var}(\bar{X}) = \text{Var}(X_i)/n = p(1 - p)/n$

Normal Population Distribution

Proposition

*Let X_1, X_2, \dots, X_n be a random sample from a **normal** distribution with mean μ and standard deviation σ . Then for any n , \bar{X} is normally distributed with mean μ and variance σ^2/n .*

Normal Population Distribution

Proposition

*Let X_1, X_2, \dots, X_n be a random sample from a **normal** distribution with mean μ and standard deviation σ . Then for any n , \bar{X} is normally distributed with mean μ and variance σ^2/n .*

A random variable X is said to have a normal distribution with mean μ and variance σ^2 , denoted by $N(\mu, \sigma^2)$, if its probability density function is given by

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/(2\sigma^2)}.$$

Normal Population Distribution

Proposition

Let X_1, X_2, \dots, X_n be a random sample from a **normal** distribution with mean μ and standard deviation σ . Then for any n , \bar{X} is normally distributed with mean μ and variance σ^2/n .

A random variable X is said to have a normal distribution with mean μ and variance σ^2 , denoted by $N(\mu, \sigma^2)$, if its probability density function is given by

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2/(2\sigma^2)}.$$

If $X_1 \sim N(\mu_1, \sigma_1^2)$ and $X_2 \sim N(\mu_2, \sigma_2^2)$ are independent, then

$$c_1 X_1 + c_2 X_2 \sim N(c_1 \mu_1 + c_2 \mu_2, c_1^2 \sigma_1^2 + c_2^2 \sigma_2^2).$$

Example

The distribution of egg weights of certian type is normal with mean value 53 and standard deviation 0.3.

Example

The distribution of egg weights of certain type is normal with mean value 53 and standard deviation 0.3.

Let X_1, X_2, \dots, X_{12} be the weights of a dozen randomly selected eggs.

Let $T = X_1 + X_2 + \dots + X_{12}$ be the total weight of the dozen eggs.

Example

The distribution of egg weights of certain type is normal with mean value 53 and standard deviation 0.3.

Let X_1, X_2, \dots, X_{12} be the weights of a dozen randomly selected eggs.

Let $T = X_1 + X_2 + \dots + X_{12}$ be the total weight of the dozen eggs.

$$E(T) = 12 \times 53 = 636, \quad Var(T) = 12 \times 0.3^2 = 1.08.$$

Example

The distribution of egg weights of certain type is normal with mean value 53 and standard deviation 0.3.

Let X_1, X_2, \dots, X_{12} be the weights of a dozen randomly selected eggs.

Let $T = X_1 + X_2 + \dots + X_{12}$ be the total weight of the dozen eggs.

$$E(T) = 12 \times 53 = 636, \quad Var(T) = 12 \times 0.3^2 = 1.08.$$

The probability that the total weight of the dozen eggs is between 635 and 640 is

$$P(635 < T < 640) = P\left(\frac{635 - 636}{\sqrt{1.08}} < Z < \frac{640 - 636}{\sqrt{1.08}}\right) = P(-0.96 < Z < 3.85) = 0.8315,$$

where $Z \sim N(0, 1)$ follows the **standard normal distribution**.

Central Limit Theorem

Theorem (Central Limit Theorem (CLT))

Let X_1, X_2, \dots, X_n be a random sample from a population with mean μ and variance σ^2 . Then if n is sufficiently large, \bar{X} has approximately a normal distribution with mean μ and variance σ^2/n , and T also has approximately a normal distribution with mean $n\mu$ and variance $n\sigma^2$. The larger the value of n , the better the approximation.

Central Limit Theorem

Theorem (Central Limit Theorem (CLT))

Let X_1, X_2, \dots, X_n be a random sample from a population with mean μ and variance σ^2 . Then if n is sufficiently large, \bar{X} has approximately a normal distribution with mean μ and variance σ^2/n , and T also has approximately a normal distribution with mean $n\mu$ and variance $n\sigma^2$. The larger the value of n , the better the approximation.

A shorter version:

$$\sqrt{n}(\bar{X} - \mu) \xrightarrow{\mathcal{D}} N(0, 1) \quad \text{as } n \rightarrow \infty$$

Central Limit Theorem

- ▶ **Implicit Assumption:** The population distribution has finite mean and finite variance.

Central Limit Theorem

- ▶ **Implicit Assumption:** The population distribution has finite mean and finite variance.
- ▶ **Sequential Interpretation:** The CLT applies to a sequence of i.i.d. random variables.

Central Limit Theorem

- ▶ **Implicit Assumption:** The population distribution has finite mean and finite variance.
- ▶ **Sequential Interpretation:** The CLT applies to a sequence of i.i.d. random variables.
- ▶ **Reparametrization:** When n is large enough, $Y = \sqrt{n}(\bar{X} - \mu)$ is approximately standard normal, where the limit distribution does not depend on n .

Central Limit Theorem

- ▶ **Implicit Assumption:** The population distribution has finite mean and finite variance.
- ▶ **Sequential Interpretation:** The CLT applies to a sequence of i.i.d. random variables.
- ▶ **Reparametrization:** When n is large enough, $Y = \sqrt{n}(\bar{X} - \mu)$ is approximately standard normal, where the limit distribution does not depend on n .
- ▶ **Approximation:** The approximated distribution should be interpreted that the c.d.f. of Y , $P(Y \leq t)$, converges to the c.d.f. of $N(0, 1)$ as $n \rightarrow \infty$ for any t .

Central Limit Theorem

- ▶ **Implicit Assumption:** The population distribution has finite mean and finite variance.
- ▶ **Sequential Interpretation:** The CLT applies to a sequence of i.i.d. random variables.
- ▶ **Reparametrization:** When n is large enough, $Y = \sqrt{n}(\bar{X} - \mu)$ is approximately standard normal, where the limit distribution does not depend on n .
- ▶ **Approximation:** The approximated distribution should be interpreted that the c.d.f. of Y , $P(Y \leq t)$, converges to the c.d.f. of $N(0, 1)$ as $n \rightarrow \infty$ for any t .
- ▶ **Proof:** The proof of the Central Limit Theorem is beyond the scope of this course. It is a result from the characteristic function and the Lévy's convergence theorem.

Central Limit Theorem

- ▶ **Implicit Assumption:** The population distribution has finite mean and finite variance.
- ▶ **Sequential Interpretation:** The CLT applies to a sequence of i.i.d. random variables.
- ▶ **Reparametrization:** When n is large enough, $Y = \sqrt{n}(\bar{X} - \mu)$ is approximately standard normal, where the limit distribution does not depend on n .
- ▶ **Approximation:** The approximated distribution should be interpreted that the c.d.f. of Y , $P(Y \leq t)$, converges to the c.d.f. of $N(0, 1)$ as $n \rightarrow \infty$ for any t .
- ▶ **Proof:** The proof of the Central Limit Theorem is beyond the scope of this course. It is a result from the characteristic function and the Lévy's convergence theorem.
- ▶ **Rule of Thumb:** $n \geq 30$ is often considered as a sufficiently large sample size.

Example

Let Y be a Binomial random variable with parameters $n = 100$ and $p = 0.5$.
We want to estimate the probability $P(40 < Y < 60)$.

Example

Let Y be a Binomial random variable with parameters $n = 100$ and $p = 0.5$. We want to estimate the probability $P(40 < Y < 60)$.

Recall our discussion on tossing a coin. Let X_i be the indicator of the i -th toss. Then $T = X_1 + X_2 + \cdots + X_{100}$ follows a Binomial distribution with parameters $n = 100$ and $p = 0.5$. That is, $T \sim Y$.

Example

Let Y be a Binomial random variable with parameters $n = 100$ and $p = 0.5$. We want to estimate the probability $P(40 < Y < 60)$.

Recall our discussion on tossing a coin. Let X_i be the indicator of the i -th toss. Then $T = X_1 + X_2 + \cdots + X_{100}$ follows a Binomial distribution with parameters $n = 100$ and $p = 0.5$. That is, $T \sim Y$. From the central limit theorem,

$$\bar{X} \approx N(p, p(1-p)/n) \sim N(0.5, 0.0025).$$

Therefore, $T = n\bar{X} \sim N(50, 25)$.

Example

Let Y be a Binomial random variable with parameters $n = 100$ and $p = 0.5$. We want to estimate the probability $P(40 < Y < 60)$.

Recall our discussion on tossing a coin. Let X_i be the indicator of the i -th toss. Then $T = X_1 + X_2 + \cdots + X_{100}$ follows a Binomial distribution with parameters $n = 100$ and $p = 0.5$. That is, $T \sim Y$. From the central limit theorem,

$$\bar{X} \approx N(p, p(1-p)/n) \sim N(0.5, 0.0025).$$

Therefore, $T = n\bar{X} \sim N(50, 25)$.

We have

$$P(40 < Y < 60) = P(40 < T < 60) \approx P\left(\frac{40 - 50}{\sqrt{25}} < Z < \frac{60 - 50}{\sqrt{25}}\right) = P(-2 < Z < 2)$$

From Probabilistic Model to Statistical Inference

► **Probabilistic Model:**

Given the (known) parameters of the population (e.g. mean and variance), we can model the distribution of the sample.

From Probabilistic Model to Statistical Inference

- ▶ **Probabilistic Model:**

Given the (known) parameters of the population (e.g. mean and variance), we can model the distribution of the sample.

- ▶ **Statistical Inference:**

Given the sample, we want to estimate the (unknown) parameters of the population.

From Probabilistic Model to Statistical Inference

- **Probabilistic Model:**

Given the (known) parameters of the population (e.g. mean and variance), we can model the distribution of the sample.

- **Statistical Inference:**

Given the sample, we want to estimate the (unknown) parameters of the population.

