

STAT 576 Bayesian Analysis

Lecture 6: Model Checking

Chencheng Cai

Washington State University

Model Checking Methods

Goal:

- ▶ Assess the fit of the model to the data.
- ▶ Assess the fit of the model to our substantive knowledge.
- ▶ Assess the adequacy/robustness of the model.

Model Checking Methods

Goal:

- ▶ Assess the fit of the model to the data.
- ▶ Assess the fit of the model to our substantive knowledge.
- ▶ Assess the adequacy/robustness of the model.

Methods:

- ▶ Sensitivity Analysis.
 - ▶ Check whether other models generate a similar posterior.
- ▶ External Validation.
 - ▶ Posterior predictive checking.
- ▶ Internal Validation.
 - ▶ Cross-validation predictive checking.


Sensitivity Analysis

- ▶ How the results are affected by different choices of the model structure?
 - ▶ different models (binomial v.s. Poisson, normal v.s. t)
 - ▶ different priors
 - ▶ different structures (hierarchical v.s. separate)
 - ▶ different distribution families (Gaussian v.s. mixed Gaussian)

Sensitivity Analysis


- ▶ How the results are affected by different choices of the model structure?
 - ▶ different models (binomial v.s. Poisson, normal v.s. t)
 - ▶ different priors
 - ▶ different structures (hierarchical v.s. separate)
 - ▶ different distribution families (Gaussian v.s. mixed Gaussian)
- ▶ Compare the sensitivity of essential inference quantities.
 - ▶ extreme quantities v.s. mean/median.
 - ▶ extrapolation v.s. interpolation.

Example: Election Prediction




- ▶ Posterior winning probability of Bill Clinton at each state in Oct. 1992.
- ▶ Hierarchical linear regression model.

Example: Election Prediction



- ▶ Posterior winning probability of Bill Clinton at each state in Oct. 1992.
- ▶ Hierarchical linear regression model.
- ▶ The model seems wrong at Texas and Florida.

Example: Election Prediction



- ▶ Posterior winning probability of Bill Clinton at each state in Oct. 1992.
- ▶ Hierarchical linear regression model.
- ▶ The model seems wrong at Texas and Florida.
- ▶ It is much easier to evaluate the performance afterwards.

Posterior Predictive Checking

- ▶ Idea: check the discrepancy between the predicted values and the observed values.

Posterior Predictive Checking


- ▶ Idea: check the discrepancy between the predicted values and the observed values.
- ▶ Procedure:
 - ▶ Generate simulated samples from the **joint posterior predictive distribution**
 - ▶ Compare the samples with the observed data.
 - ▶ Systematic differences imply the failings of the model.

Example: Light Speed Experiment

- ▶ Simon Newcomb set up an experiment in 1882 to measure the light speed.
- ▶ The travel time of light was recorded for the round-trip between
 - ▶ his lab on the Potomac river
 - ▶ a mirror at the base of the Washington Monument
- ▶ The total travel distance is 7422 meters.

Example: Light Speed Experiment

- ▶ Simon Newcomb set up an experiment in 1882 to measure the light speed.
- ▶ The travel time of light was recorded for the round-trip between
 - ▶ his lab on the Potomac river
 - ▶ a mirror at the base of the Washington Monument
- ▶ The total travel distance is 7422 meters.
- ▶ The measurement was repeated $n = 66$ times.



Histogram for deviations from 24800 ns

Example: Light Speed Experiment

- We model the travel time by a normal distribution:

$$y_i \sim \mathcal{N}(\mu, \sigma^2)$$

Example: Light Speed Experiment

- ▶ We model the travel time by a normal distribution:

$$y_i \sim \mathcal{N}(\mu, \sigma^2)$$

- ▶ We can choose a noninformative prior for μ and σ^2 :

$$p(\mu, \sigma^2) \propto \frac{1}{\sigma^2}$$

Example: Light Speed Experiment

- ▶ We model the travel time by a normal distribution:

$$y_i \sim \mathcal{N}(\mu, \sigma^2)$$

- ▶ We can choose a noninformative prior for μ and σ^2 :

$$p(\mu, \sigma^2) \propto \frac{1}{\sigma^2}$$

- ▶ Recall our previous results for multiparameter Bayesian inference. The marginal posterior for μ is

$$\mu \mid y \sim t_{66} \left(\bar{y}, \frac{65}{66^2} s^2 \right)$$

Example: Light Speed Experiment

- ▶ We model the travel time by a normal distribution:

$$y_i \sim \mathcal{N}(\mu, \sigma^2)$$

- ▶ We can choose a noninformative prior for μ and σ^2 :

$$p(\mu, \sigma^2) \propto \frac{1}{\sigma^2}$$

- ▶ Recall our previous results for multiparameter Bayesian inference. The marginal posterior for μ is

$$\mu \mid y \sim t_{66} \left(\bar{y}, \frac{65}{66^2} s^2 \right)$$

- ▶ A 95% credible interval is $[23.6, 28.8]$.
- ▶ We know the true value should be around 33.0.

Example: Light Speed Experiment


Generate posterior predictive replicates y^{rep}

- ▶ Draw $\mu^{(s)}, \sigma^{2(s)}$ from the joint posterior distribution $p(\mu, \sigma^2 \mid y)$.
- ▶ Draw $y^{rep(s)}$ from $\mathcal{N}(\mu^{(s)}, \sigma^{2(s)})$.
- ▶ Repeat the drawing to get n replicates of y^{rep} .

Example: Light Speed Experiment


Generate posterior predictive replicates y^{rep}

- ▶ Draw $\mu^{(s)}, \sigma^{2(s)}$ from the joint posterior distribution $p(\mu, \sigma^2 | y)$.
- ▶ Draw $y^{rep(s)}$ from $\mathcal{N}(\mu^{(s)}, \sigma^{2(s)})$.
- ▶ Repeat the drawing to get n replicates of y^{rep} .




Example: Light Speed Experiment

We get the histogram of the **smallest** travel time for all replicates.



Example: Light Speed Experiment


We get the histogram of the **smallest** travel time for all replicates.



- Can hardly observe an occurrence that is less than -20.

Example: Light Speed Experiment


We get the histogram of the **smallest** travel time for all replicates.



- ▶ Can hardly observe an occurrence that is less than -20.
- ▶ Decide: whether the **data** was wrong or the **model** was wrong?

Example: Light Speed Experiment

We get the histogram of the **smallest** travel time for all replicates.



- ▶ Can hardly observe an occurrence that is less than -20.
- ▶ Decide: whether the **data** was wrong or the **model** was wrong?
- ▶ The model was wrong: should use heavy-tailed distribution or contaminated normal (mixed Gaussian).

Posterior Predictive Checking

- ▶ Replicated datasets:

$$p(y^{rep} | y) = \int \underbrace{p(y^{rep} | \theta)}_{\text{obs. model}} \underbrace{p(\theta | y)}_{\text{posterior}} d\mu(\theta)$$

Posterior Predictive Checking

- ▶ Replicated datasets:

$$p(y^{rep} | y) = \int \underbrace{p(y^{rep} | \theta)}_{\text{obs. model}} \underbrace{p(\theta | y)}_{\text{posterior}} d\mu(\theta)$$

- ▶ **Test quantity** (or discrepancy measure) $T(y, \theta)$
 - ▶ Summary quantity for the observed data $T(y, \theta)$
 - ▶ Summary quantity for a replicated data $T(y^{rep}, \theta)$.

Posterior Predictive Checking

- ▶ Replicated datasets:

$$p(y^{rep} | y) = \int \underbrace{p(y^{rep} | \theta)}_{obs. \ model} \underbrace{p(\theta | y)}_{posterior} d\mu(\theta)$$

- ▶ **Test quantity** (or discrepancy measure) $T(y, \theta)$
 - ▶ Summary quantity for the observed data $T(y, \theta)$
 - ▶ Summary quantity for a replicated data $T(y^{rep}, \theta)$.
- ▶ The frequentist counter-part is known as **test statistics** $T(y)$, which only depends on the data.

Posterior Predictive Checking

- ▶ Replicated datasets:

$$p(y^{rep} | y) = \int \underbrace{p(y^{rep} | \theta)}_{obs. \ model} \underbrace{p(\theta | y)}_{posterior} d\mu(\theta)$$

- ▶ **Test quantity** (or discrepancy measure) $T(y, \theta)$
 - ▶ Summary quantity for the observed data $T(y, \theta)$
 - ▶ Summary quantity for a replicated data $T(y^{rep}, \theta)$.
- ▶ The frequentist counter-part is known as **test statistics** $T(y)$, which only depends on the data.
- ▶ In the light speed example, we choose $T(y, \theta) = \min(y)$ (also a test statistic).

Posterior Predictive Checking

Classical p-values:

$$p_C = \mathbb{P}[T(y^{rep}) \geq T(y) \mid \theta]$$

Posterior Predictive Checking

Classical p-values:

$$p_C = \mathbb{P}[T(y^{rep}) \geq T(y) \mid \theta]$$

Posterior predictive p-values:

$$p_B = \mathbb{P}[T(y^{rep}, \theta) \geq T(y, \theta) \mid y]$$

Posterior Predictive Checking

Classical p-values:

$$p_C = \mathbb{P}[T(y^{rep}) \geq T(y) \mid \theta]$$

Posterior predictive p-values:

$$p_B = \mathbb{P}[T(y^{rep}, \theta) \geq T(y, \theta) \mid y]$$

- ▶ The classical p-values measure how likely the data is coming from the null model.
- ▶ The posterior predictive p-values measure how likely the data is similar to the posterior predictive replicates.

Posterior Predictive Checking

Classical p-values:

$$p_C = \mathbb{P}[T(y^{rep}) \geq T(y) \mid \theta]$$

Posterior predictive p-values:


$$p_B = \mathbb{P}[T(y^{rep}, \theta) \geq T(y, \theta) \mid y]$$

- ▶ The classical p-values measure how likely the data is coming from the null model.
- ▶ The posterior predictive p-values measure how likely the data is similar to the posterior predictive replicates.
- ▶ In Bayesian, θ is also random. p_B can be estimated by joint samples of (y^{rep}, θ) .

$$\begin{aligned} p_B &= \iint \mathbb{I}\{T(y^{rep}, \theta) \geq T(y, \theta)\} p(y^{rep} \mid \theta) p(\theta \mid y) d\mu(\theta) d\mu(y^{rep}) \\ &\approx \frac{1}{S} \sum_{s=1}^S \mathbb{I}\{T(y^{rep(s)}, \theta^{(s)}) \geq T(y, \theta^{(s)})\} \end{aligned}$$

Example: Light Speed Experiment

If we use the sample variance as the test quantity:



Cannot tell the discrepancy — because the sample variance is a sufficient statistics.

Posterior Predictive Checking

- ▶ A **good** test statistic is ancilliary
 - ▶ ancilliary: depends on the observed data but independent of the parameters.
- ▶ A **bad** test statistic is highly dependent of the parameters.
 - ▶ i.e. sufficient statistics.

Posterior Predictive Checking

- ▶ A **good** test statistic is ancilliary
 - ▶ ancilliary: depends on the observed data but independent of the parameters.
- ▶ A **bad** test statistic is highly dependent of the parameters.
 - ▶ i.e. sufficient statistics.
- ▶ If we have multiple test statistics, we do not conduct p-value justification.
 - ▶ See the smoking example in the textbook.

Posterior Predictive Checking

- ▶ A **good** test statistic is ancilliary
 - ▶ ancilliary: depends on the observed data but independent of the parameters.
- ▶ A **bad** test statistic is highly dependent of the parameters.
 - ▶ i.e. sufficient statistics.
- ▶ If we have multiple test statistics, we do not conduct p-value justification.
 - ▶ See the smoking example in the textbook.
- ▶ An extreme p-value often suggests the weakness of the current model. The next step is to revise the model.

Example: Educational Testing

Data: the effects of coaching programs for the SAT-V scores for students in 8 schools.

| School | Estimated treatment effect, y_j | Standard error of effect estimate, σ_j |
|--------|---|---|
| A | 28 | 15 |
| B | 8 | 10 |
| C | -3 | 16 |
| D | 7 | 11 |
| E | -1 | 9 |
| F | 1 | 11 |
| G | 18 | 10 |
| H | 12 | 18 |

Example: Educational Testing

Separate estimation:

- ▶ Some schools have moderate effects (18-28).
- ▶ Most schools have small effects (0-12).
- ▶ Two have negative effects.
- ▶ Difficult to distinguish because of large variance.

Example: Educational Testing

Separate estimation:

- ▶ Some schools have moderate effects (18-28).
- ▶ Most schools have small effects (0-12).
- ▶ Two have negative effects.
- ▶ Difficult to distinguish because of large variance.

Pooled estimation:

- ▶ All schools have identical effect θ .
- ▶ Use noninformative prior.
- ▶ Posterior mean: 7.7 with s.e. 4.1

Example: Educational Testing

Separate estimation:

- ▶ Some schools have moderate effects (18-28).
- ▶ Most schools have small effects (0-12).
- ▶ Two have negative effects.
- ▶ Difficult to distinguish because of large variance.

Pooled estimation:

- ▶ All schools have identical effect θ .
- ▶ Use noninformative prior.
- ▶ Posterior mean: 7.7 with s.e. 4.1

Hierarchical model:

- ▶ $\theta_1, \dots, \theta_8 \sim \mathcal{N}(\mu, \tau^2)$ i.i.d.
- ▶ $y_j \mid \theta_j \sim (\theta_j, \sigma_j^2)$ independent.
- ▶ choose flat prior $p(\mu, \tau) \propto 1$.

Example: Educational Testing

Hierarchical model:

- ▶ By drawing posterior samples:
 - ▶ draw $\mu^{(s)}, \tau^{(s)}$ from $p(\mu, \tau \mid y)$
 - ▶ draw $\theta_1^{(s)}, \dots, \theta_8^{(s)}$ from $p(\theta_1, \dots, \theta_8 \mid \mu^{(s)}, \tau^{(s)}, y)$

Example: Educational Testing

Hierarchical model:

- ▶ By drawing posterior samples:
 - ▶ draw $\mu^{(s)}, \tau^{(s)}$ from $p(\mu, \tau \mid y)$
 - ▶ draw $\theta_1^{(s)}, \dots, \theta_8^{(s)}$ from $p(\theta_1, \dots, \theta_8 \mid \mu^{(s)}, \tau^{(s)}, y)$
- ▶ we have the posterior quantiles for each school:

| School | Posterior quantiles | | | | |
|--------|---------------------|-----|--------|-----|-------|
| | 2.5% | 25% | median | 75% | 97.5% |
| A | -2 | 7 | 10 | 16 | 31 |
| B | -5 | 3 | 8 | 12 | 23 |
| C | -11 | 2 | 7 | 11 | 19 |
| D | -7 | 4 | 8 | 11 | 21 |
| E | -9 | 1 | 5 | 10 | 18 |
| F | -7 | 2 | 6 | 10 | 28 |
| G | -1 | 7 | 10 | 15 | 26 |
| H | -6 | 3 | 8 | 13 | 33 |

Example: Educational Testing — Model Checking





- ▶ Assumptions:
 - ▶ normality of y_j .
 - ▶ exchangeability of the priors for θ_j 's.
 - ▶ normality of prior of θ_j .
 - ▶ flat hyperprior.

Example: Educational Testing — Model Checking

- ▶ Assumptions:
 - ▶ normality of y_j .
 - ▶ exchangeability of the priors for θ_j 's.
 - ▶ normality of prior of θ_j .
 - ▶ flat hyperprior.
- ▶ Comparing posterior inferences to substantive knowledge:
 - ▶ Individual effects between 5 and 10 seems reasonable.
 - ▶ Some lower bounds go to negative.

Example: Educational Testing — Model Checking


- ▶ Posterior predictive checking.
 - ▶ $y^{rep} = (y_1^{rep}, \dots, y_8^{rep})$
 - ▶ Test statistics: max, min, mean, s.d.



Example: Educational Testing — Model Checking

Sensitivity Analysis:


- ▶ Uniform prior for τ : the marginal posterior for τ
— no significant change if we multiply it by another prior



Example: Educational Testing — Model Checking

Sensitivity Analysis:

- ▶ Uniform prior for τ : the marginal posterior for τ
— no significant change if we multiply it by another prior




- ▶ normality of $y_j \mid \theta_j, \sigma_j$: ensured by experimental design and CLT.

Example: Educational Testing — Model Checking

Sensitivity Analysis:

- ▶ Uniform prior for τ : the marginal posterior for τ
— no significant change if we multiply it by another prior



- ▶ normality of $y_j \mid \theta_j, \sigma_j$: ensured by experimental design and CLT.
- ▶ normality of the prior for θ_j 's:
One may consider other heavy-tailed distributions. But needs advanced sampling techniques.