

Set 12 - Confidence intervals

STAT 401 (Engineering) - Iowa State University

February 19, 2017

Confidence intervals

Definition

The **coverage** of an interval estimator is the probability the interval will contain the true value of the parameter *when the data are considered to be random*. If an interval estimator has $100(1 - \alpha)\%$ coverage, then we call it a $100(1 - \alpha)\%$ **confidence interval** and $1 - \alpha$ is the **confidence level**.

Normal model

If $Y_i \stackrel{ind}{\sim} N(\mu, \sigma^2)$ and we assume the default prior $p(\mu, \sigma^2) \propto 1/\sigma^2$, then a $100(1 - a)\%$ credible interval for μ is given by

$$\bar{y} \pm t_{n-1, a/2} s / \sqrt{n}.$$

When the data are considered random

$$T = \frac{\bar{Y} - \mu}{S/\sqrt{n}} \sim t_{n-1} \sim t_{n-1}(0, 1)$$

thus the probability μ is within our credible interval is

$$\begin{aligned} & P\left(\bar{Y} - t_{n-1, a/2} S / \sqrt{n} < \mu < \bar{Y} + t_{n-1, a/2} S / \sqrt{n}\right) \\ &= P\left(-t_{n-1, a/2} < \frac{\bar{Y} - \mu}{S/\sqrt{n}} < t_{n-1, a/2}\right) \\ &= P\left(-t_{n-1, a/2} < T < t_{n-1, a/2}\right) \\ &= 1 - a. \end{aligned}$$

Thus, this $100(1 - a)\%$ credible interval is also a $100(1 - a)\%$ confidence interval.

Data example

Recall the corn yield example from Set 11 with 9 randomly selected fields in Iowa whose sample average yield is 205 and sample standard deviation is 21. Then a 95% confidence interval for the is

$$205 \pm 2.31 \times 21/\sqrt{9} = (221, 189).$$

This confidence interval tells us nothing about the true mean yield of fields in Iowa. Instead, it tells us that if we use this **procedure** repeatedly on different data sets, then 95% of the time, the interval will contain the true parameter.

Sampling distribution

Definition

The **sampling distribution** of a statistic is the distribution of the statistic *when the data are considered random*.

Example

If $Y_i \stackrel{ind}{\sim} N(\mu, \sigma^2)$, then

$$T = \frac{\bar{Y} - \mu}{S/\sqrt{n}} \sim t_{n-1}(0, 1)$$

where \bar{Y} and S are the sample average and sample standard deviation over n observations.

Most of the time we don't know the sampling distribution of the statistic.

Approximate sampling distributions

If the estimator $\hat{\theta} = \hat{\theta}(Y)$ is based on an average or a sum, then the Central Limit Theorem tells us what its **approximate** sampling distribution is, i.e.

$$\hat{\theta} \sim N(E[\hat{\theta}], Var[\hat{\theta}]).$$

Example

If $Y_i \stackrel{ind}{\sim} N(\mu, \sigma^2)$, then

$$\hat{\theta} = \bar{Y} \sim N(E[\bar{Y}], Var[\bar{Y}])$$

where

$$\begin{aligned} E[\bar{Y}] &= \mu \\ Var[\bar{Y}] &= \sigma^2/n. \end{aligned}$$

Standard error

Definition

The **standard error** of an estimator is an *estimate* of the standard deviation of the sampling distribution of the estimator.

Example

If $Y_i \stackrel{\text{ind}}{\sim} N(\mu, \sigma^2)$ and $\hat{\theta} = \bar{Y}$, then

$$\text{Var}[\hat{\theta}] = \text{Var}[\bar{Y}] = \sigma^2/n \quad \text{and} \quad \sqrt{\text{Var}[\hat{\theta}]} = \sigma/\sqrt{n}.$$

We have the following *consistent* estimators

$$S^2 \xrightarrow{P} \sigma^2, \quad S \xrightarrow{P} \sigma, \quad \text{and} \quad S/\sqrt{n} \xrightarrow{P} \sigma/\sqrt{n}$$

and thus

$$SE[\hat{\theta}] = \sqrt{\widehat{\text{Var}[\hat{\theta}]}} = S/\sqrt{n}.$$

where S is the sample standard deviation.

Approximate confidence intervals

If an estimator has an asymptotic normal distribution, then we can construct an **approximate** $100(1 - \alpha)\%$ confidence interval for $E[\hat{\theta}]$ using

$$\hat{\theta} \pm z_{\alpha/2} SE[\hat{\theta}].$$

where $SE(\hat{\theta}) = \sqrt{Var[\hat{\theta}]}$ is the **standard error** of the estimator.

This comes from the fact that if $\hat{\theta} \sim N(E[\hat{\theta}], Var[\hat{\theta}])$, then

$$\begin{aligned} &P\left(\hat{\theta} - z_{\alpha/2} SE(\hat{\theta}) < E[\hat{\theta}] < \hat{\theta} + z_{\alpha/2} SE(\hat{\theta})\right) \\ &= P\left(-z_{\alpha/2} < \frac{\hat{\theta} - E[\hat{\theta}]}{SE(\hat{\theta})} < z_{\alpha/2}\right) \\ &= P\left(-z_{\alpha/2} < \frac{\hat{\theta} - E[\hat{\theta}]}{\sqrt{Var[\hat{\theta}]}} < z_{\alpha/2}\right) \\ &\approx P\left(-z_{\alpha/2} < Z < z_{\alpha/2}\right) \\ &= 1 - \alpha. \end{aligned}$$

Normal example

If $Y_i \stackrel{\text{ind}}{\sim} N(\mu, \sigma^2)$ and we have the estimator $\hat{\theta} = \bar{Y}$, then

$$\begin{aligned} E[\hat{\theta}] &= \mu \\ SE[\hat{\theta}] &= S/\sqrt{n} \end{aligned}$$

Thus an **approximate** $100(1 - a)\%$ confidence interval for $\mu = E[\hat{\theta}]$ is


$$\bar{Y} \pm z_{a/2} SE[\hat{\theta}] = \bar{Y} \pm z_{a/2} S/\sqrt{n}.$$

Note that this is almost identical to the **exact** $100(1 - a)\%$ confidence interval for μ ,

$$\bar{Y} \pm t_{n-1, a/2} S/\sqrt{n}$$

and when n is large $z_{a/2} \approx t_{n-1, a/2}$.

T critical values vs Z critical values



Binomial example

Suppose $Y \sim \text{Bin}(n, \theta)$ and we are interested in a confidence interval for θ . An estimator for θ is

$$\hat{\theta} = \frac{Y}{n}$$

since $E[\hat{\theta}] = \theta$. The variance when the data are considered random of this estimator is

$$\text{Var}[\hat{\theta}] = \frac{\theta(1 - \theta)}{n}.$$

A standard error for this estimator is

$$SE[\hat{\theta}] = \sqrt{\frac{\hat{\theta}(1 - \hat{\theta})}{n}}$$

since $\hat{\theta} = Y/n$ is a consistent estimator for θ

Approximate confidence interval for binomial proportion

If $Y \sim \text{Bin}(n, \theta)$, then an approximate $100(1 - \alpha)\%$ confidence interval for θ is

$$\hat{\theta} \pm z_{\alpha/2} \sqrt{\frac{\hat{\theta}(1 - \hat{\theta})}{n}}.$$

where $\hat{\theta} = Y/n$.

Example

The Gallup poll is a poll of approximately 1,500 (randomly selected?) U.S. adults. In the post recent poll (2017/02/19), 32.1% of respondents indicated that they were engaged at work. Thus an approximate 95% confidence interval for the proportion of all U.S. adults is

$$0.321 \pm 1.96 \times \sqrt{\frac{.321(1 - .321)}{1500}} = (0.30, 0.34).$$

But the confidence interval actually says nothing about where the true proportion is. Instead it says something about how many of our intervals would cover the truth if we were to repeat this procedure over and over.

Confidence interval summary

Model	Estimator	Confidence Interval	Type
$Y_i \stackrel{ind}{\sim} N(\mu, \sigma^2)$	$\hat{\mu} = \bar{y}$	$\hat{\mu} \pm t_{n-1, a/2} s / \sqrt{n}$	exact
$Y_i \stackrel{ind}{\sim} N(\mu, \sigma^2)$	$\hat{\mu} = \bar{y}$	$\hat{\mu} \pm z_{a/2} s / \sqrt{n}$	approximate
$Y_i \stackrel{ind}{\sim} Ber(\theta)$	$\hat{\theta} = \bar{y}$	$\hat{\theta} \pm z_{a/2} \sqrt{\hat{\theta}(1 - \hat{\theta})/n}$	approximate
$Y \sim Bin(n, \theta)$	$\hat{\theta} = y/n$	$\hat{\theta} \pm z_{a/2} \sqrt{\hat{\theta}(1 - \hat{\theta})/n}$	approximate

Bayesian credible intervals based on the priors we have discussed are generally approximate confidence intervals