

Midterm review

Dr. Jarad Niemi

Iowa State University

March 1, 2017

What we have covered

Chapters

- Probability and inference (Ch 1)
- Single-parameter models (Ch 2)
- Introduction to multiparameter models (Ch 3)
- Asymptotics and connections to non-Bayesian approaches (Ch 4)
- Hierarchical models (Ch 5)
- Model checking (Ch 6)
- Bayesian hypothesis tests (Sec 7.4)
- Decision theory (Sec 9.1)
- Stan

Probability and inference (Ch 1)

- Three steps of Bayesian data analysis (Sec 1.1)
 - Set up a full probability model: $p(y|\theta)$ and $p(\theta)$
 - Condition on observed data: $p(\theta|y)$
 - Evaluate the fit of the model: $p(y^{rep}|y)$
- Bayesian inference via Bayes' rule (Sec 1.3)
 - Parameter posteriors: $p(\theta|y) \propto p(y|\theta)p(\theta)$
 - Predictions: $p(\tilde{y}|y) = \int p(\tilde{y}|\theta)p(\theta|y)d\theta$
 - Model probabilities $p(M|y) \propto p(y|M)p(M)$ where $p(y|M) = \int p(y|\theta, M)p(\theta|M)d\theta$.
- Interpreting Bayesian probabilities (Sec 1.5)
 - Epistemic probability: my belief
 - Frequency probability: long run percentage
- Computation (Sec 1.9)
 - Inference via simulations

Single-parameter models (Ch 2)

General

- Priors
 - Conjugate (Sec 2.4)
 - Default - Jeffreys (Sec 2.8)
 - Weakly informative (Sec 2.9)
- Posteriors
 - Compromise between data and prior (2.2)
 - Point estimation
 - Credible intervals (Sec 2.3)

Specific models

- Binomial (Sec 2.1–2.4)
- Normal, unknown mean (Sec 2.5)
- Normal, unknown variance (Sec 2.6)
- Poisson (Sec 2.6)
- Exponential (Sec 2.6)
- Poisson with exposure (Sec 2.7)

Single-parameter models (Ch 2)

Additional comments:

- Deriving posteriors using the **kernel**
- Discrete priors are conjugate
- Mixtures of conjugate priors are conjugate
- Point estimation depends on utility function
 - Mean minimizes squared error
 - Median minimizes absolute error
 - Mode is obtained as a limit of minimizing a sequence of 0-1 errors
- Credible intervals
 - One-tailed
 - Equal-tailed
 - Highest posterior density

Introduction to multiparameter models (Ch 3)

- Joint posterior

$$p(\theta_1, \dots, \theta_n | y) \propto p(y | \theta_1, \dots, \theta_n) p(\theta_1, \dots, \theta_n)$$

- Marginal posterior

$$p(\theta_1 | y) = \int \cdots \int p(\theta_1, \dots, \theta_n | y) d\theta_2 \cdots d\theta_n$$

- Conditional posteriors

$$p(\theta_2, \dots, \theta_n | \theta_1, y) \propto p(\theta_1, \dots, \theta_n | y)$$

- Posterior decomposition, e.g.

$$p(\theta_1, \dots, \theta_n | y) = p(\theta_1 | y) \prod_{i=2}^n p(\theta_i | \theta_{1:i-1}, y)$$

where $1 : i - 1 = 1, 2, \dots, i - 1$.

- Conditional independence, e.g.

$$p(\theta_i | \theta_{1:i-1}, y) = p(\theta_i | \theta_{i-1}, y)$$

Normal model

- Normal model with default prior (Sec 3.2)

$$y_i \stackrel{iid}{\sim} N(\mu, \sigma^2) \quad p(\mu, \sigma^2) \propto 1/\sigma^2$$

results in

$$p(\mu, \sigma^2 | y) = N(\bar{y}, \sigma^2/n) \text{Inv-}\chi^2(n-1, s^2)$$

where $s^2 = \frac{1}{n-1} \sum_{i=1}^n (y_i - \bar{y})^2$.

- Normal model with conjugate prior (Sec 3.3)

$$y \stackrel{iid}{\sim} N(\mu, \sigma^2) \quad \mu | \sigma^2 \sim N(\mu_0, \sigma^2/\kappa_0) \quad \sigma^2 \sim \text{Inv-}\chi^2(\nu_0, \sigma_0^2)$$

results in

$$p(\mu, \sigma^2 | y) = N\left(\frac{\kappa_0 \mu_0 + n \bar{y}}{\kappa_0 + n}, \frac{\sigma^2}{\kappa_0 + n}\right) \text{Inv-}\chi^2(\nu_0 + n, \sigma_n^2)$$

where $\sigma_n^2 = \left[\nu_0 \sigma_0^2 + (n-1)s^2 + \frac{\kappa_0 n}{\kappa_0 + n} (\bar{y} - \mu_0)^2 \right] / (\nu_0 + n)$.

Data asymptotics (Ch 4)

Consider a model $y_i \stackrel{iid}{\sim} p(y|\theta_0)$ for some true value θ_0 .

- Posterior convergence:

If A is a neighborhood of θ_0 , then $Pr(\theta \in A|y) \rightarrow 1$.

- Point estimation:

$$\hat{\theta}_{Bayes} \rightarrow \hat{\theta}_{MLE} \xrightarrow{P} \theta_0$$

- Limiting distribution:

$$\theta|y \xrightarrow{d} N\left(\hat{\theta}, \frac{1}{n}I(\hat{\theta})^{-1}\right)$$

Asymptotics - What can go wrong?

- Not unique to Bayesian statistics
 - Unidentified parameters
 - Number of parameters increase with sample size
 - Aliasing
 - Unbounded likelihoods
 - Tails of the distribution
 - True sampling distribution is not $p(y|\theta)$
- Unique to Bayesian statistics
 - Improper posterior
 - Prior distributions that exclude the point of convergence
 - Convergence to the edge of the parameter space

Hierarchical models (Ch 5)

- Hierarchical model (Ch 5):

$$p(\theta, \phi | y) \propto p(y | \theta) p(\theta | \phi) p(\phi)$$

- Exchangeability (Sec 5.2)

$$p(y_1, \dots, y_n) = p(y_{\pi_1}, \dots, y_{\pi_n})$$

- Hierarchical binomial model (Sec 5.3):

$$y_i \stackrel{iid}{\sim} \text{Bin}(n_i, \theta_i) \quad \theta_i \stackrel{iid}{\sim} \text{Be}(\alpha, \beta)$$

- Hierarchical Poisson (with exposure) model

$$y_i \stackrel{iid}{\sim} \text{Po}(x_i \lambda_i) \quad \lambda_i \stackrel{iid}{\sim} \text{Ga}(\mu\beta, \beta)$$

- Hierarchical normal model (Sec 5.4)

$$y_{ij} \stackrel{iid}{\sim} N(\mu_j, \sigma_j^2) \quad \mu_j \stackrel{iid}{\sim} N(\eta, \tau^2) \quad \sigma_j^2 \stackrel{iid}{\sim} \text{Ga}(\alpha, \beta)$$

Model checking (Ch 6)

- Data replications

$$p(y^{rep}|y) = \int p(y^{rep}|\theta)p(\theta|y)d\theta$$

- Graphical posterior predictive checks (Sec 6.4)
- Posterior predictive pvalues (Sec 6.3)

$$p_B = P(T(y^{rep}, \theta) \geq T(y, \theta)|y)$$

for a test statistic $T(y, \theta)$.

Hypothesis testing (Section 7.4)

From a Bayesian perspective,

Simple: $H_i : \theta = \theta_i$ Composite: $H_i : \theta \in (\theta_i, \theta_{i+1}]$

Treat all simple (or all composite) hypotheses as formal Bayesian parameter estimation. Treat a mix of simple and composite hypotheses as formal Bayesian tests.

Formal Bayesian tests

- require prior probabilities for each hypothesis, $p(H_i)$,
- require priors for parameters in non-point hypotheses, $p(\theta|H_i)$, and
- calculate posterior probabilities $p(H_i|y)$ which depend on
- the marginal likelihood, $p(y|H_i)$.

Decision theory (Sec 9.1)

In order to make a decision, a utility (or loss) function, i.e.

$U(\theta, \delta) = -L(\theta, \delta)$, must be set. Then the optimal Bayesian decision is to maximize expected utility (or minimize expected loss), i.e.

$$\operatorname{argmax}_{\delta} \int U(\theta, \delta) p(\theta) d\theta$$

where $p(\theta)$ represents your current state of belief, i.e. it could be a prior or a posterior depending on your perspective.

Stan

```
model = "  
data {  
  int<lower=0> N;  
  int<lower=0> n[N];  
  int<lower=0> y[N];  
  real s;  
}  
parameters {  
  real<lower=0,upper=1> mu;  
  real<lower=0> eta;  
}  
transformed parameters {  
  real<lower=0> alpha;  
  real<lower=0> beta;  
  alpha <- eta * mu;  
  beta <- eta * (1-mu);  
}  
model {  
  mu ~ beta(20,30);  
  eta ~ lognormal(0,s);  
  y ~ beta_binomial(n,alpha,beta);  
}  
generated quantities {  
  real<lower=0,upper=1> theta[N];  
  for (i in 1:N) theta[i] <- beta_rng(alpha+y[i], beta+n[i]-y[i]);  
}  
"
```