

# STAT 401A - Statistical Methods for Research Workers

## Regression diagnostics

Jarad Niemi (Dr. J)

Iowa State University

last updated: October 22, 2014

# All models are wrong!

George Box (Empirical Model-Building and Response Surfaces, 1987):

*All models are wrong, but some are useful.*

# All models are wrong!

George Box (Empirical Model-Building and Response Surfaces, 1987):

*All models are wrong, but some are useful.*

http:

[//stats.stackexchange.com/questions/57407/what-is-the-meaning-of-all-models-are-wrong-but-some-are-useful](http://stats.stackexchange.com/questions/57407/what-is-the-meaning-of-all-models-are-wrong-but-some-are-useful)

*“All models are wrong” that is, every model is wrong because it is a simplification of reality. Some models, especially in the “hard” sciences, are only a little wrong. They ignore things like friction or the gravitational effect of tiny bodies. Other models are a lot wrong - they ignore bigger things.*

# All models are wrong!

George Box (Empirical Model-Building and Response Surfaces, 1987):

*All models are wrong, but some are useful.*

http:

[//stats.stackexchange.com/questions/57407/what-is-the-meaning-of-all-models-are-wrong-but-some-are-useful](http://stats.stackexchange.com/questions/57407/what-is-the-meaning-of-all-models-are-wrong-but-some-are-useful)

*"All models are wrong" that is, every model is wrong because it is a simplification of reality. Some models, especially in the "hard" sciences, are only a little wrong. They ignore things like friction or the gravitational effect of tiny bodies. Other models are a lot wrong - they ignore bigger things.*

*"But some are useful" - simplifications of reality can be quite useful. They can help us explain, predict and understand the universe and all its various components.*

# All models are wrong!

George Box (Empirical Model-Building and Response Surfaces, 1987):

*All models are wrong, but some are useful.*

http:

[//stats.stackexchange.com/questions/57407/what-is-the-meaning-of-all-models-are-wrong-but-some-are-useful](http://stats.stackexchange.com/questions/57407/what-is-the-meaning-of-all-models-are-wrong-but-some-are-useful)

*“All models are wrong” that is, every model is wrong because it is a simplification of reality. Some models, especially in the “hard” sciences, are only a little wrong. They ignore things like friction or the gravitational effect of tiny bodies. Other models are a lot wrong - they ignore bigger things.*

*“But some are useful” - simplifications of reality can be quite useful. They can help us explain, predict and understand the universe and all its various components.*

*This isn’t just true in statistics! Maps are a type of model; they are wrong. But good maps are very useful.*

# Regression

The simpler linear regression model is

$$Y_i \stackrel{ind}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

# Regression

The simpler linear regression model is

$$Y_i \stackrel{ind}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

this can be rewritten as

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{ind}{\sim} N(0, \sigma^2)$$

# Regression

The simpler linear regression model is

$$Y_i \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

this can be rewritten as

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{\text{ind}}{\sim} N(0, \sigma^2)$$

where we estimate the errors via the residuals

$$r_i = \hat{e}_i = Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i).$$



# Regression

The simpler linear regression model is

$$Y_i \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

this can be rewritten as

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{\text{ind}}{\sim} N(0, \sigma^2)$$

where we estimate the errors via the residuals

$$r_i = \hat{e}_i = Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i).$$

Key assumptions are:

- Normality of the errors

# Regression

The simpler linear regression model is

$$Y_i \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

this can be rewritten as

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{\text{ind}}{\sim} N(0, \sigma^2)$$

where we estimate the errors via the residuals

$$r_i = \hat{e}_i = Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i).$$

Key assumptions are:

- Normality of the errors
- Constant variance of the errors

# Regression

The simpler linear regression model is

$$Y_i \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

this can be rewritten as

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{\text{ind}}{\sim} N(0, \sigma^2)$$

where we estimate the errors via the residuals

$$r_i = \hat{e}_i = Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i).$$

Key assumptions are:

- Normality of the errors
- Constant variance of the errors
- Independence of the errors

# Regression

The simpler linear regression model is

$$Y_i \stackrel{ind}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

this can be rewritten as

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{ind}{\sim} N(0, \sigma^2)$$


where we estimate the errors via the residuals

$$r_i = \hat{e}_i = Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i).$$

Key assumptions are:

- Normality of the errors
- Constant variance of the errors
- Independence of the errors
- Linearity between mean response and explanatory variable

# Histograms with best fitting bell curves



# Normal QQ-plot

## Definition

The quantile-quantile or qq-plot is an exploratory graphical device used to check the validity of a distributional assumption for a data set.

# Normal QQ-plot

## Definition

The quantile-quantile or qq-plot is an exploratory graphical device used to check the validity of a distributional assumption for a data set.

A normal qq-plot graphs the theoretical quantiles from a normal distribution versus the observed quantiles.

# Normal QQ-plot

## Definition


The quantile-quantile or qq-plot is an exploratory graphical device used to check the validity of a distributional assumption for a data set.

A normal qq-plot graphs the theoretical quantiles from a normal distribution versus the observed quantiles.

**Remark** The bottom line is that, if the distribution assumption is satisfied, the points should fall roughly along the  $y=x$  line.




# Normal




SAS swaps the x and y axes

# Normal ( $n=10$ )




SAS swaps the x and y axes

# Normal( $n=100$ )



SAS swaps the x and y axes


# Normal ( $n=1000$ )




SAS swaps the x and y axes

# Not normal ( $n=10$ )


normal




right-skewed



left-skewed




heavy-tailed



SAS swaps the x and y axes


# Not normal ( $n=100$ )



SAS swaps the x and y axes

# Not normal ( $n=1000$ )


normal




right-skewed



left-skewed



heavy-tailed



SAS swaps the x and y axes

# Constant variance

Recall the model

$$Y_i = \beta_0 + \beta_x X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$

so the variance for the  $e_i$  is constant.



# Constant variance

Recall the model

$$Y_i = \beta_0 + \beta_x X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$

so the variance for the  $e_i$  is constant.

To assess this assumption, we look at plots of residuals vs anything and look for patterns that show different “spreads”

# Constant variance

Recall the model

$$Y_i = \beta_0 + \beta_x X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$

so the variance for the  $e_i$  is constant.

To assess this assumption, we look at plots of residuals vs anything and look for patterns that show different “spreads”, e.g.

- funnels
- football shapes

# Constant variance

Recall the model

$$Y_i = \beta_0 + \beta_x X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$

so the variance for the  $e_i$  is constant.

To assess this assumption, we look at plots of residuals vs anything and look for patterns that show different “spreads”, e.g.

- funnels
- football shapes

The most common way this assumption is violated is by having increasing variance with increasing mean

# Constant variance

Recall the model

$$Y_i = \beta_0 + \beta_x X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$


so the variance for the  $e_i$  is constant.

To assess this assumption, we look at plots of residuals vs anything and look for patterns that show different “spreads”, e.g.


- funnels
- football shapes

The most common way this assumption is violated is by having increasing variance with increasing mean, thus we often look at a residuals vs predicted (fitted) mean plot.


# Constant variance



Predicted mean




Predicted mean




Predicted mean


# Constant variance



# Extreme non-constant variance (funnel)




# Extreme non-constant variance (funnel)







# Non-constant variance ( $n=10$ , $\sigma_2/\sigma_1 = 4$ )




# Non-constant variance ( $n=100$ , $\sigma_2/\sigma_1 = 4$ )



Predicted mean




Predicted mean




Predicted mean


# Non-constant variance ( $n=1000$ , $\sigma_2/\sigma_1 = 4$ )



Predicted mean




Predicted mean



Predicted mean

# Extreme non-constant variance (football)



# Independence

Lack of independence includes

- Cluster effect
- Serial correlation
- Spatial association

# Independence

Lack of independence includes

- Cluster effect
- Serial correlation
- Spatial association

Make plots of residuals vs relevant explanatory variables and look for patterns,

# Independence

Lack of independence includes

- Cluster effect
- Serial correlation
- Spatial association

Make plots of residuals vs relevant explanatory variables and look for patterns, e.g.

- Residuals vs groups

# Independence

Lack of independence includes

- Cluster effect
- Serial correlation
- Spatial association

Make plots of residuals vs relevant explanatory variables and look for patterns, e.g.

- Residuals vs groups
- Residuals vs time (or observation number)



# Independence

Lack of independence includes

- Cluster effect
- Serial correlation
- Spatial association

Make plots of residuals vs relevant explanatory variables and look for patterns, e.g.

- Residuals vs groups
- Residuals vs time (or observation number)
- Residuals vs spatial variable

# Summary

Often the best strategy is graphical exploration of the data,

# Summary

Often the best strategy is graphical exploration of the data, here are some relevant graphs:


# Summary

Often the best strategy is graphical exploration of the data, here are some relevant graphs:

- transformed response vs transformed explanatory
- transformed response vs transformed explanatory
- qqplot of residuals
- residual vs fitted value
- residual vs explanatory
- residual vs observation number
- residual vs any other variable

# Linearity

Assess using scatterplots of (transformed) response vs (transformed) explanatory variable:



# Testing Composite hypotheses

Comparing two models

- $H_0$  : (reduced)
- $H_1$  : (full)

# Testing Composite hypotheses

Comparing two models

- $H_0$  : (reduced)
- $H_1$  : (full)

Do the following

1. Calculate extra sum of squares.
2. Calculate extra degrees of freedom
3. Calculate

$$\text{F-statistic} = \frac{\text{Extra sum of squares} / \text{Extra degrees of freedom}}{\hat{\sigma}_{full}^2}$$

4. Compare this to an F-distribution with

- numerator degrees of freedom = extra degrees of freedom
- denominator degrees of freedom = degrees of freedom in estimating  $\hat{\sigma}_{full}^2$

# Lack-of-fit F-test

Let  $Y_{ij}$  be the  $i^{th}$  observation from the  $j^{th}$  group where the group is defined by those observations having the same explanatory variable value ( $X_j$ ).



## Lack-of-fit F-test

Let  $Y_{ij}$  be the  $i^{th}$  observation from the  $j^{th}$  group where the group is defined by those observations having the same explanatory variable value ( $X_j$ ).

Two models:

$$\text{ANOVA:} \quad Y_{ij} \stackrel{\text{ind}}{\sim} N(\mu_j, \sigma^2)$$

## Lack-of-fit F-test

Let  $Y_{ij}$  be the  $i^{th}$  observation from the  $j^{th}$  group where the group is defined by those observations having the same explanatory variable value ( $X_j$ ).

Two models:

$$\text{ANOVA: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\mu_j, \sigma^2) \quad (\text{full})$$

$$\text{Regression: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_j, \sigma^2) \quad (\text{reduced})$$

## Lack-of-fit F-test

Let  $Y_{ij}$  be the  $i^{th}$  observation from the  $j^{th}$  group where the group is defined by those observations having the same explanatory variable value ( $X_j$ ).

Two models:

$$\text{ANOVA: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\mu_j, \sigma^2) \quad (\text{full})$$

$$\text{Regression: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_j, \sigma^2) \quad (\text{reduced})$$

## Lack-of-fit F-test

Let  $Y_{ij}$  be the  $i^{th}$  observation from the  $j^{th}$  group where the group is defined by those observations having the same explanatory variable value ( $X_j$ ).

Two models:

$$\text{ANOVA: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\mu_j, \sigma^2) \quad (\text{full})$$

$$\text{Regression: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_j, \sigma^2) \quad (\text{reduced})$$

- Regression model is reduced:

## Lack-of-fit F-test

Let  $Y_{ij}$  be the  $i^{th}$  observation from the  $j^{th}$  group where the group is defined by those observations having the same explanatory variable value ( $X_j$ ).

Two models:

$$\text{ANOVA: } Y_{ij} \overset{\text{ind}}{\sim} N(\mu_j, \sigma^2) \quad (\text{full})$$

$$\text{Regression: } Y_{ij} \overset{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_j, \sigma^2) \quad (\text{reduced})$$

- Regression model is reduced:
  - ANOVA has  $J$  parameters for the mean

## Lack-of-fit F-test

Let  $Y_{ij}$  be the  $i^{th}$  observation from the  $j^{th}$  group where the group is defined by those observations having the same explanatory variable value ( $X_j$ ).

Two models:

$$\text{ANOVA: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\mu_j, \sigma^2) \quad (\text{full})$$

$$\text{Regression: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_j, \sigma^2) \quad (\text{reduced})$$

- Regression model is reduced:
  - ANOVA has  $J$  parameters for the mean
  - Regression has 2 parameters for the mean

## Lack-of-fit F-test

Let  $Y_{ij}$  be the  $i^{th}$  observation from the  $j^{th}$  group where the group is defined by those observations having the same explanatory variable value ( $X_j$ ).

Two models:

$$\text{ANOVA: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\mu_j, \sigma^2) \quad (\text{full})$$

$$\text{Regression: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_j, \sigma^2) \quad (\text{reduced})$$

- Regression model is reduced:
  - ANOVA has  $J$  parameters for the mean
  - Regression has 2 parameters for the mean
  - Set  $\mu_j = \beta_0 + \beta_1 X_j$ .

# Lack-of-fit F-test

Let  $Y_{ij}$  be the  $i^{th}$  observation from the  $j^{th}$  group where the group is defined by those observations having the same explanatory variable value ( $X_j$ ).

Two models:

$$\text{ANOVA: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\mu_j, \sigma^2) \quad (\text{full})$$

$$\text{Regression: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_j, \sigma^2) \quad (\text{reduced})$$

- Regression model is reduced:
  - ANOVA has  $J$  parameters for the mean
  - Regression has 2 parameters for the mean
  - Set  $\mu_j = \beta_0 + \beta_1 X_j$ .
- Small pvalues indicate a lack-of-fit, i.e. the reduced model is not adequate.



# Lack-of-fit F-test

Let  $Y_{ij}$  be the  $i^{th}$  observation from the  $j^{th}$  group where the group is defined by those observations having the same explanatory variable value ( $X_j$ ).


Two models:

$$\text{ANOVA: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\mu_j, \sigma^2) \quad (\text{full})$$


$$\text{Regression: } Y_{ij} \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_j, \sigma^2) \quad (\text{reduced})$$

- Regression model is reduced:
  - ANOVA has  $J$  parameters for the mean
  - Regression has 2 parameters for the mean
  - Set  $\mu_j = \beta_0 + \beta_1 X_j$ .
- Small pvalues indicate a lack-of-fit, i.e. the reduced model is not adequate.
- Lack-of-fit F-test requires multiple observations at a few  $X_j$  values!

# Telomere length



# Telomere length



# SAS code

```
DATA t;
  INFILE 'telomeres.csv' DSD FIRSTOBS=2;
  INPUT years length;

PROC REG DATA=t;
  MODEL length = years / CLB LACKFIT;
  RUN;
```

The REG Procedure  
 Model: MODEL1  
 Dependent Variable: length

Number of Observations Read	39
Number of Observations Used	39

## Analysis of Variance

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	1	0.22777	0.22777	8.42	0.0062
Error	37	1.00033	0.02704		
Lack of Fit	9	0.18223	0.02025	0.69	0.7093
Pure Error	28	0.81810	0.02922		
Corrected Total	38	1.22810			

```
# Use as.factor to turn a continuous variable into a categorical variable
m_anova = lm(telomere.length ~ as.factor(years), Telomeres)
m_reg   = lm(telomere.length ~ years, Telomeres)
anova(m_reg, m_anova)
```

#### Analysis of Variance Table

Model 1: telomere.length ~ years

Model 2: telomere.length ~ as.factor(years)

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	37	1.000				
2	28	0.818	9	0.182	0.69	0.71

```
# Use as.factor to turn a continuous variable into a categorical variable
m_anova = lm(telomere.length ~ as.factor(years), Telomeres)
m_reg   = lm(telomere.length ~ years, Telomeres)
anova(m_reg, m_anova)
```

Analysis of Variance Table

Model 1: telomere.length ~ years

Model 2: telomere.length ~ as.factor(years)

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)
1	37	1.000				
2	28	0.818	9	0.182	0.69	0.71

No evidence of a lack of fit.

# Lack-of-fit F-test summary

- Lack-of-fit F-test tests the assumption of linearity

# Lack-of-fit F-test summary

- Lack-of-fit F-test tests the assumption of linearity
- Needs multiple observations at various explanatory variable values



# Lack-of-fit F-test summary

- Lack-of-fit F-test tests the assumption of linearity
- Needs multiple observations at various explanatory variable values
- Small pvalue indicates a lack-of-fit, i.e. means are not linear

# Lack-of-fit F-test summary

- Lack-of-fit F-test tests the assumption of linearity
- Needs multiple observations at various explanatory variable values
- Small pvalue indicates a lack-of-fit, i.e. means are not linear
  - Transform response, e.g. log

# Lack-of-fit F-test summary

- Lack-of-fit F-test tests the assumption of linearity
- Needs multiple observations at various explanatory variable values
- Small pvalue indicates a lack-of-fit, i.e. means are not linear
  - Transform response, e.g. log
  - Transform explanatory variable

# Lack-of-fit F-test summary

- Lack-of-fit F-test tests the assumption of linearity
- Needs multiple observations at various explanatory variable values
- Small pvalue indicates a lack-of-fit, i.e. means are not linear
  - Transform response, e.g. log
  - Transform explanatory variable
  - Add other explanatory variables

# Interpretations using logs

The most common transformation of either the response or explanatory is to take logarithms

# Interpretations using logs

The most common transformation of either the response or explanatory is to take logarithms because

- linearity will often then be approximately true,

# Interpretations using logs

The most common transformation of either the response or explanatory is to take logarithms because

- linearity will often then be approximately true,
- the variance will likely be approximately constant, and

# Interpretations using logs

The most common transformation of either the response or explanatory is to take logarithms because

- linearity will often then be approximately true,
- the variance will likely be approximately constant, and
- there is a (relatively) convenient interpretation.



# Interpretations using logs

The most common transformation of either the response or explanatory is to take logarithms because

- linearity will often then be approximately true,
- the variance will likely be approximately constant, and
- there is a (relatively) convenient interpretation.

# Interpretations using logs

The most common transformation of either the response or explanatory is to take logarithms because

- linearity will often then be approximately true,
- the variance will likely be approximately constant, and
- there is a (relatively) convenient interpretation.

We will talk about interpretation of  $\beta_0$  and  $\beta_1$  when

- only the response is logged,

# Interpretations using logs

The most common transformation of either the response or explanatory is to take logarithms because

- linearity will often then be approximately true,
- the variance will likely be approximately constant, and
- there is a (relatively) convenient interpretation.

We will talk about interpretation of  $\beta_0$  and  $\beta_1$  when

- only the response is logged,
- only the explanatory variable is logged, and

# Interpretations using logs

The most common transformation of either the response or explanatory is to take logarithms because

- linearity will often then be approximately true,
- the variance will likely be approximately constant, and
- there is a (relatively) convenient interpretation.

We will talk about interpretation of  $\beta_0$  and  $\beta_1$  when

- only the response is logged,
- only the explanatory variable is logged, and
- when both are logged.

# Neither response nor explanatory variable are logged

If

$$E[Y|X] = \beta_0 + \beta_1 X$$

# Neither response nor explanatory variable are logged

If

$$E[Y|X] = \beta_0 + \beta_1 X$$

, then

- $\beta_0$  is the expected response when  $X$  is zero and

## Neither response nor explanatory variable are logged

If

$$E[Y|X] = \beta_0 + \beta_1 X$$

, then

- $\beta_0$  is the expected response when  $X$  is zero and
- $\beta_1$  is the expected change in the response for a one unit increase in the explanatory variable.

## Neither response nor explanatory variable are logged

If

$$E[Y|X] = \beta_0 + \beta_1 X$$

, then

- $\beta_0$  is the expected response when  $X$  is zero and
- $\beta_1$  is the expected change in the response for a one unit increase in the explanatory variable.



# Neither response nor explanatory variable are logged

If

$$E[Y|X] = \beta_0 + \beta_1 X$$

, then

- $\beta_0$  is the expected response when  $X$  is zero and
- $\beta_1$  is the expected change in the response for a one unit increase in the explanatory variable.

For the following discussion,

- $Y$  is always going to be the original response and

## Neither response nor explanatory variable are logged

If

$$E[Y|X] = \beta_0 + \beta_1 X$$

, then

- $\beta_0$  is the expected response when  $X$  is zero and
- $\beta_1$  is the expected change in the response for a one unit increase in the explanatory variable.

For the following discussion,

- $Y$  is always going to be the original response and
- $X$  is always going to be the original explanatory variable.

# Example

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

## Example

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

Then, if

$$E[Y|X] = \beta_0 + \beta_1 X$$

- $\beta_0$  is the expected corn yield per acre when fertilizer level is zero and

# Example

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

Then, if

$$E[Y|X] = \beta_0 + \beta_1 X$$

- $\beta_0$  is the expected corn yield per acre when fertilizer level is zero and
- $\beta_1$  is the expected change in corn yield per acre when fertilizer is increase by 1 lbs/acre.

# Response is logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 X \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 X}$$

# Response is logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 X \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 X}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $X$  is zero and

# Response is logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 X \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 X}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $X$  is zero and
- $\beta_1$  is the expected change in  $\log(Y)$  for a one unit increase in the explanatory variable.



# Response is logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 X \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 X}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $X$  is zero and
- $\beta_1$  is the expected change in  $\log(Y)$  for a one unit increase in the explanatory variable.

# Response is logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 X \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 X}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $X$  is zero and
- $\beta_1$  is the expected change in  $\log(Y)$  for a one unit increase in the explanatory variable.

Alternatively,

- $\exp(\beta_0)$  is the median of  $Y$  when  $X$  is zero and

# Response is logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 X \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 X}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $X$  is zero and
- $\beta_1$  is the expected change in  $\log(Y)$  for a one unit increase in the explanatory variable.

Alternatively,

- $\exp(\beta_0)$  is the median of  $Y$  when  $X$  is zero and
- $\exp(\beta_1)$  is the multiplicative effect on the median of  $Y$  for a one unit increase in the explanatory variable.

# Response is logged

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

# Response is logged

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

Then, if

$$E[\log(Y)|X] = \beta_0 + \beta_1 X \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 X}$$

- $\exp(\beta_0)$  is the median corn yield per acre when fertilizer level is zero and

# Response is logged

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

Then, if

$$E[\log(Y)|X] = \beta_0 + \beta_1 X \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 X}$$

- $\exp(\beta_0)$  is the median corn yield per acre when fertilizer level is zero and
- $\exp(\beta_1)$  is the multiplicative effect in median corn yield per acre when fertilizer is increase by 1 lbs/acre.

## Explanatory variable is logged

If

$$E[Y|X] = \beta_0 + \beta_1 \log(X)$$

## Explanatory variable is logged

If

$$E[Y|X] = \beta_0 + \beta_1 \log(X)$$

, then

- $\beta_0$  is the expected response when  $\log(X)$  is zero and



## Explanatory variable is logged

If

$$E[Y|X] = \beta_0 + \beta_1 \log(X)$$

, then

- $\beta_0$  is the expected response when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in the response for a one unit increase in  $\log(X)$ .

## Explanatory variable is logged

If

$$E[Y|X] = \beta_0 + \beta_1 \log(X)$$

, then

- $\beta_0$  is the expected response when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in the response for a one unit increase in  $\log(X)$ .

## Explanatory variable is logged

If

$$E[Y|X] = \beta_0 + \beta_1 \log(X)$$

, then

- $\beta_0$  is the expected response when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in the response for a one unit increase in  $\log(X)$ .

Alternatively,

- $\beta_0$  is the expected response when  $X$  is 1 and

## Explanatory variable is logged

If

$$E[Y|X] = \beta_0 + \beta_1 \log(X)$$

, then

- $\beta_0$  is the expected response when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in the response for a one unit increase in  $\log(X)$ .

Alternatively,

- $\beta_0$  is the expected response when  $X$  is 1 and
- $\beta_1 \log(d)$  is the expected change in the response when  $X$  increase multiplicatively by  $d$ , e.g.

## Explanatory variable is logged

If

$$E[Y|X] = \beta_0 + \beta_1 \log(X)$$

, then

- $\beta_0$  is the expected response when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in the response for a one unit increase in  $\log(X)$ .

Alternatively,

- $\beta_0$  is the expected response when  $X$  is 1 and
- $\beta_1 \log(d)$  is the expected change in the response when  $X$  increase multiplicatively by  $d$ , e.g.
  - $\beta_1 \log(2)$  is the expected change in the response for each doubling of  $X$   
or

## Explanatory variable is logged

If

$$E[Y|X] = \beta_0 + \beta_1 \log(X)$$

, then

- $\beta_0$  is the expected response when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in the response for a one unit increase in  $\log(X)$ .

Alternatively,

- $\beta_0$  is the expected response when  $X$  is 1 and
- $\beta_1 \log(d)$  is the expected change in the response when  $X$  increase multiplicatively by  $d$ , e.g.
  - $\beta_1 \log(2)$  is the expected change in the response for each doubling of  $X$  or
  - $\beta_1 \log(10)$  is the expected change in the response for each ten-fold increase in  $X$ .

# Explanatory variable is logged

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

## Explanatory variable is logged

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

Then, if

$$E[Y|X] = \beta_0 + \beta_1 \log(X)$$

- $\beta_0$  is the median corn yield per acre when fertilizer level is 1 lb/acre and



## Explanatory variable is logged

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

Then, if

$$E[Y|X] = \beta_0 + \beta_1 \log(X)$$

- $\beta_0$  is the median corn yield per acre when fertilizer level is 1 lb/acre and
- $\beta_1 \log(2)$  is the expected change in corn yield when fertilizer level is doubled.

## Both response and explanatory variables are logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 \log(X) \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 \log(X)} = e^{\beta_0} X^{\beta_1}$$

## Both response and explanatory variables are logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 \log(X) \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 \log(X)} = e^{\beta_0} X^{\beta_1}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $\log(X)$  is zero and

## Both response and explanatory variables are logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 \log(X) \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 \log(X)} = e^{\beta_0} X^{\beta_1}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in  $\log(Y)$  for a one unit increase in  $\log(X)$ .

## Both response and explanatory variables are logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 \log(X) \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 \log(X)} = e^{\beta_0} X^{\beta_1}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in  $\log(Y)$  for a one unit increase in  $\log(X)$ .

## Both response and explanatory variables are logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 \log(X) \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 \log(X)} = e^{\beta_0} X^{\beta_1}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in  $\log(Y)$  for a one unit increase in  $\log(X)$ .

Alternatively,

- $\exp(\beta_0)$  is the median of  $Y$  when  $X$  is 1 and

## Both response and explanatory variables are logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 \log(X) \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 \log(X)} = e^{\beta_0} X^{\beta_1}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in  $\log(Y)$  for a one unit increase in  $\log(X)$ .

Alternatively,

- $\exp(\beta_0)$  is the median of  $Y$  when  $X$  is 1 and
- $d^{\beta_1}$  is the multiplicative change in the median of the response when  $X$  increase multiplicatively by  $d$ , e.g.

## Both response and explanatory variables are logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 \log(X) \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 \log(X)} = e^{\beta_0} X^{\beta_1}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in  $\log(Y)$  for a one unit increase in  $\log(X)$ .

Alternatively,

- $\exp(\beta_0)$  is the median of  $Y$  when  $X$  is 1 and
- $d^{\beta_1}$  is the multiplicative change in the median of the response when  $X$  increase multiplicatively by  $d$ , e.g.
  - $2^{\beta_1}$  is the multiplicative effect on the median of the response for each doubling of  $X$  or



## Both response and explanatory variables are logged

If

$$E[\log(Y)|X] = \beta_0 + \beta_1 \log(X) \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 \log(X)} = e^{\beta_0} X^{\beta_1}$$

, then

- $\beta_0$  is the expected  $\log(Y)$  when  $\log(X)$  is zero and
- $\beta_1$  is the expected change in  $\log(Y)$  for a one unit increase in  $\log(X)$ .

Alternatively,

- $\exp(\beta_0)$  is the median of  $Y$  when  $X$  is 1 and
- $d^{\beta_1}$  is the multiplicative change in the median of the response when  $X$  increase multiplicatively by  $d$ , e.g.
  - $2^{\beta_1}$  is the multiplicative effect on the median of the response for each doubling of  $X$  or
  - $10^{\beta_1}$  is the multiplicative effect on the median of the response for each ten-fold increase in  $X$ .

# Both response and explanatory variables are logged

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

## Both response and explanatory variables are logged

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

Then, if

$$E[\log(Y)|X] = \beta_0 + \beta_1 \log(X) \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 \log(X)} = e^{\beta_0} X^{\beta_1}$$

- $\exp(\beta_0)$  is the median corn yield per acre when fertilizer level is 1 lb/acre and

## Both response and explanatory variables are logged

Suppose

- $Y$  is corn yield per acre
- $X$  is fertilizer level in lbs/acre

Then, if

$$E[\log(Y)|X] = \beta_0 + \beta_1 \log(X) \quad \text{or} \quad \text{Median}\{Y|X\} = e^{\beta_0} e^{\beta_1 \log(X)} = e^{\beta_0} X^{\beta_1}$$

- $\exp(\beta_0)$  is the median corn yield per acre when fertilizer level is 1 lb/acre and
- $2^{\beta_1}$  is the multiplicative effect on median corn yield per acre when fertilizer level doubles.