

M5S4 - Practice with CIs

Professor Jarad Niemi

STAT 226 - Iowa State University

October 17, 2018

Outline

- Constructing confidence intervals
 - Review
 - Bayesian

Confidence Interval Review

Two methods of constructing confidence intervals for the population mean μ :

$$\bar{x} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \quad \text{and} \quad \bar{x} \pm t_{n-1, \alpha/2} \frac{s}{\sqrt{n}}$$

where

- \bar{x} is the sample mean,
- s is the sample standard deviation,
- n is the sample size,
- σ is the known population standard deviation,
- $z_{\alpha/2}$ is the z critical value such that $P(Z > z_{\alpha/2}) = \alpha/2$,
- $t_{n-1, \alpha/2}$ is the t critical value such that $P(T_{n-1} > t_{n-1, \alpha/2}) = \alpha/2$ and $n - 1$ is the degrees of freedom,
- α is the significance (error) level, and
- $100(1 - \alpha)\%$ is the confidence level.

The interpretation of a $100(1 - \alpha)\%$ confidence interval is that, on average, $100(1 - \alpha)\%$ of the intervals constructed with this procedure will cover μ .

Deciding which method to use

Recall that all our confidence interval formulas require the observations be independent and identically distributed. We usually accomplish this by taking a **random sample** from the population.

Data	σ	Sample size	Interval
Normal	Known	any	z is exact
Normal	Unknown	any	t is exact
Not normal	Known	large	z is approximate
Not normal	Unknown	any	t is approximate

Estimator for a proportion

Let $X_i \stackrel{iid}{\sim} \text{Ber}(p)$, then $Y = \sum_{i=1}^n X_i \sim \text{Bin}(n, p)$. An estimator for p is

$$\hat{p} = \frac{Y}{n}$$

with

$$E[\hat{p}] = E\left[\frac{Y}{n}\right] = \frac{E[Y]}{n} = \frac{np}{n}$$

thus \hat{p} is an unbiased estimator and

$$\text{Var}[\hat{p}] = \text{Var}\left[\frac{Y}{n}\right] = \frac{1}{n^2} \text{Var}[Y] = \frac{np(1-p)}{n^2} = \frac{p(1-p)}{n}$$

thus

$$SD[\hat{p}] = \sqrt{\text{Var}[\hat{p}]} = \sqrt{\frac{p(1-p)}{n}}.$$

Confidence interval for a proportion

To construct a $100(1 - \alpha)\%$ confidence interval for p , we use the formula

$$\hat{p} \pm z_{\alpha/2} \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$$

where $SE[\hat{p}] = \sqrt{\hat{p}(1 - \hat{p})/n}$, i.e. our estimate of the SD.

It is common in polling to report \hat{p} and the **margin of error**
 $z_{\alpha/2} \sqrt{\hat{p}(1 - \hat{p})/n}$.

2018 Iowa Governor Poll

In the most recent Des Moines register poll of 555 likely voters

https://www.realclearpolitics.com/epolls/2018/governor/ia/iowa_governor_reynolds_vs_hubbell-6477.html

43% indicated they would vote for Fred Hubbell with a margin of error of 4.2.

Thus a 95% confidence interval for the actual proportion who say they would vote for Fred Hubbell is

$$0.43 \pm 0.042 = (0.388, 0.472) = (38.8\%, 47.2\%).$$

The margin of error calculation is

$$2 \cdot \sqrt{\frac{0.43(1 - 0.43)}{555}} = 0.042 = 4.2\%.$$

The best resource for combining all the information from polls is 538:

<https://projects.fivethirtyeight.com/2018-midterm-election-forecast/governor/>