

STAT 401A - Statistical Methods for Research Workers

Simple linear regression

Jarad Niemi (Dr. J)

Iowa State University

last updated: October 20, 2014

Simple Linear Regression

Recall the one-way ANOVA model:

$$Y_{ij} \stackrel{ind}{\sim} N(\mu_j, \sigma^2)$$

where Y_{ij} is the observation for individual i in group j .

Simple Linear Regression

Recall the one-way ANOVA model:

$$Y_{ij} \stackrel{ind}{\sim} N(\mu_j, \sigma^2)$$

where Y_{ij} is the observation for individual i in group j .

The **simple linear regression** model is

$$Y_i \stackrel{ind}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

where Y_i and X_i are the response and explanatory variable, respectively, for individual i .

Simple Linear Regression

Recall the one-way ANOVA model:

$$Y_{ij} \stackrel{\text{ind}}{\sim} N(\mu_j, \sigma^2)$$

where Y_{ij} is the observation for individual i in group j .


The **simple linear regression** model is

$$Y_i \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

where Y_i and X_i are the response and explanatory variable, respectively, for individual i .

Terminology (all of these are equivalent):

response	explanatory
outcome	covariate
dependent	independent
endogenous	exogenous



Telomere length

<http://www.pnas.org/content/101/49/17312>

People who are stressed over long periods tend to look haggard, and it is commonly thought that psychological stress leads to premature aging and the earlier onset of diseases of aging.

...

This design allowed us to examine the importance of perceived stress and measures of objective stress (caregiving status and chronicity of caregiving stress based on the number of years since a child's diagnosis).

...

Telomere length values were measured from DNA by a quantitative PCR assay that determines the relative ratio of telomere repeat copy number to single-copy gene copy number (T/S ratio) in experimental samples as compared with a reference DNA sample.

Parameter interpretation

$$E[Y_i|X_i = x] = \beta_0 + \beta_1 x \quad V[Y_i|X_i = x] = \sigma^2$$

Parameter interpretation

$$E[Y_i|X_i = x] = \beta_0 + \beta_1 x \quad V[Y_i|X_i = x] = \sigma^2$$

- If $X_i = 0$, then $E[Y_i|X_i = 0] = \beta_0$.

Parameter interpretation

$$E[Y_i|X_i = x] = \beta_0 + \beta_1 x \quad V[Y_i|X_i = x] = \sigma^2$$

- If $X_i = 0$, then $E[Y_i|X_i = 0] = \beta_0$.

β_0 is the expected response when the explanatory variable is zero.

Parameter interpretation

$$E[Y_i|X_i = x] = \beta_0 + \beta_1 x \quad V[Y_i|X_i = x] = \sigma^2$$

- If $X_i = 0$, then $E[Y_i|X_i = 0] = \beta_0$.
 β_0 is the expected response when the explanatory variable is zero.
- If X_i increases from x to $x + 1$, then

$$E[Y_i|X_i = x + 1] = \beta_0 + \beta_1 x + \beta_1$$

Parameter interpretation

$$E[Y_i|X_i = x] = \beta_0 + \beta_1 x \quad V[Y_i|X_i = x] = \sigma^2$$

- If $X_i = 0$, then $E[Y_i|X_i = 0] = \beta_0$.
 β_0 is the expected response when the explanatory variable is zero.
- If X_i increases from x to $x + 1$, then

$$\begin{aligned} E[Y_i|X_i = x + 1] &= \beta_0 + \beta_1 x + \beta_1 \\ E[Y_i|X_i = x] &= \beta_0 + \beta_1 x \end{aligned}$$

Parameter interpretation

$$E[Y_i|X_i = x] = \beta_0 + \beta_1 x \quad V[Y_i|X_i = x] = \sigma^2$$

- If $X_i = 0$, then $E[Y_i|X_i = 0] = \beta_0$.
 β_0 is the expected response when the explanatory variable is zero.
- If X_i increases from x to $x + 1$, then

$$\begin{array}{rcl} E[Y_i|X_i = x + 1] & = & \beta_0 + \beta_1 x + \beta_1 \\ - E[Y_i|X_i = x] & = & \beta_0 + \beta_1 x \\ \hline & = & \beta_1 \end{array}$$

Parameter interpretation

$$E[Y_i|X_i = x] = \beta_0 + \beta_1 x \quad V[Y_i|X_i = x] = \sigma^2$$

- If $X_i = 0$, then $E[Y_i|X_i = 0] = \beta_0$.
 β_0 is the expected response when the explanatory variable is zero.
- If X_i increases from x to $x + 1$, then

$$\begin{array}{rcl} E[Y_i|X_i = x + 1] & = & \beta_0 + \beta_1 x + \beta_1 \\ - E[Y_i|X_i = x] & = & \beta_0 + \beta_1 x \\ \hline & = & \beta_1 \end{array}$$

β_1 is the expected increase in the response for each unit increase in the explanatory variable.

Parameter interpretation

$$E[Y_i|X_i = x] = \beta_0 + \beta_1 x \quad V[Y_i|X_i = x] = \sigma^2$$

- If $X_i = 0$, then $E[Y_i|X_i = 0] = \beta_0$.
 β_0 is the expected response when the explanatory variable is zero.
- If X_i increases from x to $x + 1$, then

$$\begin{array}{rcl} E[Y_i|X_i = x + 1] & = & \beta_0 + \beta_1 x + \beta_1 \\ - E[Y_i|X_i = x] & = & \beta_0 + \beta_1 x \\ \hline & = & \beta_1 \end{array}$$

β_1 is the expected increase in the response for each unit increase in the explanatory variable.

- σ is the standard deviation of the response for a fixed value of the explanatory variable.

Remove the mean:

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$

Remove the mean:

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$

So the error is

$$e_i = Y_i - (\beta_0 + \beta_1 X_i)$$

Remove the mean:

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$

So the error is

$$e_i = Y_i - (\beta_0 + \beta_1 X_i)$$

which we approximate by the **residual**

$$r_i = \hat{e}_i = Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i)$$

Remove the mean:

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$

So the error is

$$e_i = Y_i - (\beta_0 + \beta_1 X_i)$$

which we approximate by the **residual**

$$r_i = \hat{e}_i = Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i)$$

The least squares, maximum likelihood, and Bayesian estimators are

Remove the mean:

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$

So the error is

$$e_i = Y_i - (\beta_0 + \beta_1 X_i)$$

which we approximate by the **residual**

$$r_i = \hat{e}_i = Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i)$$

The least squares, maximum likelihood, and Bayesian estimators are

$$\begin{aligned}\hat{\beta}_1 &= SXY / SXX \\ \hat{\beta}_0 &= \bar{Y} - \hat{\beta}_1 \bar{X} \\ \hat{\sigma}^2 &= SSE / (n - 2) \quad df = n - 2\end{aligned}$$

Remove the mean:

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$

So the error is

$$e_i = Y_i - (\beta_0 + \beta_1 X_i)$$

which we approximate by the **residual**

$$r_i = \hat{e}_i = Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i)$$

The least squares, maximum likelihood, and Bayesian estimators are

$$\begin{aligned} \hat{\beta}_1 &= SXY / SXX \\ \hat{\beta}_0 &= \bar{Y} - \hat{\beta}_1 \bar{X} \\ \hat{\sigma}^2 &= SSE / (n - 2) \quad df = n - 2 \end{aligned}$$

$$\begin{aligned} \bar{X} &= \frac{1}{n} \sum_{i=1}^n X_i \\ \bar{Y} &= \frac{1}{n} \sum_{i=1}^n Y_i \end{aligned}$$

Remove the mean:

$$Y_i = \beta_0 + \beta_1 X_i + e_i \quad e_i \stackrel{iid}{\sim} N(0, \sigma^2)$$

So the error is

$$e_i = Y_i - (\beta_0 + \beta_1 X_i)$$

which we approximate by the **residual**

$$r_i = \hat{e}_i = Y_i - (\hat{\beta}_0 + \hat{\beta}_1 X_i)$$

The least squares, maximum likelihood, and Bayesian estimators are

$$\begin{aligned}\hat{\beta}_1 &= SXY / SXX \\ \hat{\beta}_0 &= \bar{Y} - \hat{\beta}_1 \bar{X} \\ \hat{\sigma}^2 &= SSE / (n - 2) \quad df = n - 2\end{aligned}$$

$$\begin{aligned}\bar{X} &= \frac{1}{n} \sum_{i=1}^n X_i \\ \bar{Y} &= \frac{1}{n} \sum_{i=1}^n Y_i\end{aligned}$$

$$\begin{aligned}SXY &= \sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y}) \\ SXX &= \sum_{i=1}^n (X_i - \bar{X})(X_i - \bar{X}) = \sum_{i=1}^n (X_i - \bar{X})^2 \\ SSE &= \sum_{i=1}^n r_i^2\end{aligned}$$

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

We quantify this uncertainty using their standard errors:

$$SE(\beta_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} \quad df = n - 2$$

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

We quantify this uncertainty using their standard errors:

$$SE(\beta_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} \quad df = n - 2$$

$$SE(\beta_1) = \hat{\sigma} \sqrt{\frac{1}{(n-1)s_X^2}} \quad df = n - 2$$

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

We quantify this uncertainty using their standard errors:

$$SE(\beta_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} \quad df = n - 2$$

$$SE(\beta_1) = \hat{\sigma} \sqrt{\frac{1}{(n-1)s_X^2}} \quad df = n - 2$$

$$s_X^2 = SXX / (n - 1)$$

$$s_Y^2 = SY / (n - 1)$$

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

We quantify this uncertainty using their standard errors:

$$SE(\beta_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} \quad df = n - 2$$

$$SE(\beta_1) = \hat{\sigma} \sqrt{\frac{1}{(n-1)s_X^2}} \quad df = n - 2$$

$$s_X^2 = SXX / (n - 1)$$

$$s_Y^2 = SY / (n - 1)$$

$$SY = \sum_{i=1}^n (Y_i - \bar{Y})^2$$

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

We quantify this uncertainty using their standard errors:

$$SE(\beta_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} \quad df = n - 2$$

$$SE(\beta_1) = \hat{\sigma} \sqrt{\frac{1}{(n-1)s_X^2}} \quad df = n - 2$$

$$s_X^2 = SXX / (n - 1)$$

$$s_Y^2 = SYY / (n - 1)$$

$$SYY = \sum_{i=1}^n (Y_i - \bar{Y})^2$$

$$r_{XY} = \frac{SXY / (n-1)}{s_X s_Y}$$

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

We quantify this uncertainty using their standard errors:

$$SE(\beta_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} \quad df = n - 2$$

$$SE(\beta_1) = \hat{\sigma} \sqrt{\frac{1}{(n-1)s_X^2}} \quad df = n - 2$$

$$s_X^2 = SXX / (n - 1)$$

$$s_Y^2 = SYY / (n - 1)$$

$$SYY = \sum_{i=1}^n (Y_i - \bar{Y})^2$$

$$r_{XY} = \frac{SXY / (n-1)}{s_X s_Y}$$

correlation coefficient

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

We quantify this uncertainty using their standard errors:

$$SE(\beta_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} \quad df = n - 2$$

$$SE(\beta_1) = \hat{\sigma} \sqrt{\frac{1}{(n-1)s_X^2}} \quad df = n - 2$$

$$s_X^2 = SXX / (n - 1)$$

$$s_Y^2 = SYY / (n - 1)$$

$$SYY = \sum_{i=1}^n (Y_i - \bar{Y})^2$$

$$r_{XY} = \frac{SXY / (n-1)}{s_X s_Y}$$

correlation coefficient

$$R^2 = r_{XY}^2$$

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

We quantify this uncertainty using their standard errors:

$$SE(\beta_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} \quad df = n - 2$$

$$SE(\beta_1) = \hat{\sigma} \sqrt{\frac{1}{(n-1)s_X^2}} \quad df = n - 2$$

$$s_X^2 = SXX / (n - 1)$$

$$s_Y^2 = SYY / (n - 1)$$

$$SYY = \sum_{i=1}^n (Y_i - \bar{Y})^2$$

$$r_{XY} = \frac{SXY / (n-1)}{s_X s_Y}$$

$$R^2 = r_{XY}^2 = \frac{SST - SSE}{SST}$$

correlation coefficient

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

We quantify this uncertainty using their standard errors:

$$SE(\beta_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} \quad df = n - 2$$

$$SE(\beta_1) = \hat{\sigma} \sqrt{\frac{1}{(n-1)s_X^2}} \quad df = n - 2$$

$$s_X^2 = SXX / (n - 1)$$

$$s_Y^2 = SY / (n - 1)$$

$$SYY = \sum_{i=1}^n (Y_i - \bar{Y})^2$$

$$r_{XY} = \frac{SXY / (n-1)}{s_X s_Y}$$

$$R^2 = r_{XY}^2 = \frac{SST - SSE}{SST} \quad \begin{array}{l} \text{correlation coefficient} \\ \text{coefficient of determination} \end{array}$$

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

We quantify this uncertainty using their standard errors:

$$SE(\beta_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} \quad df = n - 2$$

$$SE(\beta_1) = \hat{\sigma} \sqrt{\frac{1}{(n-1)s_X^2}} \quad df = n - 2$$

$$s_X^2 = SXX / (n - 1)$$

$$s_Y^2 = SYY / (n - 1)$$

$$SYY = \sum_{i=1}^n (Y_i - \bar{Y})^2$$

$$r_{XY} = \frac{SXY / (n-1)}{s_X s_Y}$$

correlation coefficient

$$R^2 = r_{XY}^2$$

$$= \frac{SST - SSE}{SST}$$

coefficient of determination

$$SST = SYY = \sum_{i=1}^n (Y_i - \bar{Y})^2$$

How certain are we about $\hat{\beta}_0$ and $\hat{\beta}_1$ being equal to β_0 and β_1 ?

We quantify this uncertainty using their standard errors:

$$SE(\beta_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} \quad df = n - 2$$

$$SE(\beta_1) = \hat{\sigma} \sqrt{\frac{1}{(n-1)s_X^2}} \quad df = n - 2$$

$$s_X^2 = SXX / (n - 1)$$

$$s_Y^2 = SY / (n - 1)$$

$$SYY = \sum_{i=1}^n (Y_i - \bar{Y})^2$$

$$r_{XY} = \frac{SXY / (n-1)}{s_X s_Y}$$

correlation coefficient

$$R^2 = r_{XY}^2$$


$$= \frac{SST - SSE}{SST}$$

coefficient of determination

$$SST = SYY = \sum_{i=1}^n (Y_i - \bar{Y})^2$$

The coefficient of determination (R^2) is the proportion of the total response variation explained by the explanatory variable(s).

Telomere length vs years post diagnosis



Pvalues and confidence interval

We can compute two-sided pvalues via

$$2P\left(t_{n-2} < -\left|\frac{\hat{\beta}_0}{SE(\beta_0)}\right|\right) \quad \text{and} \quad 2P\left(t_{n-2} < -\left|\frac{\hat{\beta}_1}{SE(\beta_1)}\right|\right)$$

Pvalues and confidence interval

We can compute two-sided pvalues via

$$2P\left(t_{n-2} < -\left|\frac{\hat{\beta}_0}{SE(\beta_0)}\right|\right) \quad \text{and} \quad 2P\left(t_{n-2} < -\left|\frac{\hat{\beta}_1}{SE(\beta_1)}\right|\right)$$

These test the null hypothesis that the corresponding parameter is zero.

Pvalues and confidence interval

We can compute two-sided pvalues via

$$2P\left(t_{n-2} < -\left|\frac{\hat{\beta}_0}{SE(\beta_0)}\right|\right) \quad \text{and} \quad 2P\left(t_{n-2} < -\left|\frac{\hat{\beta}_1}{SE(\beta_1)}\right|\right)$$

These test the null hypothesis that the corresponding parameter is zero.

We can construct $100(1 - \alpha)\%$ two-sided confidence intervals via

$$\hat{\beta}_0 \pm t_{n-2}(1 - \alpha/2)SE(\beta_0) \quad \text{and} \quad \hat{\beta}_1 \pm t_{n-2}(1 - \alpha/2)SE(\beta_1)$$

These provide ranges of the parameters consistent with the data.

Calculations by hand

	n	Xbar	Ybar	s _X	s _Y	r _{XY}
1	39	5.59	1.22	2.935	0.1798	-0.4307

$$SXX = (n-1)s_X^2 = (39-1) \times 2.935427^2 = 327.4358$$

$$SYY = (n-1)s_Y^2 = (39-1) \times 0.1797731^2 = 1.228098$$

$$SXY = (n-1)s_X s_Y r_{XY} = (39-1) \times 2.935427 \times 0.1797731 \times -0.4306534 = -8.635897$$

$$\hat{\beta}_1 = SXY / SXX = -8.635897 / 327.4358 = -0.02637432$$

$$\hat{\beta}_0 = \bar{Y} - \hat{\beta}_1 \bar{X} = 1.220256 - (-0.02637432) \times 5.589744 = 1.367682$$

$$R^2 = r_{XY}^2 = (-0.4306534)^2 = 0.1854624$$

$$SSE = SYY(1 - R^2) = 1.228098(1 - 0.1854624) = 1.000332$$

$$\hat{\sigma}^2 = SSE / (n-2) = 1.000332 / (39-2) = 0.027036$$

$$\hat{\sigma} = \sqrt{\hat{\sigma}^2} = \sqrt{0.027036} = 0.1644263$$

$$SE(\hat{\beta}_0) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{\bar{X}^2}{(n-1)s_X^2}} = 0.1644263 \sqrt{\frac{1}{39} + \frac{5.589744^2}{327.4358}} = 0.05721115$$

$$SE(\hat{\beta}_1) = \hat{\sigma} \sqrt{\frac{1}{(n-1)s_X^2}} = 0.1644263 \sqrt{\frac{1}{327.4358}} = 0.009086742$$

$$PH_0:\beta_0=0 = 2P\left(t_{n-2} < -\left|\frac{1.367682}{0.05721115}\right|\right) = 2P(t_{37} < -23.90586) < 0.0001$$

$$PH_0:\beta_1=0 = 2P\left(t_{n-2} < -\left|\frac{-0.02637432}{0.009086742}\right|\right) = 2P(t_{37} < -2.902506) < 0.0062$$

$$CI_{95\%} \beta_0 = \hat{\beta}_0 \pm t_{n-2}(1 - \alpha/2)SE(\hat{\beta}_0) = 1.367682 \pm 2.026192 \times 0.05721115 = (1.251761, 1.483603)$$

$$CI_{95\%} \beta_1 = \hat{\beta}_1 \pm t_{n-2}(1 - \alpha/2)SE(\hat{\beta}_1) = -0.02637432 \pm 2.026192 \times 0.009086742 = (-0.044785804 - 0.007962836)$$

```
DATA t;
  INFILE 'telomeres.csv' DSD FIRSTOBS=2;
  INPUT years length;
```

```
PROC CORR DATA=t;
  VAR length;
  WITH years;
  RUN;
```

The CORR Procedure

```
1 With Variables:  years
1   Variables:    length
```

Simple Statistics

Variable	N	Mean	Std Dev	Sum	Minimum	Maximum
years	39	5.58974	2.93543	218.00000	1.00000	12.00000
length	39	1.22026	0.17977	47.59000	0.84000	1.63000

Pearson Correlation Coefficients, N = 39
 Prob > |r| under H0: Rho=0

```
length
years  -0.43065
      0.0062
```

```
PROC GLM DATA=t;
  MODEL length = years / CLPARM;
  RUN;
```

The GLM Procedure

Number of Observations Read	39
Number of Observations Used	39

Dependent Variable: length

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	1	0.22776588	0.22776588	8.42	0.0062
Error	37	1.00033156	0.02703599		
Corrected Total	38	1.22809744			

R-Square	Coeff Var	Root MSE	length Mean
0.185462	13.47473	0.164426	1.220256

Source	DF	Type I SS	Mean Square	F Value	Pr > F
years	1	0.22776588	0.22776588	8.42	0.0062

Source	DF	Type III SS	Mean Square	F Value	Pr > F
years	1	0.22776588	0.22776588	8.42	0.0062

Parameter	Estimate	Standard Error	t Value	Pr > t	95% Confidence Limits	
Intercept	1.367682067	0.05721112	23.91	<.0001	1.251761335	1.483602799
years	-0.026374315	0.00908674	-2.90	0.0062	-0.044785794	-0.007962836

Regression in R

```
m = lm(telomere.length~years, Telomeres)
with(Telomeres, cor(telomere.length,years))
```

```
[1] -0.4307
```

```
anova(m)
```

Analysis of Variance Table

Response: telomere.length

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
years	1	0.228	0.228	8.42	0.0062 **
Residuals	37	1.000	0.027		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Regression in R

```
m = lm(telomere.length~years, Telomeres)
summary(m)
```

```
Call:
lm(formula = telomere.length ~ years, data = Telomeres)
```

```
Residuals:
```

	Min	1Q	Median	3Q	Max
	-0.4222	-0.0854	0.0206	0.1074	0.2887

```
Coefficients:
```

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	1.36768	0.05721	23.9	<2e-16 ***
years	-0.02637	0.00909	-2.9	0.0062 **

```
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Residual standard error: 0.164 on 37 degrees of freedom
```

```
Multiple R-squared:  0.185, Adjusted R-squared:  0.163
```

```
F-statistic: 8.42 on 1 and 37 DF,  p-value: 0.0062
```

```
confint(m)
```

	2.5 %	97.5 %
(Intercept)	1.25176	1.483603
years	-0.04479	-0.007963

Conclusion

Telomere length at the time of diagnosis of a child's chronic illness is estimated to be 1.37 with a 95% confidence interval of (1.25, 1.48).

Conclusion

Telomere length at the time of diagnosis of a child's chronic illness is estimated to be 1.37 with a 95% confidence interval of (1.25, 1.48). For each year since diagnosis, the telomere length decreases by 0.026 with a 95% confidence interval of (0.008, 0.045) **on average**.

Conclusion

Telomere length at the time of diagnosis of a child's chronic illness is estimated to be 1.37 with a 95% confidence interval of (1.25, 1.48). For each year since diagnosis, the telomere length decreases by 0.026 with a 95% confidence interval of (0.008, 0.045) **on average**. The proportional of variability in telomere length described by years since diagnosis is 18.5%.

Conclusion

Telomere length at the time of diagnosis of a child's chronic illness is estimated to be 1.37 with a 95% confidence interval of (1.25, 1.48). For each year since diagnosis, the telomere length decreases by 0.026 with a 95% confidence interval of (0.008, 0.045) **on average**. The proportional of variability in telomere length described by years since diagnosis is 18.5%.

<http://www.pnas.org/content/101/49/17312>

The zero-order correlation between chronicity of caregiving [years] and mean telomere length, r , is -0.445 ($P < 0.01$). [$R^2 = 0.198$ was shown in the plot.]

Conclusion

Telomere length at the time of diagnosis of a child's chronic illness is estimated to be 1.37 with a 95% confidence interval of (1.25, 1.48). For each year since diagnosis, the telomere length decreases by 0.026 with a 95% confidence interval of (0.008, 0.045) **on average**. The proportional of variability in telomere length described by years since diagnosis is 18.5%.

<http://www.pnas.org/content/101/49/17312>

The zero-order correlation between chronicity of caregiving [years] and mean telomere length, r , is -0.445 ($P < 0.01$). [$R^2 = 0.198$ was shown in the plot.]

Remark I'm guessing our analysis and that reported in the paper don't match exactly due to a discrepancy in the data.

Summary

- The simple linear regression model is

$$Y_i \stackrel{ind}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

where Y_i and X_i are the response and explanatory variable, respectively, for individual i .

Summary

- The simple linear regression model is

$$Y_i \stackrel{ind}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

where Y_i and X_i are the response and explanatory variable, respectively, for individual i .

- Know how to use SAS/R to obtain $\hat{\beta}_0$, $\hat{\beta}_1$, $\hat{\sigma}^2$, R^2 , pvalues, CIs, etc.

Summary

- The simple linear regression model is

$$Y_i \stackrel{ind}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

where Y_i and X_i are the response and explanatory variable, respectively, for individual i .

- Know how to use SAS/R to obtain $\hat{\beta}_0$, $\hat{\beta}_1$, $\hat{\sigma}^2$, R^2 , pvalues, CIs, etc.
- Interpret SAS output

Summary

- The simple linear regression model is

$$Y_i \stackrel{ind}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

where Y_i and X_i are the response and explanatory variable, respectively, for individual i .

- Know how to use SAS/R to obtain $\hat{\beta}_0$, $\hat{\beta}_1$, $\hat{\sigma}^2$, R^2 , pvalues, CIs, etc.
- Interpret SAS output
 - At a value of zero for the explanatory variable ($X_i = 0$), β_0 is the expected value for the response (Y_i).

Summary

- The simple linear regression model is

$$Y_i \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

where Y_i and X_i are the response and explanatory variable, respectively, for individual i .

- Know how to use SAS/R to obtain $\hat{\beta}_0$, $\hat{\beta}_1$, $\hat{\sigma}^2$, R^2 , pvalues, CIs, etc.
- Interpret SAS output
 - At a value of zero for the explanatory variable ($X_i = 0$), β_0 is the expected value for the response (Y_i).
 - For each unit increase in the explanatory variable value, β_1 is the expected increase in the response.

Summary

- The simple linear regression model is

$$Y_i \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

where Y_i and X_i are the response and explanatory variable, respectively, for individual i .

- Know how to use SAS/R to obtain $\hat{\beta}_0$, $\hat{\beta}_1$, $\hat{\sigma}^2$, R^2 , pvalues, CIs, etc.
- Interpret SAS output
 - At a value of zero for the explanatory variable ($X_i = 0$), β_0 is the expected value for the response (Y_i).
 - For each unit increase in the explanatory variable value, β_1 is the expected increase in the response.
 - At a constant value of the explanatory variable, σ^2 is the variance of the responses.

Summary

- The **simple linear regression** model is

$$Y_i \stackrel{\text{ind}}{\sim} N(\beta_0 + \beta_1 X_i, \sigma^2)$$

where Y_i and X_i are the response and explanatory variable, respectively, for individual i .

- Know how to use SAS/R to obtain $\hat{\beta}_0$, $\hat{\beta}_1$, $\hat{\sigma}^2$, R^2 , pvalues, CIs, etc.
- Interpret SAS output
 - At a value of zero for the explanatory variable ($X_i = 0$), β_0 is the expected value for the response (Y_i).
 - For each unit increase in the explanatory variable value, β_1 is the expected increase in the response.
 - At a constant value of the explanatory variable, σ^2 is the variance of the responses.
 - The coefficient of determination (R^2) is the percentage of the total response variation explained by the explanatory variable(s).

What is $E[Y|X = x]$?

We know $\beta_0 = E[Y|X = 0]$, but what about $X = x$?

What is $E[Y|X = x]$?

We know $\beta_0 = E[Y|X = 0]$, but what about $X = x$?

$$E[Y|X = x] = \beta_0 + \beta_1 x$$

What is $E[Y|X = x]$?

We know $\beta_0 = E[Y|X = 0]$, but what about $X = x$?

$$E[Y|X = x] = \beta_0 + \beta_1 x$$

which we can estimate via

$$E[\widehat{Y}|X = x] = \hat{\beta}_0 + \hat{\beta}_1 x$$

What is $E[Y|X = x]$?

We know $\beta_0 = E[Y|X = 0]$, but what about $X = x$?

$$E[Y|X = x] = \beta_0 + \beta_1 x$$

which we can estimate via

$$E[\widehat{Y}|X = x] = \hat{\beta}_0 + \hat{\beta}_1 x$$

but there is uncertainty in both β_0 and β_1 .

What is $E[Y|X = x]$?

We know $\beta_0 = E[Y|X = 0]$, but what about $X = x$?

$$E[Y|X = x] = \beta_0 + \beta_1 x$$

which we can estimate via

$$E[\widehat{Y|X = x}] = \hat{\beta}_0 + \hat{\beta}_1 x$$

but there is uncertainty in both β_0 and β_1 . So the standard error of $E[Y|X = x]$ is

$$SE(E[Y|X = x]) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{(\bar{X} - x)^2}{(n-1)s_X^2}}$$

and a $100(1 - \alpha)\%$ confidence interval is

$$\hat{\beta}_0 + \hat{\beta}_1 x \pm t_{n-2}(1 - \alpha/2)SE(E[Y|X = x])$$

What do we predict about Y at $X = x$?

On the last slide, we calculated $E[Y|X = x]$ and it's uncertainty

What do we predict about Y at $X = x$?

On the last slide, we calculated $E[Y|X = x]$ and it's uncertainty, but if we are trying to predict a new observation, we need to account for the sampling variability σ^2 .

What do we predict about Y at $X = x$?

On the last slide, we calculated $E[Y|X = x]$ and it's uncertainty, but if we are trying to predict a new observation, we need to account for the sampling variability σ^2 . Thus a prediction about Y at a new $X = x$ is still

$$\text{Pred}\{Y|X = x\} = \hat{\beta}_0 + \hat{\beta}_1 x$$

but the uncertainty includes the variability due to σ^2 .

What do we predict about Y at $X = x$?

On the last slide, we calculated $E[Y|X = x]$ and it's uncertainty, but if we are trying to predict a new observation, we need to account for the sampling variability σ^2 . Thus a prediction about Y at a new $X = x$ is still

$$\text{Pred}\{Y|X = x\} = \hat{\beta}_0 + \hat{\beta}_1 x$$

but the uncertainty includes the variability due to σ^2 . So the standard error of $\text{Pred}\{Y|X = x\}$ is

$$SE(\text{Pred}\{Y|X = x\}) = \hat{\sigma} \sqrt{1 + \frac{1}{n} + \frac{(\bar{X} - x)^2}{(n-1)s_X^2}}$$

and a $100(1 - \alpha)\%$ confidence interval is

$$\hat{\beta}_0 + \hat{\beta}_1 x \pm t_{n-2}(1 - \alpha/2)SE(\text{Pred}\{Y|X = x\}).$$

```
DATA tnew;
  INPUT years;
  DATALINES;
4
;
```

```
DATA combined;
  SET t tnew;
  RUN;
```

```
PROC PRINT DATA=combined;
  WHERE years=4;
  RUN;
```

Obs	years	length
10	4	1.51
11	4	1.31
15	4	1.03
16	4	0.84
40	4	.

```
PROC GLM DATA=combined;
  MODEL length = years;
  OUTPUT OUT=combinedreg PREDICTED=predicted  LCLM=lclm UCLM=uclm LCL=lcl UCL=ucl;
  RUN;
```

```
PROC PRINT DATA=combinedreg;
  WHERE length=.;          /* . is missing data in SAS */
  RUN;
```


Obs	years	length	predicted	lclm	uclm	lcl	ucl
40	4	.	1.26218	1.20133	1.32303	0.92351	1.60086


```
m = lm(telomere.length~years, Telomeres)
new = data.frame(years=4)
predict(m, new, interval="confidence")
```

```
      fit    lwr    upr
1 1.262 1.201 1.323
```

```
predict(m, new, interval="prediction")
```

```
      fit    lwr    upr
1 1.262 0.9235 1.601
```



Shifting the intercept

The intercept (β_0) is the expected response when the explanatory variable is zero.

Shifting the intercept

The intercept (β_0) is the expected response when the explanatory variable is zero.

So, if we change our explanatory variable, we change the interpretation of our intercept

Shifting the intercept

The intercept (β_0) is the expected response when the explanatory variable is zero.

So, if we change our explanatory variable, we change the interpretation of our intercept, e.g. if, instead of using number of years since diagnosis, we use “number of years since diagnosis **minus 4**”, then our intercept is the expected response at 4 years since diagnosis.

Shifting the intercept

The intercept (β_0) is the expected response when the explanatory variable is zero.

So, if we change our explanatory variable, we change the interpretation of our intercept, e.g. if, instead of using number of years since diagnosis, we use “number of years since diagnosis **minus 4**”, then our intercept is the expected response at 4 years since diagnosis.

Let x be number of years since diagnosis, then

$$E[Y|X = x] = \tilde{\beta}_0 + \tilde{\beta}_1(x - 4) = (\beta_0 - 4\beta_1) + \beta_1x$$

so our new parameters for the mean are

- intercept $\tilde{\beta}_0 = (\beta_0 - 4\beta_1)$

Shifting the intercept

The intercept (β_0) is the expected response when the explanatory variable is zero.

So, if we change our explanatory variable, we change the interpretation of our intercept, e.g. if, instead of using number of years since diagnosis, we use “number of years since diagnosis **minus 4**”, then our intercept is the expected response at 4 years since diagnosis.

Let x be number of years since diagnosis, then

$$E[Y|X = x] = \tilde{\beta}_0 + \tilde{\beta}_1(x - 4) = (\beta_0 - 4\beta_1) + \beta_1 x$$

so our new parameters for the mean are

- intercept $\tilde{\beta}_0 = (\beta_0 - 4\beta_1)$ and
- slope $\tilde{\beta}_1 = \beta_1$ (unchanged).

```
DATA t;
  INFILE "telomeres.csv" DSD FIRSTOBS=2;
  INPUT years length;
  years4 = years-4;
```

```
PROC GLM DATA=combined;
  MODEL length = years;
  RUN;
```

```
PROC GLM DATA=combined;
  MODEL length = years4;
  RUN;
```

Parameter	Estimate	Standard Error	t Value	Pr > t	95% Confidence Limits	
Intercept	1.367682067	0.05721112	23.91	<.0001	1.251761335	1.483602799
years	-0.026374315	0.00908674	-2.90	0.0062	-0.044785794	-0.007962836
Intercept	1.262184808	0.03003174	42.03	<.0001	1.201334726	1.323034890
years4	-0.026374315	0.00908674	-2.90	0.0062	-0.044785794	-0.007962836


```
m0 = lm(telomere.length ~ years, Telomeres)
m4 = lm(telomere.length ~ I(years-4), Telomeres)
```

```
coef(m0)
```

```
(Intercept)      years
   1.36768      -0.02637
```

```
coef(m4)
```

```
(Intercept) I(years - 4)
   1.26218      -0.02637
```

```
confint(m0)
```

```
              2.5 %    97.5 %
(Intercept)  1.25176  1.483603
years        -0.04479 -0.007963
```

```
confint(m4)
```

```
              2.5 %    97.5 %
(Intercept)  1.20133  1.323035
I(years - 4) -0.04479 -0.007963
```