

Introduction to Bayesian Computation

Dr. Jarad Niemi

Iowa State University

March 21, 2016

Bayesian computation

Goals:

- $E_{\theta|y}[h(\theta)|y] = \int h(\theta)p(\theta|y)d\theta$
- $p(y) = \int p(y|\theta)p(\theta)d\theta = E_{\theta}[p(y|\theta)]$

Approaches:

- Deterministic approximation
- Monte Carlo approximation
 - Theoretical justification
 - Gridding
 - Inverse CDF
 - Accept-reject

Example: Normal-Cauchy model

Let $Y \sim N(\theta, 1)$ with $\theta \sim \text{Ca}(0, 1)$. The posterior is

$$p(\theta|y) \propto p(y|\theta)p(\theta) \propto \frac{\exp(-(y - \theta)^2/2)}{1 + \theta^2}$$

which is not a known distribution. We might be interested in

1. normalizing this posterior, i.e. calculating

$$p(y) = \int p(y|\theta)p(\theta)d\theta$$

2. or in calculating the posterior mean, i.e.

$$E[\theta|y] = \int \theta p(\theta|y)d\theta.$$

Numerical integration

We have

$$E[h(\theta)|y] = \int h(\theta)p(\theta|y)d\theta \approx \sum_{s=1}^S w_s h\left(\theta^{(s)}\right) p\left(\theta^{(s)}|y\right)$$

Approaches:

- deterministic methods where
 - $\theta^{(s)}$ are selected points,
 - w_s is the weight given to the point $\theta^{(s)}$, and
 - the error can be bounded.
- Monte Carlo (simulation) methods where
 - $\theta^{(s)} \stackrel{iid}{\sim} g(\theta)$ (for some proposal distribution g),
 - $w_s = p(\theta|y)/g(\theta)$,
 - and we have SLLN and CLT.

Marginal likelihood

```

y = 1 # Data

q = function(theta,y,log=FALSE) {
  out = -(y-theta)^2/2-log(1+theta^2)
  if (log) return(out)
  return(exp(out))
}

# Find marginal likelihood for y
w = 0.1
theta = seq(-5,5,by=w)+y
(py = sum(q(theta,y)*w))

[1] 1.305608


integrate(function(x) q(x,y), -Inf, Inf)

1.305609 with absolute error < 0.00013

```

Example: Normal-Cauchy model

```
curve(q(x,y), -5, 5, n=1001)
points(theta,rep(0,length(theta)), cex=0.5, pch=19)
segments(theta,0,theta,q(theta,y))
```




Posterior expectation

$$E[h(\theta)|y] \approx \sum_{s=1}^S w_s h\left(\theta^{(s)}\right) p\left(\theta^{(s)}|y\right) = \sum_{s=1}^S w_s h\left(\theta^{(s)}\right) \frac{p\left(\theta^{(s)}|y\right)}{p(y)}$$

```
h = function(theta) theta
sum(w*h(theta)*q(theta,y)/py)
```

```
[1] 0.5542021
```

Posterior expectation as a function of observed data



Monte Carlo integration

Consider evaluating the integral

$$E[h(\theta)] = \int_{\Theta} h(\theta)p(\theta)d\theta$$

using the Monte Carlo estimate

$$\hat{h}_J = \frac{1}{J} \sum_{j=1}^J h(\theta^{(j)})$$

where $\theta^{(j)} \stackrel{\text{ind}}{\sim} g(\theta)$. We know

- SLLN: \hat{h}_J converges almost surely to $E[h(\theta)]$.
- CLT: if h^2 has finite expectation, then

$$\hat{h}_J \xrightarrow{d} N(E[h(\theta)], v_J)$$

where

$$v_J = \frac{1}{J} \widehat{V[h(\theta)]} \approx \frac{1}{J^2} \sum_{s=1}^J \left[h(\theta^{(s)}) - \hat{h}_J \right]^2.$$

Definite integral

Suppose you are interested in evaluating

$$I = \int_0^1 e^{-\theta^2/2} d\theta.$$

Then set

- $h(\theta) = e^{-\theta^2/2}$ and
- $p(\theta) = 1$, i.e. $\theta \sim \text{Unif}(0, 1)$.



and approximate by a Monte Carlo estimate via

1. For $j = 1, \dots, J$,
 - a. sample $\theta^{(j)} \sim \text{Unif}(0, 1)$ and
 - b. calculate $h(\theta^{(j)})$.

2. Calculate


$$I \approx \frac{1}{J} \sum_{j=1}^J h(\theta^{(j)}).$$

Strong law of large numbers

20 samples**200 samples**


Strong law of large numbers

Monte Carlo estimate



Central limit theorem

Monte Carlo estimate



Infinite bounds

Suppose $\theta \sim N(0, 1)$ and you are interested in evaluating

$$E[\theta] = \int_{-\infty}^{\infty} \theta \frac{1}{\sqrt{2\pi}} e^{-\theta^2/2} d\theta$$

Then set



- $h(\theta) = \theta$ and
- $g(\theta) = \phi(\theta)$, i.e. $\theta \sim N(0, 1)$.

and approximate by a Monte Carlo estimate via


1. For $j = 1, \dots, J$,
 - a. sample $\theta^{(j)} \sim N(0, 1)$ and
 - b. calculate $h(\theta^{(j)})$.
2. Calculate

$$E[\theta] \approx \frac{1}{J} \sum_{j=1}^J h(\theta^{(j)}).$$

Non-uniform sampling



Monte Carlo estimate



Monte Carlo approximation via gridding

Rather than approximate $p(y)$ and then $E[\theta|y]$ via deterministic gridding (all w_i are equal), we can use the grid as a discrete approximation to the posterior, i.e.

$$p(\theta|y) \approx \sum_{i=1}^N p_i \delta_{\theta_i}(\theta) \quad p_i = \frac{q(\theta_i|y)}{\sum_{j=1}^N q(\theta_j|y)}$$

where $\delta_{\theta_i}(\theta)$ is the Dirac delta function, i.e.

$$\delta_{\theta_i}(\theta) = 0 \quad \forall \theta \neq \theta_i \quad \int \delta_{\theta_i}(\theta) d\theta = 1.$$

This discrete approximation to $p(\theta|y)$ can be used to approximate the expectation $E[h(\theta)|y]$ deterministically or via simulation, i.e.

$$E[h(\theta)|y] \approx \sum_{i=1}^N p_i h(\theta_i) \quad E[h(\theta)|y] \approx \frac{1}{S} \sum_{s=1}^S h(\theta^{(s)})$$

where $\theta^{(s)} \sim \sum_{i=1}^N p_i \delta_{\theta_i}(\theta)$ (with replacement).

Example: Normal-Cauchy model

```
# Small number of grid locations  
theta = seq(-5,5,length=1e2+1)+y; p = q(theta,y)/sum(q(theta,y)); sum(p*theta)
```

```
[1] 0.5542021
```

```
mean(sample(theta,prob=p,replace=TRUE))
```

```
[1] 0.6158416
```

```
# Large number of grid locations  
theta = seq(-5,5,length=1e6+1)+y; p = q(theta,y)/sum(q(theta,y)); sum(p*theta)
```

```
[1] 0.5542021
```

```
mean(sample(theta,1e2,prob=p,replace=TRUE)) # But small MC sample
```

```
[1] 0.5819513
```

```
# Truth  
post_expectation(1)
```

```
[1] 0.5542021
```

Inverse cumulative distribution function

Definition

The **cumulative distribution function** of a random variable X is defined by

$$F_X(x) = P_X(X \leq x) \quad \text{for all } x.$$


Suppose you want to sample $X \sim f(x)$ and you have access to the inverse cdf of X , $F^{-1}(x)$, then

Lemma

If $U \sim \text{Unif}(0, 1)$, then $X = F^{-1}(U)$ is a simulation from $f(x)$.

Inverse CDF


Standard normal CDF



Exponential example

For example, to sample $X \sim \text{Exp}(1)$,

1. Sample $U \sim \text{Unif}(0, 1)$.
2. Set $X = -\log(1 - U)$, or $X = -\log(U)$.



Sampling from a univariate truncated distribution


Suppose you wish to sample from $X \sim N(\mu, \sigma^2)I(a < X < b)$, i.e. a normal random variable with untruncated mean μ and variance σ^2 , but truncated to the interval (a, b) . Suppose the untruncated cdf is F and inverse cdf is F^{-1} .

1. Calculate endpoints $p_a = F(a)$ and $p_b = F(b)$.
2. Sample $U \sim \text{Unif}(p_a, p_b)$.
3. Set $X = F^{-1}(U)$.


This just avoids having to recalculate the normalizing constant for the pdf, i.e. $1/(F^{-1}(b) - F^{-1}(a))$, which rescales

Normal

$$X \sim N(5, 9)I(1 \leq X \leq 6)$$



$$X \sim N(5, 9)I(1 \leq X \leq 6)$$



Rejection sampling

Suppose you wish to obtain samples $\theta \sim p(\theta|y)$, rejection sampling performs the following

1. Sample a proposal $\theta^* \sim g(\theta)$ and $U \sim \text{Unif}(0, 1)$.
2. Accept $\theta = \theta^*$ as a draw from $p(\theta|y)$ if $U \leq p(\theta^*|y)/Mg(\theta^*)$, otherwise return to step 1.

where M satisfies $Mg(\theta) \geq p(\theta|y)$ for all θ .

- For a given proposal distribution $g(\theta)$, the optimal M is $M = \sup_{\theta} p(\theta|y)/g(\theta)$.
- The probability of acceptance is $1/M$.

The accept-reject idea is to create an envelope, $Mg(\theta)$, above $p(\theta|y)$.

Rejection sampling with unnormalized density

Suppose you wish to obtain samples $\theta \sim p(\theta|y) \propto q(\theta|y)$, rejection sampling performs the following

1. Sample a proposal $\theta^* \sim g(\theta)$ and $U \sim \text{Unif}(0, 1)$.
2. Accept $\theta = \theta^*$ as a draw from $p(\theta|y)$ if $U \leq q(\theta^*|y)/M^*g(\theta^*)$, otherwise return to step 1.

where M^* satisfies $M^* g(\theta) \geq q(\theta|y)$ for all θ .

- For a given proposal distribution $g(\theta)$, the optimal M^* is $M^* = \sup_{\theta} q(\theta|y)/g(\theta)$.
- The acceptance probability is $1/M = p(y)/M^*$.

The accept-reject idea is to create an envelope, $M g(\theta)$, above $q(\theta|y)$.

Example: Normal-Cauchy model

If $Y \sim N(\theta, 1)$ and $\theta \sim Ca(0, 1)$, then

$$p(\theta|y) \propto e^{-(y-\theta)^2/2} \frac{1}{(1+\theta^2)}$$

for $\theta \in \mathbb{R}$.

Choose a $N(y, 1)$ as a proposal distribution, i.e.


$$g(\theta) = \frac{1}{\sqrt{2\pi}} e^{-(\theta-y)^2/2}$$

with

$$M^* = \sup_{\theta} \frac{q(\theta|y)}{g(\theta)} = \sup_{\theta} \frac{e^{-(y-\theta)^2/2} \frac{1}{(1+\theta^2)}}{\frac{1}{\sqrt{2\pi}} e^{-(\theta-y)^2/2}} = \frac{\sqrt{2\pi}}{(1+\theta^2)} \leq \sqrt{2\pi}$$

The acceptance rate is $1/M = p(y)/M^* = 1.643545/\sqrt{2\pi} = 0.656$.

Example: Normal-Cauchy model



Acceptance rate was 0.66

Heavy-tailed proposals

Suppose our target is a standard Cauchy and our (proposed) proposal is a standard normal, then

$$\frac{p(\theta|y)}{g(\theta)} = \frac{\frac{1}{\pi(1+\theta^2)}}{\frac{1}{\sqrt{2\pi}}e^{-\theta^2/2}}$$

and

$$\frac{\frac{1}{\pi(1+\theta^2)}}{\frac{1}{\sqrt{2\pi}}e^{-\theta^2/2}} \xrightarrow{\theta \rightarrow \infty} \infty$$

since e^{-a} converges to zero faster than $1/(1+a)$. Thus, there is no value M such that $M g(\theta) \geq p(\theta|y)$ for all θ .

Bottom line: the condition $M g(\theta) \geq p(\theta|y)$ requires the proposal to have tails at least as thick (heavy) as the target.