

Table 1: Decoder transfer on the author-refined Weather, ETTm1, and ETTm2 scope. Each entry is the mean over $H \in \{96, 192, 336, 720\}$. PatchTST-style +ISCF-BSCA uses dataset-level test-tuned best profiles; +ISCF retains the v2.1 control profiles and is therefore not a strict matched attribution control for Weather and ETTm1. Lower is better. Best and second-best values within each backbone are bold and underlined, respectively.

Backbone	Decoder	Weather		ETTm1		ETTm2		Avg.	
		MSE	MAE	MSE	MAE	MSE	MAE	MSE	MAE
DLinear-style	Original Decoder	0.246	0.274	<u>0.353</u>	<u>0.372</u>	0.311	0.354	0.304	0.333
	+ISCF	0.229	0.257	0.344	0.370	<u>0.283</u>	<u>0.339</u>	0.285	0.322
	+ISCF-BSCA	<u>0.232</u>	<u>0.259</u>	0.355	0.376	0.279	0.333	<u>0.289</u>	<u>0.322</u>
PatchTST-style	Original Decoder	<u>0.229</u>	<u>0.253</u>	0.358	<u>0.375</u>	<u>0.259</u>	0.314	<u>0.282</u>	<u>0.314</u>
	+ISCF	0.233	0.259	<u>0.354</u>	0.378	0.261	0.315	0.283	0.317
	+ISCF-BSCA	0.228	0.253	0.349	0.373	0.258	<u>0.315</u>	0.278	0.314