

CS7GV2: Mathematics of Light and Sound, M.Sc. in Computer Science.

Lecture #6: Random Phasor Sums

Fergal Shevlin, Ph.D.

School of Computer Science and Statistics,
Trinity College Dublin


November 10, 2022

Random walks


Multiple wave phasors can be summed like vectors to find the *resultant* wave phasor at a point in space and time.

When wave amplitudes are independent and wave phases are independent* their summation is called a “random walk.”


Random walks



Random walks




Random walks




Random walks

Multiple wave phasors can be summed like vectors to find the *resultant* wave phasor at a point in space and time.


When wave amplitudes are independent and wave phases are independent* their summation is called a “random walk.”



Multiple phasors and their resultant sum.



Same amplitudes, different phases, smaller resultant.



Independence and randomness

Independence means that one event or value of a random quantity X (e.g. a wave's amplitude) has no effect on another, Y (e.g. its phase,)

$$\mathbf{P}(X|Y) = \mathbf{P}(X \cap Y)/\mathbf{P}(Y)$$

$$\mathbf{P}(X \cap Y) = \mathbf{P}(X|Y)\mathbf{P}(Y)$$

$$\mathbf{P}(X \cap Y) = \mathbf{P}(X)\mathbf{P}(Y) \text{ i.f.f.}$$

$$\mathbf{P}(X) = \mathbf{P}(X|Y) \text{ and } \mathbf{P}(Y) = \mathbf{P}(Y|X).$$


Independence and randomness

A random quantity is one whose value depends on the outcome of a random phenomenon.


Its occurrence may be known* to follow a particular probability density function f_X , or probability mass function p_X , with descriptive parameters μ, σ , etc.

Example PDFs are *Uniform, Normal, Exponential, Poisson, Rayleigh*.

Independence and randomness



Independence and randomness



Independence and randomness

Independence means that one event or value of a random quantity X (e.g. a wave's amplitude) has no effect on another, Y (e.g. its phase,)

$$P(X|Y) = P(X \cap Y)/P(Y)$$

$$P(X \cap Y) = P(X|Y)P(Y)$$



$$P(X \cap Y) = P(X)P(Y) \text{ i.f.f.}$$

$$P(X) = P(X|Y) \text{ and } P(Y) = P(Y|X).$$


A random quantity is one whose value depends on the outcome of a random phenomenon.

Its occurrence may be known* to follow a particular probability density function f_X , or probability mass function p_X , with descriptive parameters μ, σ , etc.


Example PDFs are *Uniform, Normal, Exponential, Poisson, Rayleigh*.




10,000 values (x, y) chosen randomly




10,000 values (x, y) chosen randomly







10,000 values (x, y) chosen randomly



10,000 values (x, y) chosen randomly



10,000 values (x, y) chosen randomly



Descriptive statistics

Expected value or *mean* of a continuous random quantity X with probability density function f_X ,

$$\mathbf{E}[X] = \int_{-\infty}^{+\infty} x f_X(x) dx.$$

And for a discrete, finite, random quantity X with probability mass function p_X ,

$$\mathbf{E}[X] = \sum_{i=1}^N x_i p_X(x_i).$$

Descriptive statistics

This is the *arithmetic mean* when probability mass function p_X is uniformly $1/N$.

$$\begin{aligned}\mathbf{E}[X] &= \sum_{i=1}^N x_i \frac{1}{N} \\ &= \frac{1}{N} \sum_{i=1}^N x_i \\ &= (x_1 + x_2 + \dots + x_N) / N\end{aligned}$$

Descriptive statistics

Linearity of expectation,

$$\mathbf{E}[X + Y] = \mathbf{E}[X] + \mathbf{E}[Y].$$

If $Y = aX + b$ for $a, b \in \mathbb{R}$,

$$\mathbf{E}[Y] = a\mathbf{E}[X] + b.$$

If X, Y independent,

$$\mathbf{E}[XY] = \mathbf{E}[X]\mathbf{E}[Y].$$

Descriptive statistics

Variance is mean distance squared to the mean
(when uniform,)

$$\begin{aligned}\sigma_X^2 &= \mathbf{E}[(X - \mathbf{E}[X])^2] \\ &= \mathbf{E}[X^2] - \mathbf{E}[X]^2 \\ &= \mathbf{E}[X^2] \text{ when } \mathbf{E}[X] = 0.\end{aligned}$$

Standard deviation $\sigma_X = \sqrt{\sigma_X^2}$.

[cf. SCFT STD vs MAD.]

Descriptive statistics

Expected value or *mean* of a continuous random quantity X with probability density function f_X ,

$$\mathbf{E}[X] = \int_{-\infty}^{+\infty} x f_X(x) dx.$$

And for a discrete, finite, random quantity X with probability mass function p_X ,

$$\mathbf{E}[X] = \sum_{i=1}^N x_i p_X(x_i).$$

This is the *arithmetic mean* when probability mass function p_X is uniformly $1/N$.

$$\begin{aligned}\mathbf{E}[X] &= \sum_{i=1}^N x_i 1/N \\ &= 1/N \sum_{i=1}^N x_i \\ &= (x_1 + x_2 + \dots + x_N)/N\end{aligned}$$

Linearity of expectation,

$$\mathbf{E}[X + Y] = \mathbf{E}[X] + \mathbf{E}[Y].$$

If $Y = aX + b$ for $a, b \in \mathbb{R}$,

$$\mathbf{E}[Y] = a\mathbf{E}[X] + b.$$

If X, Y independent,

$$\mathbf{E}[XY] = \mathbf{E}[X]\mathbf{E}[Y].$$

Variance is mean distance squared to the mean (when uniform,)

$$\begin{aligned}\sigma_X^2 &= \mathbf{E}[(X - \mathbf{E}[X])^2] \\ &= \mathbf{E}[X^2] - \mathbf{E}[X]^2 \\ &= \mathbf{E}[X^2] \text{ when } \mathbf{E}[X] = 0.\end{aligned}$$

Standard deviation $\sigma_X = \sqrt{\sigma_X^2}$.

[cf. SCFT STD vs MAD.]

Random phasor sum

Defined as a weighted sum of random phasors:

$$\begin{aligned}\frac{1}{\sqrt{N}} \sum_{n=1}^N a_n e^{j\phi_n} &= \frac{1}{\sqrt{N}} \sum_{n=1}^N \mathbf{a}_n = A e^{j\theta} \\ &= \mathbf{A} \quad (\text{the "resultant."})\end{aligned}$$

Random phasor sum

$$\begin{aligned}\mathbf{E}[\operatorname{Re}\{\mathbf{A}\}] &= \mathbf{E}\left[\frac{1}{\sqrt{N}} \sum_{n=1}^N a_n \cos \phi_n\right] \\ &= \frac{1}{\sqrt{N}} \sum \mathbf{E}[a_n \cos \phi_n] \\ &= \frac{1}{\sqrt{N}} \sum \mathbf{E}[a_n] \mathbf{E}[\cos \phi_n] \\ &= 0.\end{aligned}$$

Similarly, $\mathbf{E}[\operatorname{Im}\{\mathbf{A}\}] = 0$.

Random phasor sum

$$\sigma_{\text{Re}\{\mathbf{A}\}}^2 = \mathbf{E}[\text{Re}\{\mathbf{A}\}^2].$$

$$\begin{aligned}\text{Re}\{\mathbf{A}\}^2 &= 1/\sqrt{N}(a_1 \cos \phi_1 + a_2 \cos \phi_2 + \dots) \times \\ &\quad 1/\sqrt{N}(a_1 \cos \phi_1 + a_2 \cos \phi_2 + \dots) \\ &= 1/N \sum_n \sum_m a_n a_m \cos \phi_n \cos \phi_m.\end{aligned}$$

$$\begin{aligned}\mathbf{E}[\text{Re}\{\mathbf{A}\}^2] &= 1/N \sum_n \sum_m \mathbf{E}[a_n a_m] \times \\ &\quad \mathbf{E}[\cos \phi_n \cos \phi_m] \\ &= 1/N \sum_n \mathbf{E}[a_n^2] \mathbf{E}[\cos^2 \phi_n]\end{aligned}$$

Random phasor sum

$$\begin{aligned} & \text{(since for } n \neq m, \mathbf{E}[\cos \phi_n \cos \phi_m] \\ & \quad = \mathbf{E}[\cos \phi_n] \mathbf{E}[\cos \phi_m] = 0) \\ & = 1/N \sum_n \mathbf{E}[a_n^2] \times \mathbf{E}[1/2 + 1/2 \cos 2\phi_n] \\ & \quad \text{(since } \cos^2 \phi = (1 + \cos 2\phi)/2) \\ & = 1/N \sum_n \mathbf{E}[a_n^2]/2. \end{aligned}$$

$$\text{Similarly, } \sigma_{\text{Im}\{\mathbf{A}\}}^2 = 1/N \sum_n \mathbf{E}[a_n^2]/2.$$

Random phasor sum

Defined as a weighted sum of random phasors:

$$\begin{aligned}\frac{1}{\sqrt{N}} \sum_{n=1}^N a_n e^{j\phi_n} &= \frac{1}{\sqrt{N}} \sum_{n=1}^N \mathbf{a}_n = A e^{j\theta} \\ &= \mathbf{A} \quad (\text{the "resultant."})\end{aligned}$$

$$\begin{aligned}\mathbf{E}[\operatorname{Re}\{\mathbf{A}\}] &= \mathbf{E}\left[\frac{1}{\sqrt{N}} \sum_{n=1}^N a_n \cos \phi_n\right] \\ &= \frac{1}{\sqrt{N}} \sum \mathbf{E}[a_n \cos \phi_n] \\ &= \frac{1}{\sqrt{N}} \sum \mathbf{E}[a_n] \mathbf{E}[\cos \phi_n] \\ &= 0.\end{aligned}$$

Similarly, $\mathbf{E}[\operatorname{Im}\{\mathbf{A}\}] = 0$.

$$\begin{aligned}\sigma_{\operatorname{Re}\{\mathbf{A}\}}^2 &= \mathbf{E}[\operatorname{Re}\{\mathbf{A}\}^2]. \\ \operatorname{Re}\{\mathbf{A}\}^2 &= \frac{1}{\sqrt{N}}(a_1 \cos \phi_1 + a_2 \cos \phi_2 + \dots) \times \\ &\quad \frac{1}{\sqrt{N}}(a_1 \cos \phi_1 + a_2 \cos \phi_2 + \dots) \\ &= \frac{1}{N} \sum_n \sum_m a_n a_m \cos \phi_n \cos \phi_m. \\ \mathbf{E}[\operatorname{Re}\{\mathbf{A}\}^2] &= \frac{1}{N} \sum_n \sum_m \mathbf{E}[a_n a_m] \times \\ &\quad \mathbf{E}[\cos \phi_n \cos \phi_m] \\ &= \frac{1}{N} \sum_n \mathbf{E}[a_n^2] \mathbf{E}[\cos^2 \phi_n]\end{aligned}$$

$$\begin{aligned}&(\text{since for } n \neq m, \mathbf{E}[\cos \phi_n \cos \phi_m] \\ &\quad = \mathbf{E}[\cos \phi_n] \mathbf{E}[\cos \phi_m] = 0) \\ &= \frac{1}{N} \sum_n \mathbf{E}[a_n^2] \times \mathbf{E}[1/2 + 1/2 \cos 2\phi_n] \\ &\quad (\text{since } \cos^2 \phi = (1 + \cos 2\phi)/2) \\ &= \frac{1}{N} \sum_n \mathbf{E}[a_n^2]/2.\end{aligned}$$

Similarly, $\sigma_{\operatorname{Im}\{\mathbf{A}\}}^2 = \frac{1}{N} \sum_n \mathbf{E}[a_n^2]/2$.


Large numbers

Central Limit Theorem says that the probability density of the *sum* of N independent, identically-distributed, random quantities approaches Normal as $N \rightarrow \infty$, ,

$$f_{R,I}(r, i) = \frac{1}{2\pi\sigma^2} \exp\left\{-\frac{r^2 + i^2}{2\sigma^2}\right\}$$

Where $R = \text{Re}\{\mathbf{A}\}$ and $I = \text{Im}\{\mathbf{A}\}$ and $\sigma^2 = \sigma_R^2 = \sigma_I^2$.
[cf. SCFT p. 125.]

Large numbers



Normal $f_{R,I}(r, i), [0, 1]$.

Large numbers


Through transformation of variables,^{*} marginal statistics for A and θ are found as Rayleigh and Uniform respectively,

$$f_A(a) = a/\sigma^2 \exp\left\{\frac{a^2}{2\sigma^2}\right\}$$

$$f_\theta(\phi) = 1/2\pi$$

$$\mathbf{E}[A] = \sqrt{\pi/2} \sigma, \quad \sigma_A = (2 - \pi/2)\sigma^2$$

Large numbers




Rayleigh $f_A(a)$, $[0, 1]$ (scaled by σ_A .)

Large numbers

Central Limit Theorem says that the probability density of the *sum* of N independent, identically-distributed, random quantities approaches Normal as $N \rightarrow \infty$, ,

$$f_{R,I}(r, i) = \frac{1}{2\pi\sigma^2} \exp\left\{-\frac{r^2 + i^2}{2\sigma^2}\right\}$$

Where $R = \text{Re}\{\mathbf{A}\}$ and $I = \text{Im}\{\mathbf{A}\}$ and $\sigma^2 = \sigma_R^2 = \sigma_I^2$.
[cf. SCFT p. 125.]




Normal $f_{R,I}(r, i)$, $[0, 1]$.

Through transformation of variables,* marginal statistics for A and θ are found as Rayleigh and Uniform respectively,

$$f_A(a) = a/\sigma^2 \exp\left\{-\frac{a^2}{2\sigma^2}\right\}$$

$$f_\theta(\phi) = 1/2\pi$$

$$\mathbf{E}[A] = \sqrt{\pi/2} \sigma, \quad \sigma_A = (2 - \pi/2)\sigma^2$$



Rayleigh $f_A(a)$, $[0, 1]$ (scaled by σ_A .)