

CS7GV2: Mathematics of Light and Sound, M.Sc. in Computer Science.

5 ECTS \approx 125 hours!
 $125h / (11 + 2 + 1)w$

Lecture #1: Waves

9h/w

11 assignments
 3 hour exam questions
 based on assignments

Fergal Shevlin, Ph.D.

School of Computer Science and Statistics,
 Trinity College Dublin

September 16, 2022

2 hours Lectures
 3 hours follow-up

+ 4 hours Programming
9 hours

SciPy ~ Python
~~PLOTTING~~

①

CS7GV2: Mathematics of Light and Sound,
 M.Sc. in Computer Science.

ANACONDA
SciPy

Install on your own
 PC.

②

Lecture #1: Waves

gitlab.scss.tcd.ie

CS7GV2-2022-23
 Lecture - Notes

Ⓐ pdfs

Ⓑ SciPy
 code

Fergal Shevlin, Ph.D.



School of Computer Science and Statistics,
 Trinity College Dublin

September 16, 2022


③
 make a project
 for each
 assignment

→ add me as a "reporter"

Physical waves




A_0



$$\frac{2\pi r}{\lambda}$$

$$r^{-1} = \frac{1}{r}$$




$$\frac{A_0}{2\pi r} = A_0 \times \frac{1}{2\pi r} \propto A_0 r^{-1}$$

Physical waves

Lambda
xi


$$\lambda = p \text{ m}$$



Physical waves

$\rightarrow \approx 17 \text{ GHz}$

$A(x,y)$



Approx. amplitude at any point x in space wrt time t . Temporal period $T := q \text{ s}$. Temporal freq. $\nu = (1 \text{ s}/q \text{ s}) \text{ c/s} = 1/q \text{ s} = 1 \text{ s}/T \text{ c/s}$.

c means cycles

c/s \rightarrow Hertz
Hz


ν nu

KHz AM
MHz FM
GHz WiFi

Physical waves


WAVEFORMS

THESE ARE
NOT PHYSICAL
WAVES
square pulse sawtooth




Non-smooth (instantaneous change) or non-changing wrt space and time is non-physical.


Physical waves




Emission from a point source in space.
Amplitude $\propto r^{-1}$ for 2-D and $\propto r^{-2}$ for 3-D.



Approx. amplitude along a radial line through space. Spatial wavelength $\lambda := p \text{ m}$. Spatial freq. $\xi := (1 \text{ m}/p \text{ m}) \text{ c/m} = 1/\lambda \text{ c/m}$.




Approx. amplitude at any point x in space wrt time t . Temporal period $T := q \text{ s}$. Temporal freq. $\nu := (1 \text{ s}/q \text{ s}) \text{ c/s} = 1/q \text{ s} = 1/T \text{ c/s}$.




Non-smooth (instantaneous change) or non-changing wrt space and time is non-physical.

Wave summation, $\bar{A}_{1+2} := A_1 + A_2$





θ
 x or t
 theta


Wave summation, $A_{1+2} := A_1 + A_2$




Wave summation, $A_{1+2} := A_1 + A_2$




Wave summation, $A_{1+2} := A_1 + A_2$





Wave summation, $A_{1+2} := A_1 + A_2$




When $\lambda_1 = \lambda_2$ and $\phi_1 = \phi_2$,
 $\lambda_{1+2} = \lambda_1 = \lambda_2$ and $\phi_{1+2} = \phi_1 = \phi_2$.



When $\lambda_1 = \lambda_2$ and $\phi_1 > \phi_2$ and $A_1 > A_2$,
 $\lambda_{1+2} = \lambda_1 = \lambda_2$ and $\phi_1 > \phi_{1+2} > \phi_2$.



Wavefront propagation




wave crest
at time t_0

Δ delta


at time $t_0 + \Delta t$

at time $t_0 + 2\Delta t$


Wavefront propagation







Wavefront propagation




Wavefront propagation




Wavefront propagation




Propagation from a point where amplitude is increasing




...and from another point where amplitude is decreasing





Q Heta



$\cos(\theta)$

Point coordinates corresponding to angle θ are,
 $(\cos \theta, \sin \theta)$.

Angle θ corresponding to point coords (x, y) is,
 $\arccos x$ and $\arcsin y$.

Geometric construction is impractical and mathematical expression is complicated:


$$\sin \theta = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} \theta^{2n+1}$$

so calculators with programmed buttons or printed tables are used.

Sinusoids with same λ but arbitrary ϕ and A sum to a sinusoid with same λ .

This is how physical waves behave.

Sinusoids are *only* periodic functions with this property.



For any point (x, y) on unit circle,
 $\cos \theta := x$ and $\sin \theta := y$.

Point coordinates corresponding to angle θ are,
 $(\cos \theta, \sin \theta)$.

Angle θ corresponding to point coords (x, y) is,
 $\arccos x$ and $\arcsin y$.

Geometric construction is impractical and mathematical expression is complicated:

$$\sin \theta = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} \theta^{2n+1}$$

so calculators with programmed buttons or printed tables are used.

Sinusoids with same λ but arbitrary ϕ and A sum to a sinusoid with same λ .

This is how physical waves behave.

Sinusoids are *only* periodic functions with this property.

Wave equation

How can waves be described so their behaviour can be analysed mathematically?

“They look like sinusoids” isn’t rigorous enough.

Wave equation

$$\frac{\partial^2 A(x,t)}{\partial x^2} - \frac{\partial^2 A(x,t)}{\partial t^2}$$

DIFFERENTIAL EQUATION

We will soon derive this constraint equation from Hooke's and Newton's Laws:

$$\boxed{\frac{\partial^2 A}{\partial x^2}} - \left(\frac{1}{c^2}\right) \frac{\partial^2 A}{\partial t^2} = 0$$

$$\frac{\partial^2 A}{\partial x^2} = \frac{1}{c^2} \frac{\partial^2 A}{\partial t^2}$$


Spatial behaviour temporal behaviour

Wave equation

One of many solutions can be found algebraically as,

$$A(x, t) = R \cos(k x - \omega t) \\ + (1 - R) \cos(k x + \omega t)$$

where k and ω are constants related to (angular) wavelength and frequency and $|R| \leq 1$.



Wave equation

How can waves be described so their behaviour can be analysed mathematically?

“They look like sinusoids” isn’t rigorous enough.


We will soon derive this constraint equation from Hooke’s and Newton’s Laws:

$$\frac{\partial^2 A}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2 A}{\partial t^2} = 0$$

One of many solutions can be found algebraically as,

$$A(x, t) = R \cos(kx - \omega t) + (1 - R) \cos(kx + \omega t)$$

where k and ω are constants related to (angular) wavelength and frequency and $|R| \leq 1$.



The sum of two weighted sinusoids travelling in opposite directions. Here for $R = 0.33$.

Assignment # 1: Huygens-Fresnel construction

- ▶ Write a SciPy program to make at least one plot similar to those shown the wavefront propagation slide.
- ▶ Use Huygens-Fresnel construction to determine where the wavefront should be at different times.
- ▶ Make it into a self-contained project repository in your personal account on gitlab.scss.tcd.ie.
- ▶ Add fshevlin@tcd.ie as a member with “reporter” privileges.

Assignment # 1: Huygens-Fresnel construction

- ▶ Write a SciPy program to make at least one plot similar to those shown the wavefront propagation slide.
- ▶ Use Huygens-Fresnel construction to determine where the wavefront should be at different times.
- ▶ Make it into a self-contained project repository in your personal account on gitlab.scss.tcd.ie.
- ▶ Add fshevlin@tcd.ie as a member with “reporter” privileges.

Assignment # 1: Huygens-Fresnel construction

- ▶ Write a SciPy program to make at least one plot similar to those shown the wavefront propagation slide.
- ▶ Use Huygens-Fresnel construction to determine where the wavefront should be at different times.
- ▶ Make it into a self-contained project repository in your personal account on gitlab.scss.tcd.ie.
- ▶ Add `fshevlín@tcd.ie` as a member with “reporter” privileges.

Greek letters* often used as symbols in mathematics

α	alpha	θ	theta	\omicron	omicron	τ	tau
β	beta	ϑ	caligr. theta	π	pi	υ	upsilon
γ	gamma	ι	iota	ϖ	caligr. pi	ϕ	phi
δ	delta	κ	kappa	ρ	rho	φ	caligr. phi
ϵ	epsilon	λ	lambda	ϱ	caligr. rho	χ	chi
ε	caligr. epsilon	μ	mu	σ	sigma	ψ	psi
ζ	zeta	ν	nu	ς	caligr. sigma	ω	omega
η	eta	ξ	xi				
Γ	big gamma	Λ	big lambda	Σ	big sigma	Ψ	big psi
Δ	big delta	Ξ	big xi	Υ	big upsilon	Ω	big omega
Θ	big theta	Π	big pi	Φ	big phi		

*With their anglophone pronunciations.