

Student Online Teaching Advice Notice

- ▶ The materials and content presented within this session are intended solely for use in a context of teaching and learning at Trinity.
- ▶ Any session recorded for subsequent review is made available solely for the purpose of enhancing student learning.
- ▶ Students should not edit or modify the recording in any way, nor disseminate it for use outside of a context of teaching and learning at Trinity.
- ▶ Please be mindful of your physical environment and conscious of what may be captured by the device camera and microphone during videoconferencing calls.
- ▶ Recorded materials will be handled in compliance with Trinity's statutory duties under the Universities Act, 1997 and in accordance with the University's policies and procedures.
- ▶ Further information on data protection and best practice when using videoconferencing software is available at https://www.tcd.ie/info_compliance/data-protection/

©Trinity College Dublin 2021

1 / 16

Notes

CS7GV2: Mathematics of Light and Sound, M.Sc. in Computer Science.

Lecture #2: Sound and Vibration

Fergal Shevlin, Ph.D.

School of Computer Science and Statistics,
Trinity College Dublin


October 8, 2021

Notes


2 / 16

Acoustic waves


- ▶ Propagation of energy through matter by oscillation of pressure or of displacement. Speed $c_s = \sqrt{K/\rho} \text{ m s}^{-1}$.
- ▶ No heat or mass is transferred.
- ▶ Can be reflected, refracted, diffracted, and/or attenuated by the medium.
- ▶ For lots of interesting simulations, see <https://tinyurl.com/yyv5sajz>



Longitudinal waves have variations around equilibrium pressure due to compression and rarefaction of the medium in the direction of propagation.



Transverse waves have surface deformations perpendicular to the direction of wave propagation.



In solids, combinations of wave types can cause particles to move in elliptical trajectories with depth-dependent direction. In liquids, particles move in anti-clockwise circular trajectories.

Notes


3 / 16

Vibration


- ▶ Oscillations about equilibrium of a material, a structure, or a mechanical system.
- ▶ Can be periodic (e.g. a pendulum) or random (e.g. wheel on gravel road.)
- ▶ *Free vibration* is when there is an initial disturbance only (e.g. a tuning fork.)
- ▶ Objects have natural *frequencies* and *modes* of free vibration.

- ▶ Modes of a cantilevered I-beam. <https://tinyurl.com/yyvosd69>
- ▶ Modes of a membrane under tension. <https://tinyurl.com/yad48x9r>
- ▶ Mode-like basis functions used in linear combination to describe other shapes. <https://tinyurl.com/y4prwl9q>

Analysis of vibration of an object with one degree-of-freedom to move (out of six in three-dimensional space) usually starts with a simple mass-spring or mass-spring-damper model.



Analysis of an object with multiple degrees-of-freedom usually starts with an assembly of simple models, one for each degree.




Notes

4 / 16

Simple Harmonic Oscillator

To develop a mathematical expression for vibration, consider some physical laws that a simple model should obey,


- ▶ Hooke's law: Spring force scales linearly with distance of displacement from equilibrium. $F = -k x$.
- ▶ Newton's second law of motion: Force equals mass by acceleration. $F = m a$.



Equating the two expressions for F yields a single differential equation that relates all parameters,

$$m \frac{d^2 x}{dt^2} + k x = 0.$$

cf. "free-body diagrams."



Friction can be modelled as a damper with force $F = -c \frac{dx}{dt}$ that can be added into the equation,

$$m \frac{d^2 x}{dt^2} + c \frac{dx}{dt} + k x = 0.$$

A solution for the undamped system is,

$$x(t) = x_0 \cos(\omega t + \phi)$$

where $\omega = \sqrt{k/m}$ is angular freq. $f = \omega/2\pi$ is temporal freq. x_0 is the amplitude of initial displacement and ϕ is phase which in this case is 0.

A solution for damped with $\zeta = \frac{c}{2\sqrt{km}}$ is,


$$x(t) = x_0 e^{-\zeta \omega t} \cos(\sqrt{1 - \zeta^2} \omega t + \phi).$$

5 / 16

Notes

Wave equation*

One way to model a continuous (non-discrete) system such as a string is to consider it as a series of masses m connected by springs of lengths h and spring constants k ,



where $u(x)$ is the distance from its equilibrium position of a mass at position x .

$$\frac{\partial^2}{\partial t^2} u(x+h, t) = \frac{k}{m} [u(x+2h, t) - u(x+h, t) - u(x+h, t) + u(x, t)].$$

For N masses evenly spaced over total length $L = Nh$ and total mass $M = Nm$ and a total spring constant $K = k/N$, the rhs becomes,

$$\frac{KL^2}{M} \frac{[u(x+2h, t) - 2u(x+h, t) + u(x, t)]}{h^2}.$$

The force acting on mass m at position $x+h$ at time t can be described independently:

$$\text{Newton's, } F(x+h, t) = m \frac{\partial^2}{\partial t^2} u(x+h, t).$$

$$\begin{aligned} \text{Hooke's, } F(x+h, t) &= F(x+2h, t) - F(x, t) \\ &= k [u(x+2h, t) - u(x+h, t)] \\ &\quad - k [u(x+h, t) - u(x, t)]. \end{aligned}$$

Let $c^2 = \frac{KL^2}{M}$ and consider the continuous system situation where $N \rightarrow \infty$ (which means taking the limit as $h \rightarrow 0$.)

$$\frac{\partial^2 u(x, t)}{\partial t^2} = c^2 \frac{\partial^2 u(x, t)}{\partial x^2}.$$

*See <https://tinyurl.com/y6acqr4j>.

6 / 16

Notes

Taking the limit

$$\lim_{h \rightarrow 0} \frac{u(x+2h, t) - 2u(x+h, t) + u(x, t)}{h^2}$$

This expression is *indeterminate* because for $h = 0$ it becomes $\frac{0}{0}$ which has multiple interpretations:

0? ∞ ? 1?

Luckily, we have l'Hospital's Rule which says that, under certain conditions,

$$\lim_{x \rightarrow y} \frac{f(x)}{g(x)} = \lim_{x \rightarrow y} \frac{f'(x)}{g'(x)}.$$

So the quotient terms can be replaced by their derivatives, $\frac{d}{dh} h^2 = 2h$ and

$$\begin{aligned} \frac{\partial}{\partial h} [u(x+2h, t) - 2u(x+h, t) + u(x, t)] &= \\ 2u'(x+2h, t) - 2u'(x+h, t). \end{aligned}$$

$$\lim_{h \rightarrow 0} \frac{u(x+2h, t) - 2u(x+h, t) + u(x, t)}{h^2} =$$

$$\lim_{h \rightarrow 0} \frac{2u'(x+2h, t) - 2u'(x+h, t)}{2h} =$$

$$\lim_{h \rightarrow 0} \frac{4u''(x+2h, t) - 2u''(x+h, t)}{2} =$$

$$\lim_{h \rightarrow 0} [2u''(x+2h, t) - u''(x+h, t)] =$$

$$2 \lim_{h \rightarrow 0} u''(x+2h, t) - \lim_{h \rightarrow 0} u''(x+h, t) =$$

$$2u''(x, t) - u''(x, t) =$$

$$u''(x, t) = \frac{\partial^2 u(x, t)}{\partial x^2}.$$

7 / 16

Notes

Wave equation solution

A solution for $u(x, t)$ is:

$$R \cos(kx - \omega t) + (1 - R) \cos(kx + \omega t)$$

ω is angular frequency $2\pi\nu$ in rad s^{-1}

k is the wave number $2\pi/\lambda$ in rad m^{-1}

$|R| \leq 1$ specifies direction of travel.

Which is the superposition of two sinusoidal waves travelling in opposite directions. Non-sinusoidal solutions are possible too.

Notes

8 / 16

d'Alembert's approach

The wave equation is a hyperbolic linear second order partial differential equation (PDE,)

$$\frac{\partial^2 u}{\partial x^2} - \frac{1}{c_s^2} \frac{\partial^2 u}{\partial t^2} = 0.$$

A solution is the function $u(x, t)$ which satisfies it. One approach to finding a solution is to transform the differential equation into a form for which a solution is already known.

The first step is to change the variables, e.g. let $\xi \equiv x - c_s t$ and $\eta \equiv x + c_s t$,

$$x = \frac{1}{2}(\xi + \eta) \text{ and } t = \frac{1}{2c_s}(\eta - \xi).$$

Notes

9 / 16

By the Chain Rule,

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial \xi} \frac{\partial \xi}{\partial x} + \frac{\partial u}{\partial \eta} \frac{\partial \eta}{\partial x} = \frac{\partial u}{\partial \xi} + \frac{\partial u}{\partial \eta}$$

$$\frac{\partial u}{\partial t} = \frac{\partial u}{\partial \xi} \frac{\partial \xi}{\partial t} + \frac{\partial u}{\partial \eta} \frac{\partial \eta}{\partial t} = -c_s \frac{\partial u}{\partial \xi} + c_s \frac{\partial u}{\partial \eta}$$

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial x} \right) = \frac{\partial}{\partial \xi} \left(\frac{\partial u}{\partial \xi} + \frac{\partial u}{\partial \eta} \right)$$

$$= \left(\frac{\partial \xi}{\partial x} \frac{\partial}{\partial \xi} + \frac{\partial \eta}{\partial x} \frac{\partial}{\partial \eta} \right) \left(\frac{\partial u}{\partial \xi} + \frac{\partial u}{\partial \eta} \right) = \frac{\partial^2 u}{\partial \xi^2} + 2 \frac{\partial^2 u}{\partial \xi \partial \eta} + \frac{\partial^2 u}{\partial \eta^2}$$

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial}{\partial t} \left(\frac{\partial u}{\partial t} \right) = \frac{\partial}{\partial \eta} \left(-c_s \frac{\partial u}{\partial \xi} + c_s \frac{\partial u}{\partial \eta} \right)$$

$$= \left(-c_s \frac{\partial \xi}{\partial t} \frac{\partial}{\partial \xi} + c_s \frac{\partial \eta}{\partial t} \frac{\partial}{\partial \eta} \right) \left(-c_s \frac{\partial u}{\partial \xi} + c_s \frac{\partial u}{\partial \eta} \right) = c_s^2 \frac{\partial^2 u}{\partial \xi^2} - 2c_s^2 \frac{\partial^2 u}{\partial \xi \partial \eta} + c_s^2 \frac{\partial^2 u}{\partial \eta^2}.$$

Notes

10 / 16

So the wave equation $\frac{\partial^2 u}{\partial x^2} - \frac{1}{c_s^2} \frac{\partial^2 u}{\partial t^2} = 0$ becomes

$$\left(\frac{\partial^2 u}{\partial \xi^2} + 2 \frac{\partial u}{\partial \xi} \eta + \frac{\partial^2 u}{\partial \eta^2} \right) - \frac{1}{c_s^2} \left(c_s^2 \frac{\partial^2 u}{\partial \xi^2} - 2 c_s^2 \frac{\partial u}{\partial \xi} \eta + c_s^2 \frac{\partial^2 u}{\partial \eta^2} \right) = 0$$

$$\text{which is } \frac{\partial u}{\partial \xi} \eta = \frac{\partial}{\partial \xi} \left(\frac{\partial u}{\partial \eta} \right) = 0.$$

for which any solution is known to have the form:

$$p(\xi, \eta) = f(\eta) + g(\xi) = f(x + c_s t) + g(x - c_s t)$$

Notes

11 / 16

Specific solutions

To find functions f and g which describe a specific wave p we need to have some:

Boundary conditions e.g. $p(0, t) = 0$ and $p(L, t) = 0$.

Initial conditions e.g. $p(x, 0) = f(x)$ and $\frac{\partial u}{\partial t}(x, 0) = g(x)$.

We might revisit this topic in more detail later.

Notes

12 / 16

Chain Rule(s) of differentiation

- ▶ Rules that specify derivatives for compositions of functions, e.g. for $f(g(x))$, the derivative is $f'(g(x)) g'(x)$.

- ▶ Leibnitz notation often used: let $y \equiv f(g(x))$ and $u \equiv g(x)$,

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = f'(g(x)) g'(x).$$

- ▶ Let $z \equiv f(x, y)$ and $x \equiv g(t)$ and $y \equiv h(t)$ then

$$\frac{dz}{dt} = \frac{\partial z}{\partial x} \frac{dx}{dt} + \frac{\partial z}{\partial y} \frac{dy}{dt}.$$

- ▶ Let $z \equiv f(x, y)$ and $x \equiv g(s, t)$ and $y \equiv h(s, t)$ then

$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial s} \frac{ds}{dx} + \frac{\partial z}{\partial t} \frac{dt}{dx}$$

Notes

13 / 16

An easier approach

- ▶ To solve PDEs, you can use a mathematics-oriented symbolic programming system like Mathematica (e.g. through the free Wolfram Alpha web interface.)

```
DSolve[D[u[x, t], {x, 2}] - (1/c^2) D[u[x, t], {t, 2}]  
== 0, u[, ], {t, x}]
```

```
{u[x, t] -> C[1] [-sqrt(c^2) t + x] +  
C[2] [sqrt(c^2) t + x]}
```

For arbitrary functions $C[1]$ and $C[2]$ which can be found once initial values and/or boundary conditions are known.

- ▶ Note that this looks different to previous solution whose parameters are expressed in angular distance (radians) not linear distance (metres.)

Notes

14 / 16

Assignment # 2: Acoustic wave simulation

- ▶ Write a SciPy program to make at least one plot similar to those shown the acoustic wave slide.
- ▶ Demonstrate wave propagation over time through either a set of pictures or an animation.
- ▶ Make it into a self-contained project repository in your personal account on gitlab.scss.tcd.ie.
- ▶ For this and every other assignment, feel free to collaborate with your classmates about the the non-mathematical parts like plotting.

Notes

15 / 16

Assignment # 3: Simple harmonic oscillator simulation

- ▶ Write a SciPy program to make an animation of simple harmonic motion.
- ▶ In addition to an object moving up and down, you could show a sinusoidal plot of its amplitude.
- ▶ Make it into a self-contained project repository in your personal account on gitlab.scss.tcd.ie.
- ▶ For this and every other assignment, feel free to collaborate with your classmates about the the non-mathematical parts like plotting.

Notes

16 / 16