Macroeconomics in the Sequence Space

Chen Gao Weigao Wang

School of Economics, Peking University

April 14, 2026

Outline

1. Why Sequence Space?
2. Local Solution
   - 2.1 SSJ in Krusell-Smith Model
   - 2.2 Fake News Algorithm
   - 2.3 Theoretical progress after SSJ
   - 2.4 Application
3. Global Solution
   - 3.1 Repeated transition method
   - 3.2 DL in sequence space
   - 3.3 RL in sequence space
4. Reference

Why Sequence Space?

Sequence Space vs. State Space

Motivation

- In heterogeneous-agent (HA) models, the economy's state is the entire cross-sectional distribution — an infinite-dimensional object.
- Sequence-space methods *sidestep* this challenge: they characterize local dynamics around the steady state without ever carrying the distribution as a state variable.
- The key insight: aggregate equilibrium responses to shocks can be encoded in Jacobian matrices of aggregate outcomes with respect to aggregate inputs.

Sequence Space vs. State Space

State Space

Definition 1

A state-space representation describes the economy recursively. At date t, a state variable x_t summarizes all payoff-relevant history, and the model evolves as

$$x_{t+1} = f(x_t, \varepsilon_{t+1}), \qquad y_t = g(x_t).$$

Given x_t, the future is independent of the past.

Definition 2

A sequence-space representation characterizes equilibrium as a mapping between entire time paths. We seek sequences $\{X_t\}_{t=0}^{\infty}$ such that equilibrium conditions hold at every date:

$$F(\{X_t\}_{t=0}^{\infty}, \{Z_t\}_{t=0}^{\infty}) = \mathbf{0},$$

where $\{X_t\}$ collects endogenous variables and $\{Z_t\}$ collects exogenous shocks or policy paths.

Sequence Space vs. State Space

Comparison

Intuition

- State space asks: *what is the current state, and how does it transition?*
- Sequence space asks: *given a shock path, what equilibrium outcome path clears all markets?*

Takeaway

When the distribution is the intractable state variable, sequence space reframes a "huge state" problem as a structured Jacobian problem — amenable to fast, scalable computation.

Local Solution

Example: The Krusell-Smith Model

Setup

- The economy is populated by a mass 1 of heterogeneous households,
- Each household maximizes its expected lifetime utility:

$$\mathbb{E}_0 \left[\sum_{t=0}^{\infty} \beta^t u(c_t) \right],$$

- Standard CRRA utility,
- There exist n_e idiosyncratic states w/ transition matrix $P(e, e')$ and stationary distribution π,
- Agents supply an exogenous number of hours n, and earn wage income $w_t e n$,
- Agents can only trade in capital k, which pays a rental rate r_t net of depreciation.

Example: The Krusell-Smith Model

Household Problem

The value function of an agent entering period w/ state (e, k_-) is given by:

$$V_t(e, k_-) = \max_{c,\, k} u(c) + \beta \sum_{e'} V_{t+1}(e', k)\, P(e, e')$$

$$\text{s.t.} \qquad c + k = (1 + r_t)k_- + w_t e n$$

$$k \geq 0$$

- Denote by $c_t^*(e, k_-)$ and $k_t^*(e, k_-)$ the policy functions that solve the Bellman equation.
- Also denote by $D_t(e, K_-)$ the measure of households in state e that own capital in a set K_- at the start of date t.

Example: The Krusell-Smith Model

Household Problem

The distribution D_t has law of motion

$$D_{t+1}(e', K) = D_t(e, k_t^{*-1}(e, K))P(e, e').$$

where $k_t^{*-1}(e, K)$ denotes the inverse of $k_t^*(e, K)$.

- We assume that prior to $t = 0$, the economy is in a steady state.
- We suppose that agents start in this stationary distribution at date 0, so that $D_0 = D_{ss}$.

Example: The Krusell-Smith Model

Aggregate Capital

- For any t, the policy $k_t^*(e, k_-)$ is a function of the future path $\{r_s, w_s\}_{s=t}^{\infty}$.
- Given $D_0 = D_{ss}$, the distribution $D_t(e, K)$ at any t is a function of the entire path $\{r_s, w_s\}_{s=0}^{\infty}$.
- The aggregate household capital holdings are characterized by a capital function $\mathcal{K}_t(\{r_s, w_s\}_{s=0}^{\infty})$.

$$\mathcal{K}_t\left(\{r_s, w_s\}_{s=0}^{\infty}\right) = \sum_e \int_{k_-} k_t^*(e, k_-) D_t(e, dk_-)$$

Key Insight

The ability to reduce interactions between heterogeneous agents to functions such as $\mathcal{K}_t$, which map aggregate sequences into aggregate sequences, is key to the sequence-space Jacobian method.

Example: The Krusell-Smith Model

Production

- Firm adopts a Cobb-Douglas production function:

$$Y_t = Z_t K_t^{\alpha} N_t^{1-\alpha}$$

- The firm's first-order conditions:

$$r_t = \alpha Z_t K_{t-1}^{\alpha-1} N_t^{1-\alpha} - \delta$$
$$w_t = (1-\alpha) Z_t K_{t-1}^{\alpha} N_t^{-\alpha}$$

Key Insight

The paths of prices $\{r_t, w_t\}_{t=0}^{\infty}$ are now functions of the exogenous paths $\{Z_t, N_t\}_{t=0}^{\infty}$ and the endogenous path $\{K_t\}_{t=0}^{\infty}$.

Example: The Krusell-Smith Model

Capital Market-Clearing Condition

Combining together, we can express the capital market-clearing condition at each point in time as a function H_t:

$$H_t(\boldsymbol{K}, \boldsymbol{Z}) \equiv \mathcal{K}_t \left(\left\{ \underbrace{\alpha Z_s \left(\frac{K_{s-1}}{\sum \pi(e) en} \right)^{\alpha-1} - \delta}_{r_s}, \underbrace{(1-\alpha) Z_s \left(\frac{K_{s-1}}{\sum \pi(e) en} \right)^{\alpha}}_{w_s} \right\}_{s=0}^{\infty} \right) - K_t = 0$$

where $\boldsymbol{K}$ is the vector of capital holdings $\{K_t\}_{t=0}^{\infty}$ and $\boldsymbol{Z}$ is the vector of exogenous shocks $\{Z_t\}_{t=0}^{\infty}$.

Equilibrium Capital Path

Given initial capital K_{-1} and the exogenous path for productivity $\boldsymbol{Z}$, we have equilibrium path of capital $\{K_t\}_{t=0}^{\infty}$ by solving the nonlinear equation $H_t(\boldsymbol{K}, \boldsymbol{Z}) = 0$.

Example: The Krusell-Smith Model

IRF

By *implicit function theorem*, given an exo. shock $\mathrm{d}\boldsymbol{Z} = (\mathrm{d}Z_0, \mathrm{d}Z_1 \ldots)'$ at $t = 0$, the linear impulse response of capital to this shock is given by:

$$d\boldsymbol{K} = -H_K^{-1} H_Z \, d\boldsymbol{Z}$$

where H_K and H_Z denote the Jacobians of H with respect to $\boldsymbol{K}$ and $\boldsymbol{Z}$, evaluated at the steady state.

- Given $\mathrm{d}\boldsymbol{K}$, the impulse responses of other variables—for example, $\{Y_t, r_t, w_t\}_{t=0}^{\infty}$—follow immediately.
- In practice, $H_t(\boldsymbol{K}, \boldsymbol{Z}) = 0$ is solved up to a sufficiently large horizon T such that $\boldsymbol{K}$ and $\boldsymbol{Z}$ have approximately returned to their steady states by time T.

Example: The Krusell-Smith Model

Constructing the Jacobian

- We use the *chain rule* to relate the Jacobians H_K and H_Z to the derivatives of the $\mathcal{K}$.
- For example, differentiating $H_t(\boldsymbol{K}, \boldsymbol{Z}) = 0$ with respect to K_s
- The (t, s) entry of H_K is:

$$[H_K]_{t,s} = \frac{\partial \mathcal{K}_t}{\partial r_{s+1}} \frac{\partial r_{s+1}}{\partial K_s} + \frac{\partial \mathcal{K}_t}{\partial w_{s+1}} \frac{\partial w_{s+1}}{\partial K_s} - \mathbf{1}_{\{s=t\}}$$

- A similar expression applies to H_Z.
- In addition, the derivatives such as $\frac{\partial r_{s+1}}{\partial K_s}$ at (K^{ss}, Z^{ss}) can all be computed analytically.
- For example:

$$\frac{\partial r_{s+1}}{\partial K_s} = \alpha(\alpha - 1) Z^{ss} (K^{ss})^{\alpha-2} (N^{ss})^{1-\alpha}$$

Example: The Krusell-Smith Model

Constructing the Jacobian

Recall that:

$$d\boldsymbol{K} = -H_K^{-1} H_Z \, d\boldsymbol{Z},$$

To get $H_K^{-1} H_Z$, the only thing left is to solve the Jacobians of the $\mathcal{K}$ function with respect to its two inputs r and w.

$$[H_K]_{t,s} = \frac{\partial \mathcal{K}_t}{\partial r_{s+1}} \frac{\partial r_{s+1}}{\partial K_s} + \frac{\partial \mathcal{K}_t}{\partial w_{s+1}} \frac{\partial w_{s+1}}{\partial K_s} - \mathbf{1}_{\{s=t\}}$$

Let

$$\mathcal{J}_{t,s}^{\mathcal{K},r} = \frac{\partial \mathcal{K}_t}{\partial r_s}, \quad \mathcal{J}_{t,s}^{\mathcal{K},w} = \frac{\partial \mathcal{K}_t}{\partial w_s}$$

be the Jacobians up to some truncation horizon T.

Example: The Krusell-Smith Model

Constructing the Jacobian

Sufficient statistics

The Jacobians $\mathcal{J}^{\mathcal{K},r}$ and $\mathcal{J}^{\mathcal{K},w}$ are sufficient statistics for the household in sequence space equilibrium:

- Shocks propagate through the dynamics of the household capital distribution, but *we do not need those details if we have the Jacobians.*
- Column s of $\mathcal{J}^{\mathcal{K},r}$ is the impulse response of capital to a news shock dr_s; the Jacobian gives these for all $s = 0, \ldots, T-1$.
- These are rich objects, but *there is a fast new algorithm to compute them.*

Fake News Algorithm

General Model Representation

We represent a heterogeneous-agent problem as a mapping from a time path of aggregate inputs $\{\boldsymbol{X}_t\}$ to a time path of aggregate outputs $\{\boldsymbol{Y}_t\}$.

Assume:

- $\boldsymbol{X}_t \in \mathbb{R}^{n_x}$: aggregate inputs,
- $\boldsymbol{Y}_t \in \mathbb{R}^{n_y}$: aggregate outputs,
- the distribution is discretized on n_g grid points,
- $\boldsymbol{D}_t \in \mathbb{R}^{n_g}$: cross-sectional distribution of agents at time t.

Aggregate outputs are averages of individual outcomes against the distribution:

$$\boldsymbol{Y}_t = \boldsymbol{y}_t' \boldsymbol{D}_t,$$

where $\boldsymbol{y}_t$ is an $n_g \times n_y$ matrix of individual outcomes (for example, consumption, savings, labor supply, or other individual-level objects).

Fake News Algorithm

Three Core Equations

We assume there exist three functions v, Λ, and y such that, for a given initial distribution $\boldsymbol{D}_0$, the aggregate sequence solves

$$\boldsymbol{v}_t = v(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t),$$

$$\boldsymbol{D}_{t+1} = \Lambda(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)\, \boldsymbol{D}_t,$$

$$\boldsymbol{Y}_t = y(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)' \boldsymbol{D}_t.$$

Interpretation:

- $\boldsymbol{v}_t$: backward-looking object, typically the value function;
- $\Lambda(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)$: transition matrix governing the law of motion of the distribution;
- $y(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)$: individual outcomes used to aggregate into $\boldsymbol{Y}_t$.

Fake News Algorithm

Steady State and Transition Paths

For a constant aggregate input $\boldsymbol{X}_{ss}$, the steady state is a fixed point

$$(\boldsymbol{Y}_{ss}, \boldsymbol{v}_{ss}, \boldsymbol{D}_{ss})$$

of the system.

For convenience, define

$$\Lambda_{ss} \equiv \Lambda(\boldsymbol{v}_{ss}, \boldsymbol{X}_{ss}), \qquad y_{ss} \equiv y(\boldsymbol{v}_{ss}, \boldsymbol{X}_{ss}).$$

We consider transition paths of length T that end at the steady state:

$$\boldsymbol{X}_{T-1} = \boldsymbol{X}_{ss}, \qquad \boldsymbol{v}_T = \boldsymbol{v}_{ss}.$$

The initial distribution $\boldsymbol{D}_0$ is given and often set equal to $\boldsymbol{D}_{ss}$.

Hence, this framework is designed to study transitory shocks around a steady state.

Fake News Algorithm

Economic Meaning of the System

Step 1: Backward

Given the future object $\boldsymbol{v}_{t+1}$ and current aggregates $\boldsymbol{X}_t$, solve for the current object $\boldsymbol{v}_t$.

Step 2: Forward

Use $\Lambda(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)$ to update the cross-sectional distribution:

$$\boldsymbol{D}_{t+1} = \Lambda(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)\boldsymbol{D}_t.$$

Step 3: Aggregate

Aggregate individual outcomes across agents:

$$\boldsymbol{Y}_t = y(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)'\boldsymbol{D}_t.$$

Fake News Algorithm

Three-Step Structure of the HA Model

Fake News Algorithm

Three-Step Structure of the HA Model

Step 1: Backward

$$\boldsymbol{v}_t = v(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)$$



Fake News Algorithm

Three-Step Structure of the HA Model

Step 1: Backward

$$\boldsymbol{v}_t = v(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)$$



Step 2: Forward

$$\boldsymbol{D}_{t+1} = \Lambda(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)\boldsymbol{D}_t$$

Fake News Algorithm

Three-Step Structure of the HA Model

Step 1: Backward

$$\boldsymbol{v}_t = v(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)$$



Step 2: Forward

$$\boldsymbol{D}_{t+1} = \Lambda(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)\boldsymbol{D}_t$$

Step 3: Aggregate

$$\boldsymbol{Y}_t = y(\boldsymbol{v}_{t+1}, \boldsymbol{X}_t)'\boldsymbol{D}_t$$

Fake News Algorithm

Calculating the Jacobian: direct vs. new algorithm

- The direct algorithm requires a separate backward and forward iteration for each $s = 0, \ldots, T-1$; this is costly.
- SSJ method uses a single backward and forward iteration to get all T columns.
- *This improves speed by a factor of roughly T with no loss in accuracy.*
- The key is to exploit time symmetries in the linearized system.

The linearized system:

$$
\begin{aligned}
d\boldsymbol{Y}_t &= d\boldsymbol{y}_t' \boldsymbol{D}_{ss} + \boldsymbol{y}_{ss}' \, d\boldsymbol{D}_t \\
d\boldsymbol{D}_{t+1} &= d\Lambda_t \boldsymbol{D}_{ss} + \Lambda_{ss} d\boldsymbol{D}_t \\
\boldsymbol{D}_0 &= d\boldsymbol{D}_{ss}
\end{aligned}
$$

Since everything is linear, we can look at terms from backward and forward iterations separately, then combine them.

Fake News Algorithm

Terms from backward iteration

Holding $\boldsymbol{D}_t = \boldsymbol{D}_{ss}$ fixed, what is the effect of $d\boldsymbol{X}_s$?

$$d\boldsymbol{Y}_t = d\boldsymbol{y}_t^0 \boldsymbol{D}_{ss} + \cancel{\boldsymbol{y}'_{ss}\, d\boldsymbol{D}_t}$$
$$d\boldsymbol{D}_{t+1} = d\Lambda_t^0 \boldsymbol{D}_{ss} + \cancel{\Lambda_{ss}\, d\boldsymbol{D}_t}$$

- $d\boldsymbol{y}_t^0$ and $d\Lambda_t^0$ are purely forward-looking from the Bellman equation.
- They depend only on the time until the shock, $s - t$.

Define:

$$\mathcal{Y}_{s-t} \equiv \frac{d\boldsymbol{y}_t^0 \boldsymbol{D}_{ss}}{d\boldsymbol{X}_s} \qquad \mathcal{D}_{s-t} \equiv \frac{d\Lambda_t^0 \boldsymbol{D}_{ss}}{d\boldsymbol{X}_s}$$

Use a single backward iteration with a shock $d\boldsymbol{X}_s$ at $s = T - 1$ to obtain $d\boldsymbol{y}_t^0$ and $d\Lambda_t^0$ for all $t = T - 1, \ldots, 0$.

Fake News Algorithm

Terms from forward iteration

Holding $\boldsymbol{y}_t = \boldsymbol{y}_{ss}$ and $\Lambda_t = \Lambda_{ss}$ fixed, what is the effect of $d\boldsymbol{D}_t$?

$$d\boldsymbol{Y}_t = \cancel{d\boldsymbol{y}_t^0 \boldsymbol{D}_{ss}} + \boldsymbol{y}'_{ss}\, d\boldsymbol{D}_t$$
$$d\boldsymbol{D}_{t+1} = \cancel{d\Lambda_t^0 \boldsymbol{D}_{ss}} + \Lambda_{ss} d\boldsymbol{D}_t$$

Since everything's linear, some linear functional (row vector) maps $d\boldsymbol{D}_t$ to $d\boldsymbol{Y}_t$: what is it?

- Can write for $s \leq t$ (zero for $s > t$):

$$d\boldsymbol{Y}_t = \underbrace{\boldsymbol{y}'_{ss}(\Lambda'_{ss})^{t-s}}_{\equiv \mathcal{P}'_{t-s}} d\boldsymbol{D}_s$$

- These $\mathcal{P}_u$ can be calculated recursively as:

$$\mathcal{P}_u = \Lambda'_{ss}\mathcal{P}_{u-1}$$

- Thus, a single "transpose" forward iteration gives all $\mathcal{P}_u$.

Fake News Algorithm

Putting it together: the Jacobian

For shock $d\boldsymbol{X}_s$, we want:

$$d\boldsymbol{Y}_t = d\boldsymbol{y}_t'\boldsymbol{D}_{ss} + \boldsymbol{y}_{ss}'\, d\boldsymbol{D}_t$$

Change in policy, holding distribution constant:

$$d\boldsymbol{y}_t'\boldsymbol{D}_{ss} = \mathcal{Y}_{s-t} d\boldsymbol{X}_s$$

Change in distribution, holding policy constant:

$$\boldsymbol{y}_{ss}'\, d\boldsymbol{D}_t = (\mathcal{P}_0'\mathcal{D}_{s-t+1} + \ldots + \mathcal{P}_{t-1}'\mathcal{D}_s) d\boldsymbol{X}_s$$

sum of shocks $\mathcal{D}$ to distribution from anticipating $d\boldsymbol{X}_s$ before t, propagated forward to effect on $d\boldsymbol{Y}_t$ with $\mathcal{P}$

The full Jacobian is then:

$$\mathcal{J}_{t,s} = \mathcal{Y}_{s-t} + (\mathcal{P}_0'\mathcal{D}_{s-t+1} + \ldots + \mathcal{P}_{t-1}'\mathcal{D}_s)$$

Fake News Algorithm

Fake news matrix

Define the fake news matrix as:

$$\mathcal{F}_{t,s} \equiv \begin{cases} \mathcal{Y}_s & \text{if } t = 0, \\ \mathcal{P}'_{t-1}\mathcal{D}_s & \text{if } t > 0. \end{cases}$$

- Interpretation:
  - $t = 0$: agents get news of shock at date $s \geq 0$
  - $t \geq 1$: news of unrealized shock disappears ("fake news")
- Build Jacobian (response to news shocks) recursively from fake news matrix (response to fake news shocks) as:

$$\mathcal{J}_{t,s} \equiv \begin{cases} \mathcal{F}_{t,s} & \text{if } t = 0 \text{ or } s = 0, \\ \mathcal{F}_{t,s} + \mathcal{J}_{t-1,s-1} & \text{otherwise} \end{cases}$$

Fake News Algorithm

Algorithm

To obtain Jacobians $\mathcal{J}^{o,i}$ for many inputs dX^i and outputs dY^o:

1. For each input i, perform backward iteration with T steps to get all $\dagger_u^{o,i}$ and $\mathcal{D}_u^i$.
2. For each output o, perform transpose forward iteration with $T-1$ steps to get all $\mathcal{P}_u^o$.
3. For each pair (o,i), construct the fake news matrix $\mathcal{F}^{o,i}$ from $\dagger^{o,i}$ in the first row and the product $(\mathcal{P}^o)'\mathcal{D}^i$ for other rows.
4. For each pair (o,i), recursively compute $\mathcal{J}^{o,i}$ from $\mathcal{F}^{o,i}$.

Step 1 is almost always the bottleneck.

DAG

Krusell-Smith model as a directed acyclic graph (DAG)

Recall that we wrote equilibrium in KS model as system

$$H_t(\boldsymbol{K}, \boldsymbol{Z}) \equiv \mathcal{K}_t \left(\left\{ \underbrace{\alpha Z_s \left(\frac{K_{s-1}}{\sum \pi(e) en} \right)^{\alpha-1} - \delta}_{r_s}, \underbrace{(1-\alpha) Z_s \left(\frac{K_{s-1}}{\sum \pi(e) en} \right)^{\alpha}}_{w_s} \right\}_{s=0}^{\infty} \right) - K_t = 0$$

This corresponds to a simple graph:



DAG

DAGs combining simple and HA blocks



- Each node ("block") takes sequences as inputs and outputs.
- Inputs of later blocks are outputs of earlier blocks.
- The number of endogenous variables ("unknowns") equals the number of equations in H ("targets"); solve to get equilibrium.
- Two kinds of blocks in SHADE models:
  - Simple blocks: Analytical equations directly in aggregates, e.g. $r_t = \alpha Z_t K_{t-1}^{\alpha-1} - \delta$; Jacobians are sparse and easy to obtain.
  - HA blocks: As described previously.

DAG

DAGs combining simple and HA blocks



- Define $\mathrm{J}^{o,i}$ to be the total derivative of o along the DAG with respect to the exogenous or endogenous variable i (distinct from the block Jacobian $\mathcal{J}^{o,i}$).
- Here, for example, $\mathrm{J}^{r,K}$ equals $\mathcal{J}^{r,K}$, etc.: the direct effect is the only effect.
- Then we need the chain rule. For instance:

$$\mathrm{J}^{\mathcal{K},K} = \mathcal{J}^{\mathcal{K},r}\mathrm{J}^{r,K} + \mathcal{J}^{\mathcal{K},w}\mathrm{J}^{w,K}$$

- Finally, we obtain H_K and H_Z; then $\mathrm{G}^{K,Z} \equiv -\mathrm{H}_K^{-1}\mathrm{H}_Z$ gives the general equilibrium map from Z to K, and all $\mathrm{G}^{o,Z}$ are determined from it.

DAG

General case: solving with forward accumulation

- Initialize $\mathrm{J}^{i,i}$ as the identity for all exogenous or unknown i.
- Evaluate the chain rule following a topological sort:

$$\mathrm{J}^{o,i} = \sum_{m \in I_b} \mathcal{J}^{o,m} \mathrm{J}^{m,i}$$

  on blocks b to get $\mathrm{H}_U = \mathrm{J}^{H,U}$ and $\mathrm{H}_Z = \mathrm{J}^{H,Z}$.
- This is called "forward accumulation" in the algorithmic differentiation literature.
- Compute the general equilibrium mapping: $\mathrm{G}^{U,Z} = -\mathrm{H}_U^{-1}\mathrm{H}_Z$
- Then do forward accumulation again:

$$\mathrm{G}^{o,Z} = \sum_{m \in I_b} \mathcal{J}^{o,m} \mathrm{G}^{m,Z}$$

  to get general equilibrium mappings $\mathrm{G}^{o,Z}$ from shocks dZ to all variables o of interest.
- This gives impulse responses to every shock simultaneously.

Determinacy and Large-Scale Solutions in the Sequence Space

To be developed

2nd SSJ

To be developed

SSJ in Continuous Time

To be developed

Applications of the extended version of SSJ

- Fiscal and monetary policy [Auclert et al., 2025c, Auclert et al., 2024b]
- Het households + het firms [Winberry et al., 2025]
- Behavioural macro [Auclert et al., 2020, Angeletos et al., 2025]
- Life-cycle model
  [Bardóczy and Velásquez-Giraldo, 2024, Bardoczy and Velasquez Giraldo, 2025]
- Dynamic spatial model [Donald et al., 2025, Fan et al., 2023]
- Production network [Schaab and Tan, 2023]
- Het banks [Bellifemine et al., 2022]
- Optimal policy [Auclert et al., 2024a, McKay and Wolf, 2022]
- Macro labor [Guerreiro et al., 2024, Kekre, 2023]

Global Solution

Repeated transition method

To be developed

DL in sequence space

To be developed

RL in sequence space

To be developed

Reference

Reference (1)

Angeletos, G.-M., Guerreiro, J., and Zhang, D. R. (2025). From rank to hank, without fire. Technical report, National Bureau of Economic Research.

Auclert, A., Bardóczy, B., Rognlie, M., and Straub, L. (2021). Using the sequence-space jacobian to solve and estimate heterogeneous-agent models. *Econometrica*, 89(5):2375–2408.

Auclert, A., Cai, M., Rognlie, M., and Straub, L. (2024a). Optimal long-run fiscal policy with heterogeneous agents. *Unpublished Manuscript, Harvard University.*

Auclert, A., Majic, E., Rognlie, M., and Straub, L. (2025a). Determinacy and large-scale solutions in the sequence space. *Journal of Political Economy: Macroeconomics.*

Auclert, A., Rigato, R., Rognlie, M., and Straub, L. (2025b). Beyond certainty equivalence: Second-order dynamics in the sequence space. Presentation slides.

Auclert, A., Rognlie, M., and Straub, L. (2020). Micro jumps, macro humps: Monetary policy and business cycles in an estimated hank model. Technical report, National Bureau of Economic Research.

Auclert, A., Rognlie, M., and Straub, L. (2024b). The intertemporal keynesian cross. *Journal of Political Economy*, 132(12):4068–4121.

Reference (2)

Auclert, A., Rognlie, M., and Straub, L. (2025c). Fiscal and monetary policy with heterogeneous agents. *Annual Review of Economics*, 17.

Azinovic-Yang, M. and Žemlička, J. (2025). Deep learning in the sequence space. *arXiv preprint arXiv:2509.13623*.

Bardóczy, B. and Velásquez-Giraldo, M. (2024). Hank comes of age: Monetary policy with heterogeneous overlapping generations.

Bardoczy, B. and Velasquez Giraldo, M. (2025). Sequence-space jacobians of life cycle models. *Sequence-Space Jacobians of Life Cycle Models (May 29, 2025)*.

Bellifemine, M., Jamilov, R., and Monacelli, T. (2022). Hbank: Monetary policy with heterogeneous banks.

Bilal, A. and Goyal, S. (2025). Some pleasant sequence-space arithmetic in continuous time. Technical report, National Bureau of Economic Research.

Donald, E., Fukui, M., and Miyauchi, Y. (2025). Optimal dynamic spatial policy. Technical report, National Bureau of Economic Research.

Fan, J., Hong, S., and Parro, F. (2023). Learning and expectations in dynamic spatial economies. Technical report, National Bureau of Economic Research.

Reference (3)

Guerreiro, J., Hazell, J., Lian, C., and Patterson, C. (2024). Why do workers dislike inflation?: Wage erosion and conflict costs.

Kekre, R. (2023). Unemployment insurance in macroeconomic stabilization. *Review of Economic Studies*, 90(5):2439–2480.

Lee, H. (2025). Global nonlinear solutions in sequence space and the generalized transition function. Technical report, Working paper.

McKay, A. and Wolf, C. (2022). Optimal policy rules in hank. *Manuscript, January*, pages 121–144.

Schaab, A. and Tan, S. Y. (2023). Monetary and fiscal policy according to hank-io. *Unpublished Manuscript*.

Winberry, T., Auclert, A., Rognlie, M., and Straub, L. (2025). New keynesian economics with household and firm heterogeneity. Technical report, National Bureau of Economic Research.

Yang, Y., Wang, C., Schaab, A., and Moll, B. (2025). Structural reinforcement learning for heterogeneous agent macroeconomics. *arXiv preprint arXiv:2512.18892*.