Introduction to Economic and Social Network

Chen Gao

Personal notes from IESN, Fall 2025, Peking University.

November 01, 2026

Thanks to typst!

Contents

1. Games on networks 5
   1.1. Games on networks 5
   1.2. Complements and substitutes 6
      1.2.1. Complements 6
      1.2.2. Substitutes 7
   1.3. Strategic complements 7
      1.3.1. Lattice structure 8
      1.3.2. Deeper into Proposition 1.3.1 9
   1.4. An application in labor 9
   1.5. Strategic substitutes 10
   1.6. Summary 12
2. TA session 1 13
   2.1. Nodes and networks 13
   2.2. Neighborhood and degree 13
   2.3. Connecting nodes 13
   2.4. Component 14
   2.5. Diameter and average path length 14
   2.6. Centrality measures 15
   2.7. Prestige, influence, eigenvector-based centrality 17
      2.7.1. Intuition 17
      2.7.2. Number of walks from i to j 17
   2.8. Clustering 18
      2.8.1. Individual clustering coefficient 18
      2.8.2. Overall clustering 19
      2.8.3. Average clustering 19
3. Games on networks II 20
   3.1. Strategic substitutes (Bramoullé & Kranton, 2007) 20
   3.2. A linear quadratic model 21
   3.3. Friendship paradox 22
      3.3.1. Empirical observations 25
      3.3.2. Takeaways 26
   3.4. Friendship paradox in (Jackson, 2019) 26
   3.5. Further results 27
      3.5.1. More on (Bramoullé & Kranton, 2007) 27
      3.5.2. (Jackson, 2019) details 28
         3.5.2.1. Linear-quadratic game 28
4. Cooperation on networks 31
   4.1. Sociology of closure (Coleman, 1988) 31
      4.1.1. Clustering coefficients 31
   4.2. Cooperation incentives on networks 31
      4.2.1. Community enforcement 32
   4.3. (Jackson et al., 2012) 32

4.4. Closure and support 35
4.5. Empirical tests (Banerjee et al., 2013) 36
4.6. Multiplexity 37
4.6.1. Multiplex trap and homophily 38
5. Financial networks 40
5.1. (Allen & Gale, 2000) 40
5.2. (Acemoglu et al., 2015) 41
5.3. (Elliott et al., 2014) 41
5.3.1. Basics 42
5.3.2. Model 42
5.3.3. Bankruptcy 43
5.3.4. summary 46
5.4. Eurodebt (Elliott et al., 2014) 46
Bibliography 48
Index of Figures 49

1. Games on networks

1.1. Games on networks

Consider the following game:

- Each player chooses $x_i = 0, 1$
- Payoff depends on
  - how many neighbors choose each action
  - how many neighbors a player has

Consider cases where i's payoff is

$$u_{d_i}\left(x_i, m_{N_i}\right) \tag{1}$$

where m_{N_i} is the number of neighbors of i choosing 1 and $d_i(g)$ is the number of neighbors of i.

Let N_i be the set of i's neighbors. Then $d_i(g) = |N_i|$ and

$$m_{N_i} = \#\left\{j \in N_i, x_j = 1\right\} \tag{2}$$

Example 1.1.1 (Simple Complement): Agent i is willing to choose 1 $\iff$ at least t neighbors do:

- payoff action 0: $u_{d_i}\left(0, m_{N_i}\right) = 0$
- payoff action 1: $u_{d_i}\left(1, m_{N_i}\right) = -t + m_{N_i}$

In Example 1.1.1, the payoff difference is

$$u_{d_i}\left(1, m_{N_i}\right) - u_{d_i}\left(0, m_{N_i}\right) = -t + m_{N_i} \tag{3}$$

. The best response is:

$$x_i^*(m) = \begin{cases} 1 \text{ if } m \geq t \\ 0 \text{ o.w.} \end{cases} \tag{4}$$

Now let $t = 2$:




Figure 1: Equilibrium 1: all choose 0

An agent is willing to take action 1 if and only if at least two neighbors do. And another equilibrium is as in Figure 2:




Figure 2: Equilibrium 2: 3 nodes choose 1

1.2. Complements and substitutes

Definition 1.2.1 (Externalities):
- Others' behaviors affect my utility/welfare
- Others' behaviors affect my decisions, actions, consumptions, opinions...

Others' actions affect the relative payoffs to my actions.

1.2.1. Complements

Choice to take an action by my friends increases my relative payoff to taking that action (e.g., friend learns to play a video game)

Let $m = m_{N_i}$ as defined in Equation 2, then strategic complements means:

Definition 1.2.2 (Complements): for all d[1] and $m \geq m'$, we have

$$u_d(1,m) - u_d(0,m) \geq u_d(1,m') - u_d(0,m') \tag{5}$$

When others choose 1 more often, my incentive to choose 1 weakly increases.

1.2.2. Substitutes

Choice to take an action by my friends decreases my relative payoff to taking that action (e.g., roommate buys a stereo/fridge)

Strategic substitutes satisfy

Definition 1.2.3 (Substitutes): for all $d \wedge m \geq m'$:

$$u_d(1,m) - u_d(0,m) \leq u_d(1,m') - u_d(0,m') \tag{6}$$

Definition 1.2.4 (Nash Eq.): Every player's action is optimal for that player given the actions of others

We often look for pure strategy equilibria and may require some mixing to get existence (hopefully not...)

1.3. Strategic complements

Example 1.3.1 (Examples):
- education decisions
- care about number of neighbors, access to jobs, etc.: invest if at least k neighbors do
- smoking & other behavior among teens, peers, ...
- technology adoption: how many others are compatible
- learn a language, ...
- cheating, doping

One useful observation is that complements $\implies$ there exists a threshold $t(d)$ s.t. i prefers

$$\text{BR}_i = \begin{cases} 1 \text{ if } & m_{N_i} > t(d) \\ 0 \text{ if } & m_{N_i} < t(d) \end{cases} \tag{7}$$

[1] For a given agent i, we suppress i in the subscript.

and can be indifferent at the threshold $t(d)$.

Consider a case where t doesn't depend on d: $t \equiv 2$, then there are multiple equilibria as in Figure 3.




Figure 3: Multiple Equilibria

1.3.1. Lattice structure

The equilibria in Figure 3 have a *lattice structure*.

Definition 1.3.1 (Complete Lattice): For a set of equilibria $\mathbb{X}$,
- there exists an eq. x' s.t. $x' \geq x, \forall x \in \mathbb{X}$

and
- there exists an eq. x'' s.t. $x'' \leq x, \forall x \in \mathbb{X}$

Definition 1.3.2 (Coordinate order): For 2 action allocation: $x = (x_i), y = (y_i)$, we define $x \leq y$ as

$$x \leq y \iff x_i \leq y_i, \forall i = 1, 2, ..., n \tag{8}$$

Using Definition 1.3.2, Definition 1.3.1 gives the following assertions:
1. The maximum value at any given number of equilibrium coordinates is still in equilibrium.
2. The minimum value at any point of any number of equilibrium coordinates is still in equilibrium.
3. In particular, there exists a unique minimum and maximum equilibrium.

Proposition 1.3.1: Given the game of strategic complements where the individual strategy sets are complete lattices, we have, *the set of pure strategy eq. are a nonempty complete lattice*.

Proposition 1.3.1 implies:
1. eq. always exists
2. well-defined maximum and minimum eq.
3. these equilibria are easy to find

1.3.2. Deeper into Proposition 1.3.1

Proposition 1.3.1 requires **the individual strategy sets to be complete lattices**.

Assign a **coordinate-wise/product order** to each player's strategy space. Under this order, the space has a minimum upper bound (join) and a maximum lower bound (meet) for any subset and remains within the space. Equivalently, it contains a global minimum and a global maximum, and is closed with respect to the coordinates at `max/min`.

Formally, let $S_i \subset \mathbb{R}^{k_i}$ be the action space of agent i, and the product order is defined as in Definition 1.3.2. Then S_i is a complete lattice $\iff \forall A \subseteq S_i$, we have

$$\bigvee A \in S_i, \bigwedge A \in S_i \tag{9}$$

that is to take sup or inf at each dimension. Then we know that there are

$$\bot_i \in S_i, \top_i \in S_i \tag{10}$$

which are the minimum and the maximum of S_i.[2]

1.4. An application in labor

Dropout decisions as in (Calvó-Armengol & Jackson, 2004; 2007). Value to being in the labor market depends on # of friends in the market
- Dropout if some # of friends do so
- Some heterogeneity in threshold (different costs, natural abilities...)
- Homophily: segregation in the network

Consider this cascade model as in Figure 4, first, 2 groups exhibit homophily, and secondly, *one agent dropout if at least half of its neighbors do so*.

[2] And therefore usually $\top_i > \bigvee A$ unless $A = S_i$.




Figure 4: Cascade model

As in Figure 4, there are 2 initial dropouts. The cascade ends at Figure 5, where the entire yellow component drops out.




Figure 5: Cascade model: result

The dropout difference across groups persists.

1.5. Strategic substitutes

Similarly, consider the following Example 1.5.1:

Example 1.5.1:

- information gathering
  - e.g., payoff of 1 if anyone in neighborhood is informed, cost to being informed ($c < 1$)
- local public goods (shareable products...)
- competing firms (oligopoly with local markets)

And similar to Equation 7, we have the threshold function $t(d)$ s.t.

$$\mathrm{BR}_i = \begin{cases} 1 \text{ if } m_{N_i} < t(d) \\ 0 \text{ if } \ m_{N_i} > t(d) \end{cases} \tag{11}$$

and can be indifferent at $m_{N_i} = t(d)$.

Example 1.5.2 (Best Shot Game): One is willing to choose 1 $\iff$ no neighbors do. Each node i chooses action $x_i \in \{0, 1\}$ where $x_i = 1$ means providing a public good at cost c. The payoff is

$$u_{d_i}\left(0, m_{N_i}\right) = \begin{cases} 1 \text{ if } m_{N_i} > 0 \\ 0 \text{ o.w.} \end{cases}, u_{d_i}\left(1, m_{N_i}\right) = 1 - c \tag{12}$$

and let $c \in (0, 1)$.

Then for Example 1.5.2, suppose a network in form of Figure 6,




Figure 6: Network structure for Example 1.5.2

It's clear that one will choose $x_i = 1$ if and only if $m_{N_i} = 0$. One eq. is described in Figure 7.




Figure 7: An eq. of Example 1.5.2

Other equilibria are possible.

Consider the best shot game in Example 1.5.2 for a star network, then we have only 2 eq. as in Figure 8. The equilibria have *different utility distributions and total costs*. The eq. is related to a concept called **maximal independent set**.




Figure 8: eq. of star network

Definition 1.5.1 (Maximal Independent Set): Let $M \subseteq N$ be the set of providers:

$$M \equiv \{i \in N, x_i = 1\} \tag{13}$$

1. independent: for all $i, j \in M$, ij $\notin g$
2. maximal: for all $k \notin M$, if we add k to $M \cup \{k\}$, then $M \cup \{k\}$ is not *independent*.

1.6. Summary

In a game of strategic **complements**:
1. Pure strategy equilibria are a nonempty complete lattice

In a game of strategic **substitutes**:
1. Best shot game: pure strategy equilibria exist and are related to maximal independent sets
2. Others: pure strategy may not exist, but mixed will (with finite action spaces)
3. Equilibria usually do not form a lattice

2. TA session 1

2.1. Nodes and networks

Definition 2.1.1 (Nodes): set

$$N = \{1, 2, ..., n\} \tag{14}$$

Definition 2.1.2 (Network): A network (N, g) consists of nodes N and an $n \times n$ adjacency matrix g where:

- $g_{ij} = 1 \implies i$ and j are linked
- A network is undirected iff $g_{ij} = g_{ji}$
- Usually $g_{ii} = 0$

g can also be a collection of links, say, $g = \{12, 23\}$, then $(ij) \in g$ means i and j are linked.

2.2. Neighborhood and degree

Definition 2.2.1 (Neighborhood of i): The set

$$N_i(g) \equiv \left\{ j \in N : g_{ij} = 1 \right\} \tag{15}$$

similarly, degree is then defined as

Definition 2.2.2 (Degree):

$$d_i(g) = \#N_i(g) \tag{16}$$

2.3. Connecting nodes

Definition 2.3.1 (Walk): A walk from i to j is a sequence of nodes $\{i_t\}_{t=1}^{K}$ and sequence of links $\{i_t i_{t+1}\}_{t=1}^{K-1}$ such that

$$i_t i_{t+1} \in g, \forall k \in \{1, 2, ..., K-1\} \tag{17}$$

and $i_1 = i, i_K = j$.

Definition 2.3.2 (Path): A path is a walk with each node in $\{i_t\}_{t=1}^K$ is **distinct.**

Definition 2.3.3 (Cycle): A cycle is a walk $\{i_t\}_{t=1}^K$ w/
1. $i_1 = i_K$
2. all other nodes are distinct

Definition 2.3.4 (Geodesic): The shortest path between 2 nodes.[3]

2.4. Component

Definition 2.4.1 (Connected): A network (N, g) is connected if there is a **Path** between every 2 nodes.

Definition 2.4.2 (Component): A non-empty subnetwork (N', g') such that:$N' \subseteq N, g' \subseteq g, N' \neq \emptyset$ and
- (N', g') is connected
- if $i \in N'$ and $ij \in g$, then $j \in N'$ and $ij \in g'$

Components are the maximal connected subgraphs of a network.

Definition 2.4.3 (Complete network): All possible links are present:

$$g = \mathbf{1}\mathbf{1}^\top - I \tag{18}$$

or equivalently, $g_{ij} = 1$ for all $i \neq j$.

2.5. Diameter and average path length

Definition 2.5.1 (Diameter): The largest geodesic as defined in Definition 2.3.4 between any 2 nodes in the network.

[3] If there is no path between 2 nodes, then the distance between them is ∞.

For a disconnected network, the convention in footnote[3] gives diameter ∞. In practice, we usually report the diameter of the giant component.

For a tree with n nodes, where each node has a branching factor of d links, the diameter (D) is proportional to the logarithm of the number of nodes to the base of the branching factor.

$$D \propto \frac{\log(n)}{\log(d)} \tag{19}$$

Definition 2.5.2 (Average path length): The average geodesic between any 2 nodes in the network, or formally,

$$\sum_{i \geq j} \ell(i,j) \left[\frac{n(n-1)}{2} \right]^{-1} \tag{20}$$

2.6. Centrality measures

Centrality measures the importance of a node. Degree centrality measures *how connected a node is*.

Definition 2.6.1 (Degree Centrality): Let $d_i(g)$ be the degree of node i where

$$d_i(g) = \#\left\{ j \in N, g_{ij} = 1 \right\} \tag{21}$$

or in the normalized form:

$$\frac{1}{n-1} d_i(g) \tag{22}$$

Closeness centrality uses shortest-path distances:

Definition 2.6.2 (Closeness Centrality): Let

$$C_i(g) = \frac{n-1}{\sum_{j=1}^{n} l(i,j)} \tag{23}$$

where $l(i,j)$ is the number of links in the shortest path between i and j.

Definition 2.6.3 (Decay Centrality): Let

$$C_i^d(g) = \sum_{j=1, j\neq i}^{n} \sigma^{l(i,j)} \tag{24}$$

where $l(i,j)$ is defined similar to that of Definition 2.6.2. Its normalized form is

$$\frac{1}{(n-1)\sigma} C_i^d(g) \tag{25}$$

- $\sigma \to 1$, decay centrality measures component size.
  - to see this, note that

$$\begin{aligned} \lim_{\sigma\to 1} C_i^d(g) &= \lim_{\sigma\to 1} \sum_{j=1, j\neq i}^{n} \sigma^{l(i,j)} \\ &= \lim_{\sigma\to 1} \sum_{j=1, j\neq i}^{n} 1^{l(i,j)} \end{aligned} \tag{26}$$

    which is just the size of the component that i is in![4]
- $\sigma \to 0$, decay centrality measures degree .(gives infinitely more weight to closer nodes than farther nodes)
  - That's because

$$\lim_{\sigma\to 0} \frac{\sigma^1}{(n-1)\sigma} = \frac{1}{n-1}, \lim_{\sigma\to 0} \frac{\sigma^l}{(n-1)\sigma} = 0, \forall l \geq 2, l \in \mathbb{N} \tag{27}$$

- σ in between, decaying distance measure.(very distant nodes are weighted less than closer nodes)

Betweenness centrality measures *how much a node connects other nodes*.

Definition 2.6.4 (Betweenness Centrality): Let

$$C_k^B(g) = \frac{\sum_{i<j, i\neq k} \frac{P_k(ij)}{P(ij)}}{\binom{n-1}{2}} = \frac{2\sum_{i<j, i\neq k} \frac{P_k(ij)}{P(ij)}}{(n-1)(n-2)} \tag{28}$$

be the betweenness centrality of node k[5], where
1. $P_k(ij)$ denotes the total number of geodesics between i, j that k lies on.
2. $P(ij)$ is the total number of geodesics between i, j

[4] However, this is certainly not $d_i(g)$.

[5] The normalization $\binom{n-1}{2}$ counts pairs among the other $n-1$ nodes, whose geodesics may pass through this node.

2.7. Prestige, influence, eigenvector-based centrality

Eigenvector centrality weights connections by the centrality of neighboring nodes. A node connected to many central nodes also has high centrality.

Definition 2.7.1 (Eigenvector Centrality): Given a network (N, g), we define the eigenvector centrality of a node i as:

$$C_i^e(g) = \alpha \sum_{j \in N} g_{ij} C_j^e(g) \tag{29}$$

and in vector form:

$$C^e = \alpha g C^e \tag{30}$$

where $C^e(g) \in \mathbb{R}^n$.

Rearrange Equation 30 as $\frac{1}{\alpha} C^e = g C^e$. Setting $\lambda = \frac{1}{\alpha}$ gives the eigenvalue equation for g. We seek the eigenvector associated with the **largest** eigenvalue, which is nonnegative.

Katz-Bonacich centrality also includes indirect connections.

2.7.1. Intuition

Each node has a base value and receives additional value through direct and indirect connections.

2.7.2. Number of walks from i to j

Consider a network (N, g), then g^n indicates how many walks there are of length n between 2 nodes, see Example 2.7.1.

Example 2.7.1: Let $g^2 = \begin{pmatrix} 2 & 0 & 0 & 2 \\ 0 & 2 & 2 & 0 \\ 0 & 2 & 2 & 0 \\ 2 & 0 & 0 & 2 \end{pmatrix}$, then $g_{14} = 2$ means that there are 2 walks between $1, 4$ of length 2.

Give each node an initial value $\alpha d_i(g)$ for some $\alpha > 0$.

1. we add in all walks of length 1 from j to i w/

$$\sum_{j:ij \in g} \beta \alpha d_j(g) \tag{31}$$

2. then we add in all walks of length 2:

$$\sum_{j:ij \in g} \beta^2 \alpha d_j(g) \tag{32}$$

3. ...
4. finally it looks like

$$
\begin{aligned}
C^b(g) &= \alpha g\mathbf{1} + \alpha\beta g^2\mathbf{1} + \dots \\
&= \alpha(1 + \beta g + \beta^2 g^2 + \dots)g\mathbf{1} \\
&= \alpha(I - \beta g)^{-1} g\mathbf{1}
\end{aligned} \tag{33}
$$

and note that $g\mathbf{1} \in \mathbb{R}^{n\times 1}$ is the degree of each node.

- β captures how the value of being connected to someone decays with distance.
- α captures the base value on each node.

2.8. Clustering

"If i has relationships with both j and k, how likely is it that j and k are related ?"

This problem is central to clustering. In sociology, this is known as a **triadic closure**: If A knows B and A knows C, then B and C are very likely to know each other as well.

2.8.1. Individual clustering coefficient

Definition 2.8.1 (Individual Clustering Coefficient): Let

$$
Cl_i(g) = \frac{\#\{jk \in g | j,k \in N_i(g)\}}{\#\{jk \mid j,k \in N_i(g)\}} \tag{34}
$$

be the Individual Clustering Coefficient of node i.

The number of neighbor pairs is

$$
\#\{jk \mid j,k \in N_i(g)\} = \binom{d_i(g)}{2} \tag{35}
$$

1. In the Erdős–Rényi (E-R) random network $G(n,p)$, the probability of a connection between any two nodes is p. *Since the links in the E-R model are formed independently, the probability of a connection between two neighbors j and k of node i is the same as the probability of any connection in the network p* and therefore

   $$
   Cl_i(g) = p. \tag{36}
   $$

   This indicates that there is no clustering phenomenon exceeding the random level in the E-R model.
2. If the degree of node i is $d_i(g) \leq 1$ (i.e., it has 0 or 1 links), then it has insufficient neighbors to form any pair of neighbors. Then

   $$
   Cl_i(g) = 0 \tag{37}
   $$

2.8.2. Overall clustering

Definition 2.8.2 (Overall clustering): Let

$$Cl(g) = \frac{\sum_{i \in N} \#\{jk \in g | \, j,k \in N_i(g)\}}{\sum_{i \in N} \#\{jk \mid j,k \in N_i(g)\}} \tag{38}$$

be the overall clustering coef. of a network (N, g).

Each pair of friends receives equal weight, so high-degree nodes receive more weight.

1. High-degree nodes can form more triples, open or closed, so their clustering contributes more to the final $Cl(g)$ value.
2. "Equal weight for each friend" means that each "triple" contributes the same amount when calculating the ratio.

2.8.3. Average clustering

Definition 2.8.3 (Average clustering): Let

$$Cl^{\text{avg}}(g) = \frac{1}{n} \sum_{i=1}^{n} Cl_i(g) \tag{39}$$

be the average clustering coefficient of a network (N, g).

Equal weight for every node.

3. Games on networks II

3.1. Strategic substitutes (Bramoullé & Kranton, 2007)

You need to acquire information for an application, the payoff is

$$f\left(x_i + \sum_{j \in N_i(g)} x_j\right) - cx_i \tag{40}$$

where f is a concave "production function", FOC gives

$$f'(x^*) = c \tag{41}$$

So in all pure strategy Nash Equilibria, we have the optimization:

$$\begin{aligned} \max_{x_i} \quad & f\left(x_i + \sum_{j \in N_i(g)} x_j\right) - cx_i \\ \text{s.t.} \quad & x_i \geq 0 \end{aligned} \tag{42}$$

then KKT condition of Equation 42 gives:

$$\begin{aligned} f'\left(x_i + \sum_{j \in N_i(g)} x_j\right) &= c - \lambda_i \\ \lambda_i &\geq 0 \\ x_i &\geq 0 \\ \lambda_i x_i &= 0 \end{aligned} \tag{43}$$

and that is

$$\begin{aligned} x_i + \sum_{j \in N_i(g)} x_j &\geq f'^{-1}(c) \\ x_i = 0 \text{ if } x_i + \sum_{j \in N_i(g)} x_j &> f'^{-1}(c) \end{aligned} \tag{44}$$

There are 2 types of equilibria:

- Distributed eq.

$$x_i \in \left(0, f'^{-1}(c)\right) \tag{45}$$

  for some i's
- Specialized eq. for each i, either $x_i = 0$ or $x_i = f'^{-1}(c)$

We consider the case where $x^* = 1$, that is, everyone wants a total investment of at least 1 in her neighborhood. The equilibria is described in Figure 9.




Figure 9: Eq. of BK 07

The specialized equilibria can be characterized using Definition 1.5.1 as follows:

Proposition 3.1.1: Specialized Nash equilibria are such that a maximal independent set = the specialist set

$$\{i : i \in N, x_i = f'^{-1}(c)\} \tag{46}$$

3.2. A linear quadratic model

This comes from (Ballester et al., 2006), we set the payoff function as:

$$u(x_i, x_{-i}) = ax_i - 0.5bx_i^2 + \sum_j w_{ij}x_i x_j \tag{47}$$

and we set quadratic intrinsic value to be $v(x_i) = ax_i - 0.5bx_i^2$. This is also Strategic Complements since

$$\frac{\partial u}{\partial x_i} = a - bx_i + \sum_j w_{ij}x_j \tag{48}$$

so when x_j increases, $\frac{\partial u}{\partial x_i}$ also increases.

To solve the best response of x_i given x_{-i}, use the first-order condition:

$$a - bx_i + \sum_j w_{ij}x_j = 0 \tag{49}$$

so

$$x_i^* = \frac{1}{b}\left(a + \sum_j w_{ij}x_j\right) \tag{50}$$

consider the matrix form where $\boldsymbol{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$ and $\boldsymbol{\alpha} = \frac{a}{b}\mathbf{1}_n, \boldsymbol{g} = \left(g_{ij}\right)_{n\times n}$ where $g_{ij} = \frac{w_{ij}}{b}$, then we can rewrite Equation 49 in matrix form:

$$\boldsymbol{x} = \boldsymbol{\alpha} + \boldsymbol{g}\boldsymbol{x} \tag{51}$$

and then by simple algebra we have

$$\boldsymbol{x} = \sum_{k\geq 0}^{\infty} \boldsymbol{g}^k \boldsymbol{\alpha} \tag{52}$$

and if $(\boldsymbol{I} - \boldsymbol{g})$ is invertible then we have directly

$$\boldsymbol{x} = (\boldsymbol{I} - \boldsymbol{g})^{-1}\boldsymbol{\alpha} \tag{53}$$

The model is based on **Strategic Complements**:
- If my neighbors work harder ($x_j \uparrow$), my marginal benefit increases.
- I respond by working harder ($x_i \uparrow$).
- This causes my neighbors to work even harder in response.

Positive feedback can make equilibrium actions unbounded ($x_i \to \infty$).

A unique finite Nash equilibrium requires sufficiently high action costs relative to network benefits: **large** b and/or **small** w_{ij}.

- w_{ij}**:** Strength of network complementarity. Large w strengthens the feedback loop.
- b**:** Curvature of the cost function (diminishing returns).
  - Large b makes marginal costs rise steeply and limits increases in x_i.

Recall Katz - Bonacich centrality:

$$B(g) = (I - g)^{-1} g\mathbf{1} = \sum_{k\geq 0} g^{k+1}\mathbf{1} \tag{54}$$

so we have

$$x \propto B(g) \tag{55}$$

1. Natural feedback from complementarities, actions relate to the total feedback from various positions
2. Related to centrality: relative number of weighted influences going from one node to another

3.3. Friendship paradox

Example 3.3.1 (Friendship Paradox)**:** Who is more popular? You, or your friends? That is $\frac{1}{n}\sum_{i\in N} d_i(g)$ vs. $\frac{1}{n}\sum_{i\in N}\left(\frac{1}{d_i(g)}\sum_{j\in N_i(g)} d_j(g)\right)$.

We try to solve this rigorously :-)

For the LHS, write

$$\mu = \mathbb{E}[d_i(g)] = \frac{1}{n}\sum_{i\in N} d_i(g) \tag{56}$$

and for the RHS, we should consider the probability of a friend j to be selected. We know that in this case: $P(j) \propto d_j(g)$, and by regularity, we have $\sum_{j\in N, d_j(g)\geq 1} P(j) = 1$, so we have

$$P(j) = \frac{d_j(g)}{\sum_{k\in N, d_k(g)\geq 1} d_k} = \frac{d_j(g)}{\sum_{k\in N} d_k} \tag{57}$$

so we have the RHS as:

$$\begin{aligned}
\frac{1}{n}\sum_{i\in N}\left(\frac{1}{d_i(g)}\sum_{j\in N_i(g)} d_j(g)\right) &= \sum_{j\in N, d_j(g)\geq 1} d_j(g)P(j) \\
&= \sum_{j\in N, d_j(g)\geq 1} d_j(g)\frac{d_j(g)}{\sum_{k\in N} d_k} \\
&= \frac{1}{\sum_{k\in N} d_k}\sum_{j\in N} d_j^2(g) \\
&= \frac{1}{n\mu}\sum_{j\in N} d_j^2(g)
\end{aligned} \tag{58}$$

consider the variance, we have

$$\mathbb{V}[d_i(g)] = \mathbb{E}\left[d_i^2(g)\right] - \left(\mathbb{E}[d_i(g)]\right)^2 \tag{59}$$

so

$$\begin{aligned}
\mathbb{E}\left[d_i^2(g)\right] &= \mathbb{V}[d_i(g)] + \left(\mathbb{E}[d_i(g)]\right)^2 \\
&= \sigma^2 + \mu^2
\end{aligned} \tag{60}$$

and

$$\text{RHS} = \frac{1}{\mu}\mathbb{E}\left[d_i^2(g)\right] = \frac{1}{\mu}(\sigma^2 + \mu^2) = \frac{\sigma^2}{\mu} + \mu > \mu = \text{LHS} \tag{61}$$

which proves the result.

Example 3.3.2 (The Impact of the Friendship Paradox): Consider a 0-1 game...
- People choose solid or plaid
- Prefer to match the majority of your friends
- Go with own preference if indifferent

People typically have fewer friends than their friends

1. People with more friends are 'seen' more
2. Popular people are over-represented in the views of others

Start from the following state:




Figure 10: Own preferences

Note that in this starting state, only the influencers prefer the *solid* ones. Then each node tries to match the majority of its friends. The resulting equilibrium is:




Figure 11: Eq. of this shirt game

That is, all nodes will choose solid in the end.

In this example of Example 3.3.1, the social average *preference for solid shirts* is $\frac{4}{12} = \frac{1}{3}$.

Each node's perceived average *preference for solid shirts* is shown below:




Figure 12: Perceived averages

For 9 of the 12 agents, the perceived average exceeds the social average.

The friendship paradox makes socially active agents overrepresented among observed friends. Their behavior can then appear more common than it is in the population, the *majority illusion*.

3.3.1. Empirical observations

1. Systematic Overestimation (Perkins et al., 2005)

TABLE 1. Students' accuracy in perceiving the drinking norm at their school (comparing actual with perceived number of alcoholic drinks consumed the last time students "partied"/socialized)

Actual school norm (median drinks)	Accuracy of perceived drinking norm: Underestimate of ≥3 drinks (%)	Underestimate of 1-2 drinks (%)	Accurate estimate (%)	Overestimate of 1-2 drinks (%)	Overestimate of ≥3 drinks (%)	Total	*N* of respondents	*N* of schools
0	NA	NA	20.6	19.5	59.9	=100%	1,891	4
1	NA	10.5	3.8	28.5	57.2	=100%	2,526	6
2	NA	7.5	8.1	30.6	53.7	=100%	8,345	14
3	3.8	6.4	13.5	37.5	38.7	=100%	18,859	35
4	3.1	12.3	12.6	37.0	34.9	=100%	20,353	38
5	4.3	15.8	20.6	24.1	35.3	=100%	11,481	20
6	6.9	23.2	15.0	23.5	31.5	=100%	8,912	12
7	5.7	23.3	9.7	23.6	37.8	=100%	352	1
Total schools	3.4	11.8	13.8	31.9	39.1	=100%	72,719	130

   The vast majority of people (31.9% + 39.1% ≈ 71%) seriously overestimated their classmates' alcohol consumption.
2. Positive Correlation(degree, behavior) For middle school students (12-15 years old)...
   1. The probability of smoking increases by 5% for every additional friend you have. (Valente et al., 2005)
   2. For every additional friend you have, the likelihood of trying alcohol increases by 6%.(Tucker et al., 2013)

Those "high-degree agents" who are at the center of the network are often the ones who smoke and drink the most.

3.3.2. Takeaways

1. The friendship paradox overrepresents people with many friends in observations of the network.
2. The empirical patterns above show more drinking and smoking among these highly visible agents.

Strategic complementarity can account for the second pattern:
1. People in central positions (high degree/centrality) are most influenced by their friends (positive feedback loop), or they must exert more effort (action) to maintain their central position.
2. Maintaining a popular position and responding to peers can lead central individuals to drink or smoke more.

(Jackson, 2019) endogenizes this relationship.

3.4. Friendship paradox in (Jackson, 2019)

Consider the following model:
- there are n players, each with intrinsic preference θ_i
- each one chooses an action x_i
- in a random network
- Let $P_i(d)$ denote the distribution of population of i's potential neighbors' degrees.

Lemma 3.4.1 (Expected Degree of Neighbors): The distribution of i's neighbors' expected degrees is given by

$$P_i(d)\frac{d}{\mathbb{E}_i[d]} = P_i(d)\frac{d}{\sum_d dP_i(d)} \tag{62}$$

, so the expected degree of i's neighbors is:

$$\hat{\mathbb{E}}_i[d] = \sum_d dP_i(d)\frac{d}{\mathbb{E}_i[d]} = \frac{1}{\mathbb{E}_i[d]}\mathbb{E}_i\left[d^2\right] = \mathbb{E}_i[d] + \frac{\mathbb{V}_i[d]}{\mathbb{E}_i[d]} > \mathbb{E}_i[d] \tag{63}$$

Proposition 3.4.1: In a game with strategic complements, we have

$$x_i^{\text{prec}}(\theta_i, d_i) \geq x_i^{\text{soc}}(\theta_i, d_i), \forall \theta_i, d_i \tag{64}$$

and

$$\underbrace{\hat{\mathbb{E}}\left[x_i^{\text{prec}}(\theta_i, d_i)\right] \geq}_{\text{network perception effect}} \underbrace{\mathbb{E}\left[x_i^{\text{prec}}(\theta_i, d_i)\right] \geq \mathbb{E}[x_i^{\text{soc}}(\theta_i, d_i)]}_{\text{network eq. effect}} \tag{65}$$

In Equation 64, overestimating others' actions raises one's own action through strategic complementarity.

For Equation 65,
1. network perception effect: *You believe others are doing more than they actually are.*
2. network eq. effect: *Strategic complementarity allows you to do more in eq.*

In Proposition 3.4.1, central nodes are overrepresented among observed neighbors. With strategic complementarity, the resulting overestimate of others' actions raises equilibrium actions.

The welfare effect depends on the externalities.

3.5. Further results

3.5.1. More on (Bramoullé & Kranton, 2007)

1. Any maximal independent set is an eq.
2. How to select one?

- idea: Stability as Equilibrium Selection

A stable equilibrium is restored after small perturbations. A small perturbation of an unstable equilibrium can lead to a different equilibrium.

Consider an eq. x, and

$$x^0 = \begin{pmatrix} x_1 + \varepsilon_1 \\ x_2 + \varepsilon_2 \\ \vdots \\ x_n + \varepsilon_n \end{pmatrix} \tag{66}$$


where ε_i is a small perturbation.

Then let x^1 be the best response to x^0, and iterate forward:


$$x^t = \text{BR}\left(x^{t-1}\right) \tag{67}$$

and if after T periods the whole system goes back to x then we say it's stable.

So in the BK model, we can find that the *specialist equilibria* is stable, that is



This equilibrium is stable, whereas the following equilibria



are unstable. The proposition is:

Proposition 3.5.1: Only stable equilibria are specialist equilibria such that every non-specialist has at least two specialists in his or her neighborhood

1. If there is only one specialist, when the specialist's level drops slightly, the free-rider will start providing its own services.
2. If there are at least 2 specialists, even with minor perturbations:
   1. The free-rider still sees a neighbor that is working hard enough.
   2. It continues free-riding (remaining at 0).
   3. The specialist will also remain at 1 under the best-response condition.
   4. The system returns to the equilibrium, so it is stable.

3.5.2. (Jackson, 2019) details

3.5.2.1. Linear-quadratic game

The social-benchmark expected utility is:

$$\mathbb{E}[U_i^{\text{soc}}(x_i, x_{-i})] = \theta_i x_i + a x_i d_i \mathbb{E}\left[x_j\right] - \frac{c}{2} x_i^2 + \varphi \mathbb{E}\left[\sum_{j \neq i} x_j\right] \tag{68}$$

Its components are:

1. $\theta_i x_i$: the intrinsic preference
2. $a x_i d_i \mathbb{E}\left[x_j\right]$: the strategic complementarity gain where $\mathbb{E}\left[x_j\right]$ is the average action of i's neighbors
3. $-\frac{c}{2} x_i^2$: quadratic cost
4. $\varphi \mathbb{E}\left[\sum_{j \neq i} x_j\right]$: *Under the social benchmark (SOC), the planner is concerned with the overall level of behavior, not just the utility function of individuals.* That is, Your friends also generate positive externalities for others (not you), so social planners calculate the sum of everyone's actions when optimizing.

Perceived expected utility is:

$$\begin{aligned}\mathbb{E}\Big[U_i^{\text{prec}}(x_i, x_{-i})\Big] &= \theta_i x_i + a x_i d_i \hat{\mathbb{E}}\Big[x_j\Big] - \frac{c}{2}x_i^2 + \varphi \mathbb{E}\left[\sum_{j\neq i} x_j\right] \\ &= \mathbb{E}[U_i^{\text{soc}}(x_i, x_{-i})] + a x_i d_i \Big(\hat{\mathbb{E}}\Big[x_j\Big] - \mathbb{E}\Big[x_j\Big]\Big)\end{aligned} \tag{69}$$

Individuals overestimate their neighbors' actions.

And each agent i solves:

$$\max_{x_i} \quad \theta_i x_i + a x_i d_i \hat{\mathbb{E}}\Big[x_j\Big] - \frac{c}{2}x_i^2 \tag{70}$$

and the best response is:

$$x_i^* = \frac{1}{c}\Big(\theta_i + a d_i \hat{\mathbb{E}}\Big[x_j\Big]\Big) \tag{71}$$

which gives Equation 64.

Lemma 3.5.1: We assume the following holds:

1. $\hat{\mathbb{E}}_i[d] = \hat{\mathbb{E}}[d]$, that is *Everyone has the same perception of the average degree of their neighbors.*
2. and $c > a\hat{\mathbb{E}}[d]$: *The cost must be high enough relative to complementarity for equilibrium to be finite and unique.*

then there exists a unique eq.:

$$x_i^{\text{prec}}(\theta_i, d_i) = \frac{\theta_i}{c} + \frac{a d_i \hat{\mathbb{E}}[\theta]}{c^2 - ca\hat{\mathbb{E}}[d]} \tag{72}$$

To derive Lemma 3.5.1, start from the best response in Equation 71 and take expectations:

$$\begin{aligned} x_i &= \frac{1}{c}(\theta_i + a d_i \hat{\mathbb{E}}[x]) \\ \hat{\mathbb{E}}[x] &= \hat{\mathbb{E}}\left[\frac{1}{c}(\theta_i + a d_i \hat{\mathbb{E}}[x])\right] \\ \hat{\mathbb{E}}[x] &= \frac{1}{c}\big[\hat{\mathbb{E}}[\theta] + a\hat{\mathbb{E}}[d]\hat{\mathbb{E}}[x]\big] \end{aligned} \tag{73}$$

then we can get $\hat{\mathbb{E}}[x]$ and then plug it into the Equation 71 to get Equation 72.

So it's *higher actions for higher types, higher degree*. Similarly, we have

$$x_i^{\text{soc}}(\theta_i, d_i) = \frac{\theta_i}{c} + \frac{a d_i \mathbb{E}[\theta]}{c^2 - ca\mathbb{E}[d]} \tag{74}$$

so by Lemma 3.4.1, Equation 64 holds.

The degree distributions can also be compared by stochastic dominance.

Definition 3.5.1 (First-Order Stochastic Dominance): Let the CDF of X and Y be $F_X(x) = \Pr\{X \leq x\}$ and $F_{Y(y)} = \Pr\{Y \leq y\}$, then we say $X \underset{\text{FOSD}}{\succ} Y$ iff

$$F_X(x) \leq F_Y(x), \forall x \in \mathbb{X} \tag{75}$$

and $<$ holds strictly for some x.

Then by definition

$$\hat{P}_i(d) \equiv P_i(d)\frac{d}{\mathbb{E}[d]} \underset{\text{FOSD}}{\succ} P_i(d) \tag{76}$$

and therefore

$$\hat{\mathbb{E}}[d] > \mathbb{E}[d] \tag{77}$$

4. Cooperation on networks

4.1. Sociology of closure (Coleman, 1988)

When people in a social network know each other (closed structure), it is easier for them to form common norms, build trust, and restrain an individual's bad behavior through collective pressure.




FIG. 1.—Network without (*a*) and with (*b*) closure

Definition 4.1.1 (Collective sanction): Collective sanction refers to the collective pressure, punishment, or constraint exerted on an individual by members of a group in order to establish norms.

4.1.1. Clustering coefficients

1. Prison friendships
   - .31 (MacRae 60) vs .0134
2. Co-authorships
   - .15 math (Grossman 02) vs .00002,
   - .09 biology (Newman 01) vs .00001,
   - .19 econ (Goyal, van der Leij, Moraga 06) vs .00002,
3. Web
   - .11 for web links (Adamic 99) vs .0002

These networks have high clustering: a friend's friend is often a friend. Closure helps sustain norms, trust, and collective sanctions.

4.2. Cooperation incentives on networks

The game is called *Favor exchange*

1. cost of giving a favor: c
2. net benefit per period is v
3. total net benefit with discount factor δ:

$$V = \sum_{t=1}^{\infty} \delta^t v = \frac{\delta}{1-\delta} v \tag{78}$$

Self-enforcing This refers to cooperation that can be maintained solely through mutual interests, without the need for police or laws.

The incentive condition is:

$$c < V = \frac{\delta}{1-\delta} v \tag{79}$$

4.2.1. Community enforcement

So if it's a one to one relation ship, we need $c < V = \frac{\delta}{1-\delta} v$,but if we have a network like this:




Figure 16: Community-enforcement

In Figure 16, we have one mechanism: **Ostracize deviating agent**

If the green dot does something bad (such as refusing to help one of the white dots). Because the two white dots are friends, they share information. If one of the green dots betrays the other, **both white dots will punish the green dot simultaneously** (for example, by severing their relationship with it).

So in this case, we have


$$c < 2 \times V = 2 \times \frac{\delta}{1-\delta} v \tag{80}$$

4.3. (Jackson et al., 2012)

Ostracism without cost?

If a traitor is kicked out as punishment, the remaining group may be **too small to maintain cooperation**, ultimately causing the entire network to collapse.

Suppose $V < c < 2V$:




In this case, first 1 deviates against 2, then *1 is ostracized*. But when 3 deviates against 1, she will find that maintain the relationship with 2 is no longer profitable, so the entire network collapses.

Therefore, this threat of punishment is non-credible.


Formally, at period t:

1. At most one agent i_t is called upon to perform a favor for $j_t \in N_i(g_t)$
2. i_t keeps or deletes the link
3. others can respond: announce which (remaining) links they wish to maintain
4. links are retained iff mutually agree
5. the resulting network is g_{t+1}


We consider 2 structures:




For the left structure, if the there is one disconnection:



The disconnection eventually isolates all nodes:



that is, **autarchy**. But for the right structure, suppose the same split happens, it will end up as:




It loses a triad, preserving more links than the left network's autarchy outcome.

Definition 4.3.1 (Robustness Against Contagion): A network such that the punishment for failing to perform a favor only impacts neighbors of original players lose links

Deleting a node or experiencing a disturbance (default) has a *localized* impact.

Definition 4.3.2 (Supported Link): Link $ij \in g$ is supported if there exists k such that $ik \in g, jk \in g$.

That is, i, j, k forms a triangle.



The following theorem gives a necessary condition for robustness:

Theorem 4.3.1: If no pair of players could sustain favor exchange in isolation and a network is robust, then all of its links are supported.

Supported: In online games, this usually means that the existence of an edge is not only due to the relationship between two people, but is also monitored and supported by a "third party" or "mutual friend" (e.g., if A betrays B, C will also punish A).

4.4. Closure and support

1. Closure (Coleman, 1988)
2. Support (Jackson et al., 2012)

For closure, we need to ask a question as in Figure 22, that is w/ what frequency are a typical node A's neighbors, say B and C, neighbors of each other?




Figure 22: closure

And for support, as in Figure 23, we wonder w/ what frequency do a typical pair of connected nodes, say A and B, have a common neighbor C?





Figure 23: Support


4.5. Empirical tests (Banerjee et al., 2013)

1. Researchers surveyed 75 rural villages in Karnataka, India.
2. A microfinance agency (BSS) entered 43 of these villages.
3. Before entry, researchers mapped villagers' networks and distinguished two types of links:
   1. "**Favor**" **Networks**: These involve the exchange of favors, such as lending money, rice, or kerosene.[6]
   2. "**Social**" **Networks**: Purely for chatting, visiting others' spaces, and maintaining friendships. (Low risk).



- Support: *This means that if two people have a loan relationship (favor link) in these villages, there is an 80%-90% probability that they have at least one mutual friend.*
- Clustering: *This means that two friends of one person do not necessarily know each other.*

[6]Note: These relationships are risky; the money may not be repaid, so theoretically, "Support" is necessary.



Mutual friends strongly predict lending links: shared neighbors help enforce repayment.

Network formation for lending

- Because we need to lend and borrow from each other and are afraid that the other party will default (demand),
- we deliberately build a "triangle" relationship network (structure) with a large number of mutual friends.
- Support measures this structure, which resembles a patchwork of triangles.

4.6. Multiplexity

Multiplex to better sustain cooperation

- prisoner's dilemma: one may deviate against a partner
  - like not return money borrowed
- incentives to prevent such deviations:
  - potential interactions in future on the same link
  - punishments from other (common) friends
- **multiplex**: deviation on one relation is punished by other relations





Figure 26: 3 types of networks

multiplex and community enforcement both enhance cooperation
- multiplex is more robust: rely less on the rest of the network

In Figure 26, we have
- left: multiplex
- middle: community enforcement
- right: star network w/o support

4.6.1. Multiplex trap and homophily

Consider the following network:



so there are 2 types of relations:
1. kid care: better on one side
2. risk share: better across the river

For the green node, she will link kid care w/ the lower right node
1. then she'll decide the risk-share link
2. it's better to link the upper left one
3. but w/ multiplexity, she'll feel it better to link w/ the same lower right node:

In the end, we have the eq. as in the left of Figure 28.
- homophily expands from one relation to other relations
- so called. **multiplex trap**




Figure 28: Eq. and efficiency

The most efficient outcome on the right of Figure 28 is not an equilibrium.

5. Financial networks

5.1. (Allen & Gale, 2000)

- Viewed as a starting point
- An arrow: one holds deposits in another bank




FIG. 1.—Complete market structure




FIG. 2.—Incomplete market structure

The left one is viewed as "complete" and the right one is "incomplete".

Consider a shock to A and makes it bankrupt:

- in the right figure:

$$D \to C \to B \tag{81}$$

- in the left figure: equally shared by $B, C, D \implies$ More likely they all survive.




FIG. 3.—Disconnected incomplete market structure

Consider a shock that makes A bankrupt:

- only affect B
- less fragile than fig 2

5.2. (Acemoglu et al., 2015)




FIGURE 1. THE RING AND THE COMPLETE FINANCIAL NETWORKS

- Arrow: debt contracts

For a **small** negative shock, the ring (left one) is fragile.
- more like a ring: more fragile
- more complete: more stable

For a **large** negative shock:
- **Ring and complete networks are the least stable ones**

but once we have a separation "δ-connected":
- when δ is small, this is more stable than ring or complete.

Phase Transition: under small shocks vs. large shocks, the more stable network structures can be very different

- Pros:
  - generalization to (Allen & Gale, 2000)
  - debt contracts: widely accepted by macro/finance audience
- Cons:
  - model not very tractable
  - limited lessons regarding network structures

5.3. (Elliott et al., 2014)

- develop network-based measures of cascades
- distinguish the effects of *diversification* and *integration*.
- show non-monotonic effects of diversification and integration on contagion.
- illustrate an empirical application of the model.

5.3.1. Basics

- organizations have claims on
  - fundamental assets
  - other organizations
- Bankruptcy cost:
  - when an organization's value falls below a level, the value of other's claims on its drop discontinuously
  - drop in value of one leads to drop in values of others: **cascades**

5.3.2. Model

- organizations: $\{1, 2, ..., n\}$
- Assets: $\{1, 2, ..., m\}$
- let price of asset k be p_k
- and let D_{ik} be the holdings of asset k by organization i

Value of an organization:

- book value:

$$V_i = \sum_{k=1}^{m} D_{ik} p_k + \sum_{j=1}^{n} C_{ij} V_j \tag{82}$$

  - where $\sum_{k=1}^{m} D_{ik} p_k$ is direct asset holdings
  - and $\sum_{j=1}^{n} C_{ij} V_j$ is cross holdings

In vector form:

$$V = Dp + CV \tag{83}$$

that is,

$$V = (I - C)^{-1} Dp \tag{84}$$

- Market Value:
  - defined as value to private investors
  - let

$$v_i = \hat{C}_{ii} V_i \tag{85}$$

  - that is:

$$v = \underbrace{\hat{C}(I - C)^{-1}}_{A} Dp \tag{86}$$

  - and then A_{ij} denotes: *fraction of the returns owned by org j that ultimately accrue to private shareholders of i*

Example 5.3.1: Let there be $n = 2$ firms, each owns half of the other:

$$C = \begin{pmatrix} 0 & 0.5 \\ 0.5 & 0 \end{pmatrix} \tag{87}$$

then it must own half of itself:

$$\hat{C} = \begin{pmatrix} 0.5 & 0 \\ 0 & 0.5 \end{pmatrix} \tag{88}$$

and then

$$\begin{aligned} A = \hat{C}(I - C)^{-1} &= \begin{pmatrix} 0.5 & 0 \\ 0 & 0.5 \end{pmatrix} \begin{pmatrix} 1 & -0.5 \\ -0.5 & 1 \end{pmatrix}^{-1} \\ &= \begin{pmatrix} 0.5 & 0 \\ 0 & 0.5 \end{pmatrix} \begin{pmatrix} \frac{4}{3} & \frac{2}{3} \\ \frac{2}{3} & \frac{4}{3} \end{pmatrix} \\ &= \begin{pmatrix} \frac{2}{3} & \frac{1}{3} \\ \frac{1}{3} & \frac{2}{3} \end{pmatrix} \end{aligned} \tag{89}$$

and this is depicted as Figure 30




Figure 30: cascade of 2 firms

5.3.3. Bankruptcy

if an org's value drops below some *threshold* θ:

- its value falls discontinuously by β_i
- let the threshold be $\underline{v}_i$, then

$$b_i(v_i) = \begin{cases} \beta_i & \text{if } v_i < \underline{v}_i \\ 0 & \text{o.w.} \end{cases} \tag{90}$$

w/ bankruptcy, we have

$$v = A[Dp - b(v)] \tag{91}$$

Solution

- let $s_i = 1$ if an org. fails
  - $s_i = 0$ if not
- $s = (s_1, ..., s_n)$ is an equilibrium if

$$v = A[Dp - \text{diag}(\beta)s] \tag{92}$$

and

$$v_i < \underline{v}_i \iff s_i = 1 \tag{93}$$

that is

$$s(v) = \mathbb{I}(v < \underline{v}) \in \mathbb{R}^{n \times 1} \tag{94}$$

Equilibria

- Bankruptcies are complements:
  - one fails → others likely to fail
  - the set of equilibria forms a complete lattice
- exists a "base case" eq.
  - fewest orgs fail
- algorithm to find it:
  - identify orgs that fail even all others success
  - then identify failures due to the failure of the first org.

simulation

- one asset per org starting at value 1
- shock is to pick one org.'s direct investment to devalue to 0
- then check if it fails according to Section 5.3.3
- look at the result cascade

Let there be $n = 100$ org. and form a random network w/

$$\Pr\{g_{ij} = 1\} = \frac{d}{n-1} \tag{95}$$

- where d is the expected number of other org. that an org. cross holds
- and d is therefore *level of diversification*

Then we let c fraction of org cross-hold:

- so $1 - c$ holds privately

Then claim i has on j:

$$C_{ij} = c\frac{g_{ij}}{d_j} \tag{96}$$




Figure 31: diversification and contagion

The x-axis of Figure 31 is the degree and the y-axis of it is the percent of orgs that fail. different lines reports different threshold θ.

A cascade

Three necessary components of it:
1. a first failure
2. contagion: that is, other org. must be exposed to the failed org. sufficiently to fail too
3. wider propagation: sufficiently large components

Diversification

- has non-monotonicity
  - increases component size
  - decrease the impact of one organization on its neighbors

Intermediate diversification can produce widespread contagion:
- low diversification:
  - fragmented network
  - no widespread contagion
- high
  - little exposure to any single other org.
  - failures do not spread

however, in the middle:
- connected, contagion is possible
- exposure to only a few others makes it easy to spread.

Integration




Figure 32: integration

Here the x-axis is the degree: expected number of cross holdings.

- Low:
  - little exposure to others,
  - failures don't trigger others
- high:
  - difficult to get a first failure
  - failure of own assets doesn't trigger failure
- middle:
  - exposure to others enough to trigger contagion

5.3.4. summary

- a first failure:
  - some org. needs to fail
  - *integration decreases own-asset dependence*
- initial contagion:
  - some neighbors need to be sufficiently exposed to fail too

  1. *integration increases exposure of neighbors*
  2. *diversification decreases exposure of specific neighbors*
- Interconnection:
  - to continue to cascade widely, the network must have sufficiently large components.

5.4. Eurodebt (Elliott et al., 2014)

- 6 key countries in Europe that have large cross holdings of each other's sovereign debt
- isolated system
- what happens if values fall and debt is devalued

- Pros:
  - tractable
  - interesting implications
  - can be taken to data
- Cons
  - cross-holding based model
  - mechanical propagation through cross-holdings, without economic incentives

Bibliography

Acemoglu, D., Ozdaglar, A., & Tahbaz-Salehi, A. (2015). Systemic Risk and Stability in Financial Networks. *American Economic Review*, *105*(2), 564–608. https://doi.org/10.1257/aer.20130456

Allen, F., & Gale, D. (2000). Financial Contagion. *Journal of Political Economy*, *108*(1), 1–33. http://www.jstor.org/stable/10.1086/262109

Ballester, C., Calvó-Armengol, A., & Zenou, Y. (2006). Who's Who in Networks. Wanted: The Key Player. *Econometrica*, *74*(5), 1403–1417. http://www.jstor.org/stable/3805930

Banerjee, A., Chandrasekhar, A. G., Duflo, E., & Jackson, M. O. (2013). The Diffusion of Microfinance. *Science*, *341*(6144), 1236498. https://doi.org/10.1126/science.1236498

Bramoullé, Y., & Kranton, R. (2007). Public goods in networks. *Journal of Economic Theory*, *135*(1), 478–494. https://doi.org/10.1016/j.jet.2006.06.006

Calvó-Armengol, A., & Jackson, M. O. (2004). The Effects of Social Networks on Employment and Inequality. *American Economic Review*, *94*(3), 426–454. https://doi.org/10.1257/0002828041464542

Calvó-Armengol, A., & Jackson, M. O. (2007). Networks in labor markets: Wage and employment dynamics and inequality. *Journal of Economic Theory*, *132*(1), 27–46. https://doi.org/https://doi.org/10.1016/j.jet.2005.07.007

Coleman, J. S. (1988). Social Capital in the Creation of Human Capital. *American Journal of Sociology*, *94*(1), S95–S120. http://www.jstor.org/stable/2780243

Elliott, M., Golub, B., & Jackson, M. O. (2014). Financial Networks and Contagion. *American Economic Review*, *104*(10), 3115–3153. https://doi.org/10.1257/aer.104.10.3115

Jackson, M. O. (2019). The Friendship Paradox and Systematic Biases in Perceptions and Social Norms. *Journal of Political Economy*, *127*(2), 777–818. https://doi.org/10.1086/701031

Jackson, M. O., Rodriguez-Barraquer, T., & Tan, X. (2012). Social Capital and Social Quilts: Network Patterns of Favor Exchange. *American Economic Review*, *102*(5), 1857–1897. https://doi.org/10.1257/aer.102.5.1857

Perkins, H. W., Haines, M. P., & Rice, R. (2005). Misperceiving the college drinking norm and related problems: a nationwide study of exposure to prevention information, perceived norms and student alcohol misuse. *Journal of Studies on Alcohol*, *66*(4), 470–478.

Tucker, C. J., Finkelhor, D., Turner, H., & Shattuck, A. (2013). Association of sibling aggression with child and adolescent mental health. *Pediatrics*, *132*(1), 79–84.

Valente, T. W., Unger, J. B., & Johnson, C. A. (2005). Do popular students smoke? The association between popularity and smoking among middle school students. *Journal of Adolescent Health*, *37*(4), 323–329.

Index of Figures

Figure 1 Equilibrium 1: all choose 0 6
Figure 2 Equilibrium 2: 3 nodes choose 1 6
Figure 3 Multiple Equilibria 8
Figure 4 Cascade model 10
Figure 5 Cascade model: result 10
Figure 6 Network structure for Example 1.5.2 11
Figure 7 An eq. of Example 1.5.2 11
Figure 8 eq. of star network 12
Figure 9 Eq. of BK 07 21
Figure 10 Own preferences 24
Figure 11 Eq. of this shirt game 24
Figure 12 Perceived averages 25
Figure 13 27
Figure 14 28
Figure 15 31
Figure 16 Community-enforcement 32
Figure 17 33
Figure 18 33
Figure 19 34
Figure 20 34
Figure 21 35
Figure 22 closure 35
Figure 23 Support 36
Figure 24 36
Figure 25 37
Figure 26 3 types of networks 38
Figure 27 38
Figure 28 Eq. and efficiency 39
Figure 29 40
Figure 30 cascade of 2 firms 43
Figure 31 diversification and contagion 45
Figure 32 integration 46