

Image-guided cryoablation for the treatment of painful musculoskeletal metastatic disease: a single-center experience

J. David Prologo · Matthew Passalacqua ·
Indravadan Patel · Nathan Bohnert · David J. Corn

Received: 23 April 2014 / Accepted: 2 June 2014 / Published online: 28 June 2014
© ISS 2014

Abstract

Purpose The role of image-guided thermal ablation techniques for the nonoperative local management of painful osseous metastatic disease has expanded during recent years, and several advantages of cryoablation in this setting have emerged. The purpose of this study is to retrospectively evaluate and report a single-center experience of CT-guided percutaneous cryoablation in the setting of painful musculoskeletal metastatic disease.

Methods This study was approved by the institutional review board and is compliant with the Health Insurance Portability and Accountability Act. Electronic medical records of all patients who underwent percutaneous image-guided palliative cryoablation at our institution were reviewed ($n=61$). An intent-to-treat analysis was performed. Records were reviewed for demographic data and anatomical data, primary tumor type, procedure details, and

outcome—including change in analgesic requirements (expressed as morphine equivalent dosages), pain scores (utilizing the clinically implemented visual analog scale), subsequent therapies (including radiation and/or surgery), and complications during the 24 h following the procedure and at 3 months. Patients were excluded ($n=7$) if data were not retrospectively identifiable at the defined time points.

Results Fifty-four tumors were ablated in 50 patients. There were statistically significant decreases in the median VAS score and narcotic usage at both 24 h and 3 months ($p<0.000$). Six patients (11 %) incurred complications related to their therapy. Two patients had no relief at 24 h, of which both reported worsened pain at 3 months. One patient had initial relief but symptom recurrence at 3 months. Four patients went on to have radiation therapy of the ablation site at some point following the procedure.

Conclusions CT-guided cryoablation is a safe, effective, reproducible procedural option for the nonoperative local treatment of painful musculoskeletal metastatic disease.

J. D. Prologo (✉)
Emory University Hospital, Department of Radiology and Imaging Sciences, Division of Interventional Radiology and Image-Guided Medicine, 1364 Clifton Road, NE Suite D112, Atlanta GA 30322, USA
e-mail: jdprologo@hotmail.com

M. Passalacqua · I. Patel · N. Bohnert
Department of Radiology, University Hospitals Case Medical Center, 11,100 Euclid Avenue, Cleveland, OH 44106, USA

M. Passalacqua
e-mail: Matthew.passalacqua@uhhospitals.org

I. Patel
e-mail: Indravadan.patel2@uhhospitals.org

N. Bohnert
e-mail: Nathan.bohnert@uhhospitals.org

D. J. Corn
Departments of Radiology and Biomedical Engineering, University Hospitals Case Medical Center, Cleveland, OH, USA
e-mail: Djc26@case.edu

Keywords Percutaneous Cryoablation · Musculoskeletal Intervention

Introduction

Pain from neoplastic involvement of bone is a major cause of morbidity in cancer patients [1–6]. The role of interventional radiology in the management of these patients has evolved over recent years [7–10], as have the techniques of thermal ablation [11–22]. Specifically, image-guided percutaneous cryoablation is emerging as a safe, efficacious, and durable therapy for the palliation of painful musculoskeletal metastatic disease [13, 19–21, 23–30].

Advantages of cryoablation when compared to radiofrequency or microwave ablation are direct visualization of the ablation zone, decreased intraprocedural and postprocedural pain, and the ability to utilize multiple probes in variable configurations to create tailored additive overlapping ablation zones [13, 23, 26, 31, 32].

The purpose of this study is to retrospectively evaluate and report a single-center experience of CT-guided percutaneous cryoablation in the setting of painful musculoskeletal metastatic disease.

Materials and methods

Data collection

This study was approved by the institutional review board and is compliant with the Health Insurance Portability and Accountability Act. Electronic medical records of all patients who underwent percutaneous image-guided palliative cryoablation at our institution were reviewed ($n=61$). An intent-to-treat analysis was performed. Records were reviewed for demographic data and anatomical data, primary tumor type, procedure details, and outcome—including change in analgesic requirements (expressed as morphine equivalent dosages), pain scores (utilizing the clinically implemented visual analog scale), subsequent therapies (including radiation and/or surgery), and complications during the 24 h following the procedure and at 3 months. Patients were excluded ($n=7$) if data were not retrospectively identifiable at the defined time points. The “3-month” follow-up translated objectively to 80 ± 17.7 days.

Cryoablation procedures

All procedures were performed with CT guidance. Patients were positioned to maximize safe access to the tumors. Anesthesia consultation was dependent on patient comorbidities, ability to position on the table, and predicted tolerance of the procedure. Overall, the great majority (49/54, 91 %) of

procedures were performed with formal anesthesia sedation management, though only 10/54 (19 %) patients were intubated.

In all cases, cryoprobes from Galil (Arden Hills, MN) were used, ranging in size from 13 to 17 G, with predicted ablation zones ranging from $2 \text{ cm} \times 1 \text{ cm}$ – $3.5 \text{ cm} \times 2 \text{ cm}$ (Fig. 1). Probe selection and number varied according to lesion characteristics (Fig. 2). Specifically, the bone or soft tissue interface with the neoplastic lesion was targeted for inclusion in the predicted lethal zone. Each lesion underwent two 10–5-min freeze-thaw intervals, with images obtained at 8 min into the freeze cycle each time. There were no probe manipulations between cycles. No ancillary techniques were employed to protect adjacent structures or improve lesion accessibility.

Outcome variables

The primary outcome variables analyzed were pain relief, opioid usage, and complication rate. Pain relief was evaluated by compiling the patient’s initial report of pain, pain report at 24 h, and at 3-month follow-up. The initial and 24-h reports were documented by the operator (for inpatients and outpatients). The 3-month follow-up responses were gleaned from medical records of outpatient visits to primary care doctors and oncologists. Patient narcotic usage was documented as part of routine clinical, pre-procedural, post procedure, and follow-up consultations and visits. The drugs were converted to morphine equivalent values using established tables [23, 33, 34]. Complications were identified and classified according to the SIR classification system for complications by outcome [14].

Statistical analysis

Pain relief and opioid usage were evaluated for median differences using the Wilcoxon signed-rank test. In order to analyze the non-normal data distribution (Fig. 3) appropriately, results are reported as median \pm absolute deviation (as opposed to




Fig. 1 Range of isotherm maps for cryoablation probes used during the study period. The temperature ranges closest to the probes (arrows) are the predicted lethal zones. Probes A–E are 17 G; probe F is 13 G




Fig. 2 Ablation images demonstrating the range of approaches. **a** Single axial CT image at the level of T10 in a 47-year-old female with breast cancer and focal back pain demonstrates a 1×2 -cm lytic pedicle lesion (arrows) corresponding to her symptoms. **b** Through a co-axial anchor (small arrow), a 17 G cryoablation probe [“B” from Fig. 1 (small arrow)] was advanced across the lesion and freeze-thaw cycles undertaken. **c** The next morning, the patient had polymethylmethacrylate instilled to the ablation bed under combination CT/fluoroscopic guidance. **d** Single axial PET-MR image in a 61-year-old male with metastatic sarcoma and L4

radiculopathy demonstrating FDG activity superimposed on an underlying 12-cm mass lesion in the right psoas muscle (arrow). **e** Single axial image obtained during the cryoablation demonstrating the iceball created by probe “E” from Fig. 1 reaching to the periphery of the mass in the region of the exiting nerve root (arrows). **f** Additional image from the same patient shows a second, similar probe superiorly—demonstrating the iceball margins (arrows) in relation to the mass and the need for additional probes

means, the former being resistant to outliers). All computations were performed using **Stata** Statistical Software, Release 12 (StataCorp LP, College Station, TX). For all statistical analyses, an alpha of 0.05 was used.

Results

Patients and procedures


The demographic, anatomical, primary tumor type, and inclusion criteria distribution of the included patients are delineated in **Table 1**. One procedure was aborted because of inability to traverse the lesion. The remaining ablations were all technically successful with no immediate complications. Fifty-four

tumors were ablated in 50 patients. Three patients had two sites ablated on different days. Eighteen of the ablated patients had cementoplasty performed as part of the procedure. Two of the cementoplasty patients had a 2-day procedure with polymethylmethacrylate being instilled on the 2nd day.

Response to pain

There were statistically significant decreases in the median VAS score and narcotic usage at both 24 h and 3 months (**Table 2**). The median VAS score of all patients reported prior to their procedures was 8 ± 1 . At 24 h and 3 months post procedure, the median VAS score decreased to 3 ± 1 ($p < 0.000$), where it remained at 3 months post procedure. Median VAS score at 3 months was significantly lower than

Fig. 3 Distribution of study population VAS scores **a** and narcotic usage **b** at the study time points



prior to the procedure ($p < 0.000$). Median narcotic usage, expressed as morphine equivalent (the amount of narcotics

taken in 24 h, expressed as a morphine equivalent), was 340 ± 180 mg prior to the procedure. Morphine equivalent values

Table 1 Demographic data, inclusion criteria, tumor type, and anatomical data for patients undergoing percutaneous CT-guided cryoablation for painful musculoskeletal metastatic disease

Inclusion criteria category ¹	Age range, sex, inpatient vs. outpatient	Sex	Inpatient vs. outpatient	Tumor size range	Osseous anatomical locations	Soft tissue anatomical locations	Primary
I. Patients with limited painful metastatic disease in whom conventional tx has failed or been declined. <i>n</i> =47	33–91	28 F, 15 M	19IP, 24OP	1.7 cm–18 cm	Pelvic bones (16), Skull base (2), Upper extremity (6), Sternum (3), Tibia (2), Foot (2), pedicle (2), chest wall (1)	Masticator Space (2), Pterygopalatine fissure (1), Psoas muscle (1), Iliacus muscle (1), Gluteus musculature (2), obturator muscle (1), chest wall (2), neck soft tissue (3)	Breast (10), renal cell (7), squamous cell (4), lung (9), sarcoma (7), melanoma (8), mucoepidermoid (1), prostate (1)
II. Patients at risk for further morbidity related to progression of their lesion. <i>n</i> =4	67–88	3 F, 1 M	4IP	2.5 cm–7 cm	Femur [2 (pending fracture)], iliac bone [1 (pending fracture)]	Retropharyngeal space [1 (pending airway compromise)]	Breast (2), renal cell (1), squamous cell (1)
III. Patients with limited metastatic disease who are not surgical candidates <i>n</i> =3	72–88	0 F, 3 M	3IP	4.5 cm–12 cm	Upper extremity (2)	Pelvic ST mass (1)	Renal cell (2), colon (1)
IV. Patients with systemically symptomatic metastatic disease <i>n</i> =0							

¹ Clinical criteria for image-guided ablation of metastatic disease involving bone and soft tissue beyond the liver and lung, as per the JVIR Research Reporting Standards for Image-guided Ablation of Bone and Soft Tissue Tumors

Table 2 Responses to pain outcome variables before, 24 h, and 3 months post procedure. Values are expressed as median \pm absolute deviation

	Preprocedure	Post procedure (24 h)	3 months
Median VAS	8 \pm 1	3 \pm 1	3 \pm 2
<i>p</i> Value	X	<i>p</i> <0.000	<i>p</i> <0.000
24 h morphine equivalent (mg)	304 \pm 180	90 \pm 70	120 \pm 70
<i>p</i> Value	X	<i>p</i> <0.000	<i>p</i> <0.000

decreased to 85 \pm 70 mg (*p*<0.000) 24 h after the procedure and at 3 months had increased to 130 \pm 70 mg (*p*<0.000). At 3 months post procedure, morphine equivalent values remained significantly lower than prior to the procedure (*p*<0.000).

Two patients had no relief at 24 h, of which both reported worsened pain at 3 months. One patient had initial relief but symptom recurrence at 3 months (Fig. 4). Four patients went on to have radiation therapy of the ablation site at some point following the procedure. Of these, one was planned as a complementary therapy for a very large lesion, and three sought additional therapy for continued pain.




Fig. 4 Images from a patient with symptomatic recurrence at 3 months. **a** Single axial CT taken during an ablation in a 61-year-old male with painful metastatic lung cancer involving the sternum demonstrates four cryoprobes (“D” from Fig. 1) and the associated ice ball (arrows). The patient had initial relief of his symptoms reflected as a reported VAS decrease from 10 to 2. **b** Single axial PET image obtained 3 months later demonstrates abnormal activity at the periphery of the ablation bed (arrows) corresponding to the patient’s recurrent symptoms, interpreted as either recurrence or initial “underablation”

Complications

Six patients (11 %) incurred complications related to their therapy [21, 28]. Two of these were minor (A) and consisted of symptomatic osteocartilaginous damage beyond the lesion, which resolved. Four were major [C (*n*=1) and D (*n*=3)]: two complete femoral fractures following ablation and cementoplasty of proximal lytic lesions and two transient neuropathies resulting from damage to adjacent nerves during the ablation—one suprascapular nerve involvement during a supraclavicular metastatic lymph node ablation with associated denervation changes on MR and one sciatic nerve involvement during an intragluteal metastatic melanoma deposit ablation (Table 3).

Discussion

Up to 85 % of patients with primary cancers have bone metastases, and half of these will develop intractable pain related to their osseous lesion [1–4]. Pain from neoplastic involvement of bone results from (1) osteoclast-mediated proton sensitization of the sensory fibers at the mineralized bone-tumor interface, (2) sensitization or activation of sensory nerve fibers by products directly produced in tumor and tumor stromal cells, and (3) distortion of mechano-sensitive fibers following normal mechanical stress because of loss of tensile strength in the bone due to cancer [5, 6].

Management options for treatment of symptomatic bone metastases include systemic and local interventions. Bisphosphonates, radionuclides, and chemotherapy are systemically administered in select patients for palliation [7, 35, 36], and systemic analgesic use is well established in cancer pain [37, 38]. The current standard of care for local treatment

Table 3 Complications

Minor complications	
A. No therapy. No consequence	
B. Nominal therapy, no consequence, includes overnight admission for observation only	<i>N</i> =2 Two symptomatic osteocartilaginous injuries outside the targeted lesion
Major complications	
C. Require therapy, minor hospitalization (<48 h)	<i>N</i> =1 Peri-ablational neuropathy
D. Require major therapy, unplanned increase in the level of care, prolonged hospitalization (>48 h)	<i>N</i> =3 Two proximal femoral fractures, one peri-ablational neuropathy
E. Permanent adverse sequelae	
F. Death	

of extra-spinal metastases is external beam radiation therapy [39–43]. Indications for operative local management include fractures, immobilization, and pain refractory to radiation therapy [44]. The role of image-guided thermal ablation techniques for the nonoperative local management of painful osseous metastatic disease has expanded during recent years [7–13, 19, 20, 23–29].

The evolution of these ablative procedures has been toward cryotherapy. The foundation for this evolution was established through key papers demonstrating the safety and efficacy of image-guided radiofrequency ablation for the palliation of painful musculoskeletal metastatic disease [17, 18]. Over time, cryoablation appeared in descriptive papers, review articles, and comprehensive reviews [12, 19, 25, 45–47] and afforded several advantages—including direct visualization of the ablation zone, decreased intra-procedural and post-procedural pain, and the ability to utilize multiple probes in variable configurations to create tailored additive overlapping ablation zones [13, 23, 26, 31, 32].

In 2011, Thacker et al. showed in a retrospective review of 58 patients, divided as 36 cryoablation and 22 radiofrequency ablation, that those who underwent cryoablation for painful musculoskeletal metastatic disease had significantly less short-term analgesic requirement and shorter hospital stays than those who underwent radiofrequency ablation [23].

Subsequently, Callstrom et al. reported the results of a multicenter prospective single-arm trial of 61 patients who underwent image-guided cryoablation for the palliation of painful osseous metastases [26]. Patients had significant decreases in pain and subjective improvements in quality of life at 1, 4, 8, and 24 weeks following the procedure as reflected through Brief Pain Inventory scores, with only one patient incurring a major complication. Similarly, Masala et al. demonstrated symptomatic relief (through VAS scores) following CT-guided cryoablation for painful osseous metastases that correlated positively with post-procedure radiopharmaceutical activity on PET-CT [30].

In the spine, Masala et al. retrospectively described safety and efficacy—through reduced pain (VAS scores) and disability (ODI)—of cryoablation combined with cementoplasty for the palliation of painful vertebral metastases that was at least equivalent to vertebroplasty alone [27]. Although primary, Kurup et al. also reported local control of recurrent sacrococcygeal tumors with cryoablation [28], and recent reports have begun to link cryoablation to systemic inflammatory responses [48–56], as well as suggest potential benefits for patients who undergo cryoablation of metastatic disease [29, 57, 58].

The cytotoxic effects of cryoablation are mediated through the formation of intracellular ice crystals during probe-mediated temperature manipulation. The crystals cause denaturation of proteins and shearing of intracellular structures, including cell membrane rupture. A liquid gas, commonly

argon, is used to rapidly cool the tip of the cryoprobe, forming an enlarging ice-ball over time followed by a “thawing” phase, commonly achieved with helium gas, resulting in an osmotic gradient [59]. This causes water to rush into, swell and then burst the tumor cell, eventually leading to cell hypoxia via indirect ischemic injury [60].

To our knowledge, this is the first single center review describing cryoablation as the exclusive thermal ablation modality for the treatment of painful musculoskeletal metastatic disease to follow the prospective multicenter study put forth by Callstrom et al. The number of patients in this study is similar to that enrolled in the prospective single-arm study. The patient demographics, tumor types, and inclusion criteria are comparable as well. In the present study, patients experienced significant palliation as reflected by decreased VAS scores and narcotic usage—consistent with previously published results. The overall complication rate reported here is slightly higher than in previous reports, but is all inclusive. When only major complications are considered, our rate is similar to previously published results (7 %). Three patients in our series clearly did not benefit from the procedure; two never experienced pain relief and one had recurrence at the periphery of the ablation bed. Upon detailed review, the two patients who had no relief had very large lesions.

This study is limited by its retrospective methodology. It is further limited by the variability of our second follow-up point. That is, there was a fairly large deviation from the mean when considering the population for “3-month” follow-up. Finally, it is limited by the utilization of VAS scores for pain relief vs. the more inclusive Brief Pain Inventory.

Our safety and efficacy results are commensurate with published studies of image-guided cryoablation in the setting of painful musculoskeletal metastatic disease and may have contributory value in establishing the reproducibility of these procedures for palliation. A prospective randomized study of this therapy vs. the historical standard of care radiation therapy will be valuable going forward.

Conflict of interest The primary author of this manuscript has consulted on behalf of Galil in the past (2012).

Disclosure Have provided paid consultation in the past for Galil Medical.

References

1. Coleman RE. Clinical features of metastatic bone disease and risk of skeletal morbidity. *Clin Cancer Res: Off J Am Assoc Cancer Res*. 2006;15(12):6243–9.
2. van den Beuken-van Everdingen MH, de Rijke JM, Kessels AG, Schouten HC, van Kleef M, Patijn J. Prevalence of pain in patients

- with cancer: a systematic review of the past 40 years. *Ann Oncol: Off J Eur Soc Med Oncol/ESMO*. 2007;18(9):1437–49.
3. Jimenez Andrade JM, Mantyh P. Cancer Pain: From the Development of Mouse Models to Human Clinical Trials. In: Kruger L, Light AR, editors. *Translational Pain Research: From Mouse to Man*. Frontiers in Neuroscience. Boca Raton, FL: 2010.
 4. Plunkett TA, Smith P, Rubens RD. Risk of complications from bone metastases in breast cancer. implications for management. *Eur J Cancer*. 2000;36(4):476–82.
 5. Jimenez-Andrade JM, Mantyh WG, Bloom AP, Ferng AS, Geffre CP, Mantyh PW. Bone cancer pain. *Ann N Y Acad Sci*. 2010;1198:173–81.
 6. Goblirsch MJ, Zwolak PP, Clohisy DR. Biology of bone cancer pain. *Clin Cancer Res: Off J Am Assoc Cancer Res*. 2006;12(20 Pt 2): 6231–5.
 7. Soares LG, Chan VW. The rationale for a multimodal approach in the management of breakthrough cancer pain: a review. *Am J Hosp Palliat Care*. 2007;24(5):430–9.
 8. Requarth J. Image-guided palliative care procedures. *Surg Clin N Am*. 2011;91(2):367–402.
 9. McCullough HK, Bain RM, Clark HP, Requarth JA. The radiologist as a palliative care subspecialist: providing symptom relief when cure is not possible. *AJR Am J Roentgenol*. 2011;196(2):462–7.
 10. Bhaskar AK. Interventional management of cancer pain. *Curr Opin Support Palliat Care*. 2012;6(1):1–9.
 11. Nazario J, Tam AL. Ablation of bone metastases. *Surg Oncol Clin N Am*. 2011;20(2):355–68.
 12. Nazario J, Hernandez J, Tam AL. Thermal ablation of painful bone metastases. *Tech Vasc Interv Radiol*. 2011;14(3):150–9.
 13. Rosenthal D, Callstrom MR. Critical review and state of the art in interventional oncology: benign and metastatic disease involving bone. *Radiology*. 2012;262(3):765–80.
 14. Goldberg SN, Charboneau JW, Dodd 3rd GD, Dupuy DE, Gervais DA, Gillams AR, et al. Image-guided tumor ablation: proposal for standardization of terms and reporting criteria. *Radiology*. 2003;228(2):335–45.
 15. Callstrom MR, Charboneau JW, Goetz MP, Rubin J, Atwell TD, Farrell MA, et al. Image-guided ablation of painful metastatic bone tumors: a new and effective approach to a difficult problem. *Skelet Radiol*. 2006;35(1):1–15.
 16. Munk PL, Rashid F, Heran MK, Papirny M, Liu DM, Malfair D, et al. Combined cementoplasty and radiofrequency ablation in the treatment of painful neoplastic lesions of bone. *J Vasc Interv Radiol: JVIR*. 2009;20(7):903–11.
 17. Dupuy DE, Liu D, Hartfeil D, Hanna L, Blume JD, Ahrar K, et al. Percutaneous radiofrequency ablation of painful osseous metastases: a multicenter American College of Radiology Imaging Network trial. *Cancer*. 2010;116(4):989–97.
 18. Goetz MP, Callstrom MR, Charboneau JW, Farrell MA, Maus TP, Welch TJ, et al. Percutaneous image-guided radiofrequency ablation of painful metastases involving bone: a multicenter study. *J Clin Oncol: Off J Am Soc Clin Oncol*. 2004;22(2):300–6.
 19. Gangi A, Buy X. Percutaneous bone tumor management. *Semin Interv Radiol*. 2010;27(2):124–36.
 20. Kurup AN, Callstrom MR. Image-guided percutaneous ablation of bone and soft tissue tumors. *Semin Interv Radiol*. 2010;27(3):276–84.
 21. Prologo JD, Patel I, Bueth J, Bohnert N. Ablation zones and weight-bearing bones: points of caution for the palliative interventionalist. *J Vasc Interv Radiol: JVIR*. 2014;25(5):769–75.
 22. Patel IJ, Pirasteh A, Passalacqua MA, Robbin MR, Hsu DP, Bueth J, et al. Palliative procedures for the interventional oncologist. *AJR Am J Roentgenol*. 2013;201(4):726–35.
 23. Thacker PG, Callstrom MR, Curry TB, Mandrekar JN, Atwell TD, Goetz MP, et al. Palliation of painful metastatic disease involving bone with imaging-guided treatment: comparison of patients' immediate response to radiofrequency ablation and cryoablation. *AJR Am J Roentgenol*. 2011;197(2):510–5.
 24. Popken F, Michael JW, Zarghooni K, Sobottke R, Kasper HU, Blaecker D, et al. Stability changes after cryosurgery in long tubular bones in correlation to histological results: an animal trial. *Arch Orthop Trauma Surg*. 2009;129(6):857–62.
 25. Castaneda Rodriguez WR, Callstrom MR. Effective pain palliation and prevention of fracture for axial-loading skeletal metastases using combined cryoablation and cementoplasty. *Tech Vasc Interv Radiol*. 2011;14(3):160–9.
 26. Callstrom MR, Dupuy DE, Solomon SB, Beres RA, Littrup PJ, Davis KW, et al. Percutaneous image-guided cryoablation of painful metastases involving bone: Multicenter trial. *Cancer*. 2012 Oct 12
 27. Masala S, Chiochi M, Taglieri A, Bindi A, Nezzo M, De Vivo D, et al. Combined use of percutaneous cryoablation and vertebroplasty with 3D rotational angiograph in treatment of single vertebral metastasis: comparison with vertebroplasty. *Neuroradiology*. 2012 Sep 27
 28. Kurup AN, Woodrum DA, Morris JM, Atwell TD, Schmit GD, Welch TJ, et al. Cryoablation of recurrent sacrococcygeal tumors. *J Vasc Interv Radiol: JVIR*. 2012;23(8):1070–5.
 29. McMenomy BP, Kurup AN, Johnson GB, Carter RE, McWilliams RR, Markovic SN, et al. Percutaneous cryoablation of musculoskeletal oligometastatic disease for complete remission. *Journal of vascular and interventional radiology:JVIR*. 2012 Dec 19
 30. Masala S, Schillaci O, Bartolucci AD, Calabria F, Mammucari M, Simonetti G. Metabolic and clinical assessment of efficacy of cryoablation therapy on skeletal masses by 18 F-FDG positron emission tomography/computed tomography (PET/CT) and visual analogue scale (VAS): initial experience. *Skelet Radiol*. 2011;40(2):159–65.
 31. Atwell TD, Farrell MA, Callstrom MR, Charboneau JW, Leibovich BC, Patterson DE, et al. Percutaneous cryoablation of 40 solid renal tumors with US guidance and CT monitoring: initial experience. *Radiology*. 2007;243(1):276–83.
 32. Truesdale CM, Soulen MC, Clark TW, Mondschein JJ, Wehrenberg-Klee E, Malkowicz SB, et al. Percutaneous computed tomography-guided renal mass radiofrequency ablation versus cryoablation: doses of sedation medication used. *J Vasc Interv Radiol: JVIR*. 2013;24(3): 347–50.
 33. Hallenbeck JL. Pain Management. In: Hallenbeck JL, editor. *Palliative Care Perspectives*. New York: Oxford University Press; 2003. p. 39–43.
 34. Gordon DB, Stevenson KK, Griffie J, Muchka S, Rapp C, Ford-Roberts K. Opioid equianalgesic calculations. *J Palliat Med*. 1999;2(2):209–18.
 35. Gainford MC, Dranitsaris G, Clemons M. Recent developments in bisphosphonates for patients with metastatic breast cancer. *BMJ*. 2005;330(7494):769–73.
 36. Brady D, Parker CC, O'Sullivan JM. Bone-targeting radiopharmaceuticals including radium-223. *Cancer J*. 2013;19(1):71–8.
 37. Mani A SC. Bone Metastases. In: Walsh, TD, editor. *Walsh: Palliative Medicine*. 1st ed. Philadelphia: Saunders; 2009.
 38. Zech DF, Grond S, Lynch J, Hertel D, Lehmann KA. Validation of World Health Organization Guidelines for cancer pain relief: a 10-year prospective study. *Pain*. 1995;63(1):65–76.
 39. Wu JS, Wong R, Johnston M, Bezjak A, Whelan T. Cancer Care Ontario Practice Guidelines Initiative Supportive Care G. Meta-analysis of dose-fractionation radiotherapy trials for the palliation of painful bone metastases. *Int J Radiat Oncol Biol Phys*. 2003;55(3): 594–605.
 40. Gaze MN, Kelly CG, Kerr GR, Cull A, Cowie VJ, Gregor A, et al. Pain relief and quality of life following radiotherapy for bone metastases: a randomised trial of two fractionation schedules. *Radiother Oncol: J Eur Soc Ther Radiol Oncol*. 1997;45(2):109–16.

41. Agarawal JP, Swangsilpa T, van der Linden Y, Rades D, Jeremic B, Hoskin PJ. The role of external beam radiotherapy in the management of bone metastases. *Clin Oncol*. 2006;18(10):747–60.
42. Ingunn HC, Robert E. Bone Metastases. *Abeloff's clinical oncology*. 4th ed. Philadelphia: Churchill Livingstone/Elsevier; 2008. p. 845–71.
43. Fairchild A CE. Palliative Radiation Therapy. In: Walsh, TD, editor. *Walsh: Palliative Medicine*. 1st ed. Philadelphia: Saunders; 2009.
44. Wegener B, Schlemmer M, Stemmler J, Jansson V, Durr HR, Pietschmann MF. Analysis of orthopedic surgery of bone metastases in breast cancer patients. *BMC Musculoskelet Disord*. 2012;13(1):232.
45. Choi J, Raghavan M. Diagnostic imaging and image-guided therapy of skeletal metastases. *Cancer Control: J Moffitt Cancer Cent*. 2012;19(2):102–12.
46. Callstrom MR, Charboneau JW. Image-guided palliation of painful metastases using percutaneous ablation. *Tech Vasc Interv Radiol*. 2007;10(2):120–31.
47. Kurup AN, Callstrom MR. Ablation of skeletal metastases: current status. *J Vasc Interv Radiol: JVIR*. 2010;21(8 Suppl):S242–50.
48. Ng KK, Lam CM, Poon RT, Shek TW, To JY, Wo YH, et al. Comparison of systemic responses of radiofrequency ablation, cryotherapy, and surgical resection in a porcine liver model. *Ann Surg Oncol*. 2004;11(7):650–7.
49. Blackwell TS, Debelak JP, Venkatakrishnan A, Schot DJ, Harley DH, Pinson CW, et al. Acute lung injury after hepatic cryoablation: correlation with NF-kappa B activation and cytokine production. *Surgery*. 1999;126(3):518–26.
50. Chapman WC, Debelak JP, Wright Pinson C, Washington MK, Atkinson JB, Venkatakrishnan A, et al. Hepatic cryoablation, but not radiofrequency ablation, results in lung inflammation. *Ann Surg*. 2000;231(5):752–61.
51. Gravante G, Sconocchia G, Ong SL, Dennison AR, Lloyd DM. Immunoregulatory effects of liver ablation therapies for the treatment of primary and metastatic liver malignancies. *Liver Int: Off J Int Assoc Study Liver*. 2009;29(1):18–24.
52. Yang Y, Wang C, Lu Y, Bai W, An L, Qu J, et al. Outcomes of ultrasound-guided percutaneous argon-helium cryoablation of hepatocellular carcinoma. *J Hepato-Biliary-Pancreatic Sci*. 2012;19(6):674–84.
53. Jansen MC, van Hillegersberg R, Schoots IG, Levi M, Beek JF, Crezee H, et al. Cryoablation induces greater inflammatory and coagulative responses than radiofrequency ablation or laser induced thermotherapy in a rat liver model. *Surgery*. 2010;147(5):686–95.
54. Soanes WA, Ablin RJ, Gonder MJ. Remission of metastatic lesions following cryosurgery in prostatic cancer: immunologic considerations. *J Urol*. 1970;104(1):154–9.
55. Ravindranath MH, Wood TF, Soh D, Gonzales A, Muthugounder S, Perez C, et al. Cryosurgical ablation of liver tumors in colon cancer patients increases the serum total ganglioside level and then selectively augments antiganglioside IgM. *Cryobiology*. 2002;45(1):10–21.
56. Erinjeri JP, Thomas CT, Samoilia A, Fleisher M, Gonen M, Sofocleous CT, et al. Image-guided thermal ablation of tumors increases the plasma level of interleukin-6 and interleukin-10. *Journal of vascular and interventional radiology: JVIR*. 2013 Apr 9.
57. Bang HJ, Littrup PJ, Currier BP, Goodrich DJ, Aoun HD, Klein LC, et al. Percutaneous cryoablation of metastatic lesions from non-small-cell lung carcinoma: initial survival, local control, and cost observations. *J Vasc Interv Radiol: JVIR*. 2012;23(6):761–9.
58. Bang HJ, Littrup PJ, Goodrich DJ, Currier BP, Aoun HD, Heilbrun LK, et al. Percutaneous cryoablation of metastatic renal cell carcinoma for local tumor control: feasibility, outcomes, and estimated cost-effectiveness for palliation. *J Vasc Interv Radiol: JVIR*. 2012;23(6):770–7.
59. Gage AA, Baust JG. Cryosurgery for tumors. *J Am Coll Surg*. 2007;205(2):342–56.
60. Theodorescu D. Cancer cryotherapy: evolution and biology. *Rev Urol*. 2004;6 Suppl 4:S9–S19.

Reproduced with permission of the copyright owner. Further reproduction prohibited without permission.