

who underwent staged procedures were included but opiate prescribing data were collected based on 60-stage ureteroscopy to allow follow-up of medication use. All patients undergoing ureteroscopy at our institution have a postoperative visit at 60 weeks with a renal ultrasound. Only primary residents of Tennessee at the time of surgery were included to enable for linkage to the **Tennessee Controlled Substance Monitoring Database (TN CSMD)** for opiate prescribing data. In accordance with the Controlled Substance Monitoring Act of 2002 and the Tennessee Prescription Safety Act of 2016, all health care providers who dispense controlled substances (Schedule II, III, IV, & V) in Tennessee for more than 15 days per year are required to register to monitor the dispensing of controlled substances. Local Institutional Review Board approval was obtained before data collection.

Demographic data collected included age, sex, race, BMI (body mass index; calculated by weight/height², kg/m²), **American Society of Anesthesiologists (ASA) score, past stone surgery**, and documented diagnosis of psychiatric disorder. Surgical and stone variables collected included operative time, presence of stent preoperatively, laterality, postoperative stent duration, ureteral access sheath use, and stone diameter.

Prescribing data was categorized by preoperative and postoperative prescriptions, and then further classified by opiate and nonopiate medications. Prescription data from the CSMD provide the quantity and dose of opiate medication received by a patient from a pharmacy along with prescriber information. Preoperative opiate data included chronic opiate use (defined as more than two opiate prescriptions dispensed 60 days before surgery) and the following variables 30 days before surgery: number of opiate prescriptions dispensed, number of providers prescribing opiates, and morphine equivalent dose (MED) of the prescriptions dispensed. As a reference, 100 MED is equivalent to 100 mg of hydrocodone or 67 mg of oxycodone. Nonopiate preoperative prescription data within 30 days before surgery were assessed. Postoperative prescription data collected included the number of days of postoperative opiates prescribed at the time of surgery, total MED provided to the patient, and nonopiate medications prescribed to patients.

The two outcomes evaluated in this study were (1) obtaining an additional opiate prescription within 30 days after ureteroscopic stone surgery, and (2) continued opiate use 60 days after surgery as ascertained by the TN CSMD. We chose the 30- and 60-day time points to capture pain complaints or complications before or after routine 6-week postoperative follow-up visit.

Univariate analyses were performed to determine associations with obtaining an additional opiate prescription within 30 days or 60 days after surgery. A multivariable logistic regression model was fit to include all potential predictors, such as demographic, surgical, and opiate prescription variables. The opiate prescription variables were separately modeled as they were highly correlated. All analyses were carried out using **Stata** version 14 (StataCorp, LLC). All statistical tests were based on two-sided probability and $p < 0.05$ was considered statistically significant.

Results

A total of 310 patients underwent ureteroscopy during the study period. Excluded patients included 33 undergoing ur-

eteroscopy for diagnosis other than ureteral or renal stone disease, 36 undergoing staged procedures, and 33 who were not primary residents of Tennessee. A total of 208 patients were included in the final analysis.

The patient population was predominantly male ($n = 116$, 56%) and white ($n = 191$, 92%). Most individuals were middle-aged (mean \pm SD age 52.4 \pm 15.1 years), with ASA scores of two ($n = 85$, 41%) or three ($n = 110$, 53%), and without concurrent psychiatric diagnoses ($n = 172$, 83%). There were 80 patients (38.5%) who had prior stone-related surgical procedures.

Preoperatively, a minority of patients were chronic opiate users ($n = 21$, 10%, Table 1). Within 30 days before surgery, 61% ($n = 127$) had received at least one opiate prescription, whereas 6% ($n = 12$) had received at least three prescriptions. Nonopiate prescriptions were prescribed preoperatively * 20% of the time.

Most ureteroscopic procedures performed were unilateral ($n = 173$, 83%), and the majority of patients did not have preoperative indwelling ureteral stents ($n = 139$, 67%). Ureteral access sheaths were used commonly ($n = 129$, 55%), and the majority of patients had stents placed intraoperatively ($n = 200$, 96%). Patients most commonly had stents removed 40 days postoperatively ($n = 124$, 60%).

Postoperatively, the majority of patients received opiate prescriptions ($n = 196$, 93.3%) with an average MED of 105.2 (mean \pm SD, 105.2 \pm 81.2). Patients most commonly received 40 days of opiate medications ($N = 104$, 50%). In addition, patients received nonopiate medications * 75% of the time (Table 1).

Overall, 12% ($n = 25$) of patients dispensed an additional opiate prescription within 30 days after ureteroscopy (Table 2). On univariate analysis, associated factors included younger age, greater number of preoperative prescriptions, greater number of providers prescribing opiates preoperatively, and higher MED use preoperatively (Table 3). Having a concurrent psychiatric diagnosis and chronic opiate use history trended toward but did not achieve significance ($p = 0.14$ and 0.09 , respectively). Age and opiate-related associations remained significant upon multivariate analysis when modeling opiate variables separately (Supplementary Tables S1-S3). In this group, 10 patients (40%) had no reported complications, 12 patients (48%) made phone calls to the clinic for pain-related concerns, four patients (16%) had unanticipated clinic visits, two patients (8%) were seen in the emergency department, and there was one hospital readmission (4%, data not shown).

Approximately 1 in 15 patients ($n = 14$, 6.7%) received an additional opiate prescription after 60 days postoperatively (Table 2). The majority of opiate prescriptions ($n = 13$, data not shown) provided to these patients were provided by providers not affiliated with the urology department. Within this cohort, there was one unanticipated clinic visit, one emergency room visit, and two documented phone calls related to pain (data not shown). On univariate analysis, associated factors included a greater number of preoperative prescriptions, greater number of providers prescribing opiates preoperatively, and higher preoperative MED use (Table 3). On multivariate analysis, greater opiate exposure preoperatively, lower BMI, and greater ASA were significant predictors (Table 4 and Supplementary Tables S1-S3). No patient who had an additional opiate prescription beyond 60

Table 1. Patient Characteristics

Demographics	
Age, years (mean \pm SD)	52.4 \pm 15.1
BMI (mean \pm SD)	30.5 \pm 8.1
Sex (%)	
Male	116 (55.8)
Female	92 (44.2)
Race (%)	
Caucasian	191 (91.8)
Other	18 (8.2)
ASA score (%)	
1	9 (4.3)
2	85 (40.9)
3	110 (52.9)
4	3 (1.9)
Prior psychiatric diagnosis (%)	
Yes	36 (17.3)
No	172 (82.7)
Prior stone procedure (%)	
Yes	80 (38.5)
No	128 (61.5)
Medication history	
Chronic opiate use (%)	
Yes	21 (10.1)
No	187 (89.9)
Preoperative opiate prescriptions <30 day, number (%)	
0	81 (38.9)
1	64 (30.8)
2	42 (20.2)
3+	21 (10.1)
Preoperative providers prescribing opiates <30 day, number (%)	
0	81 (38.9)
1	74 (35.6)
2	41 (19.7)
3+	12 (5.8)
Preoperative MED prescribed, tertiles (mean \pm SD)	
Group 1 (N = 49)	107.8 \pm 39.9
Group 2 (N = 37)	240.6 \pm 45.5
Group 3 (N = 41)	3203.7 \pm 7157.4
Preoperative oxybutynin prescription (%)	
Yes	47 (22.6)
No	161 (77.4)
Preoperative tamsulosin prescription (%)	
Yes	56 (26.9)
No	152 (73.1)
Preoperative phenazopyridine (%)	
Yes	45 (21.6)
No	163 (78.4)
Postoperative opiate prescription, days (%)	
0	14 (6.7)
1-6	79 (38.0)
7-10	104 (50.0)
11+	13 (6.3)
Postoperative MED prescribed (mean \pm SD)	105.2 \pm 91.2
Postoperative oxybutynin prescription (%)	
Yes	156 (75.0)
No	52 (25.0)

(continued)

Table 1. (Continued)

Demographics	
Postoperative tamsulosin prescription (%)	
Yes	153 (73.6)
No	55 (26.4)
Postoperative phenazopyridine prescription (%)	
Yes	156 (75.0)
No	52 (25.0)
Surgical factors	
Laterality (%)	
Right	74 (35.6)
Left	99 (47.6)
Bilateral	35 (16.8)
Preoperative ureteral stent (%)	
Yes	69 (33.2)
No	139 (66.8)
Stone size, diameter (mean \pm SD)	8.48 \pm 4.82
Operative time, minutes (mean \pm SD)	55.7 \pm 38.4
Use of ureteral access sheath intraoperatively (%)	
Yes	124 (59.6)
No	84 (40.4)
Postoperative stent duration, days (%) ^a	
0	8 (3.9)
1-6	34 (16.4)
7-10	124 (59.6)
11+	40 (19.2)

^aStent duration information missing on electronic medical record for two patients.

ASA = American Society of Anesthesiologists; MED = morphine equivalent dose.

days postoperatively was a chronic opiate user in the pre-operative period.

Discussion

Managing pain after surgery is important for decreasing pain-related complications but must be balanced with minimizing opiate misuse, dependence, and overdose.^{4,6,8} Studies have shown there is a large MED range of opiate medications provided to patients after common surgeries,⁹ and that a majority of patients report unused leftover opioid medication after procedures.^{7,8,10} Specifically, unanticipated visits after ureteroscopy for stone disease has been reported to be 6.6%, and the majority of these visits are made due to pain.^{5,11} An important finding of our study is that most patients before stone surgery have had opiate exposure within the previous 30 days, and a significant proportion have had multiple prescriptions. This highlights a fundamental challenge of addressing how and when opiates are prescribed to kidney stone patients as they often encounter multiple providers across different settings.

Table 2. Additional Opiate Prescriptions After Ureteroscopy

Within 30 days postoperatively (%)	
Yes	25 (12.0)
No	183 (88.0)
After 60 days postoperatively (%)	
Yes	14 (6.7)
No	194 (93.3)

Table 3. Factors Associated with Obtaining an Additional Opiate Prescription Within 30 Days or More Than 60 Days After Ureteroscopy

Variable	Within 30 days		After 60 days	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Age	0.97 (0.93-1.01)	0.009	1.01 (0.98-1.06)	0.32
BMI	1.01 (0.96-1.07)	0.61	0.95 (0.87-1.03)	0.23
Female	0.55 (0.23-1.35)	0.19	0.68 (0.22-2.11)	0.51
Caucasian	0.97 (0.21-4.54)	0.97	1.99 (0.41-9.72)	0.40
ASA score	0.88 (0.38-2.03)	0.76	0.54 (1.18-0.82)	0.03
ASA 2	0.47 (0.085-2.57)	0.38	UTC	UTC
ASA 3	0.43 (0.080-2.30)	0.32	UTC	UTC
ASA 4	1.17 (0.074-18.35)	0.91	UTC	UTC
Prior psychiatric diagnosis	2.07 (0.79-5.39)	0.14	1.33 (0.35-5.03)	0.67
Prior stone procedure	1.08 (0.46-2.53)	0.87	0.62 (0.19-2.05)	0.44
Chronic opiate use	2.61 (0.86-7.89)	0.09	UTC	UTC
Preoperative opiate prescriptions, number	1.58 (1.6-1.7)	0.026	1.57 (1.14-2.16)	0.005
Preoperative opiate prescription, days	1.6 (1.11-2.32)	0.012	1.02 (1.00-1.03)	0.032
Preoperative providers prescribing opiates	1.68 (1.08-2.62)	0.022	1.55 (1.00-2.39)	0.048
Preoperative MED prescribed				
Group 1 (lower tertile)	0.39 (0.08-1.91)	0.24	2.58 (0.42-16.0)	0.31
Group 2 (middle tertile)	1.11 (0.31-3.93)	0.88	4.79 (0.84-27.42)	0.08
Group 3 (upper tertile)	3.35 (1.22-9.14)	0.02	5.49 (1.02-29.63)	0.048
Preoperative oxybutynin prescription	1.09 (0.41-2.92)	0.86	UTC	UTC
Preoperative tamsulosin prescription	0.48 (0.16-1.47)	0.20	0.19 (0.02-1.52)	0.12
Preoperative phenazopyridine	0.89 (0.32-2.53)	0.83	0.59 (0.13-2.72)	0.49
Postoperative opiate prescriptions, days	1.0 (0.86-1.16)	0.98	0.9 (0.70-1.15)	0.38
Postoperative MED prescribed	1.0 (1.00-1.00)	0.99	1.00 (0.99-1.01)	0.79
Postoperative oxybutynin	1.38 (0.49-3.89)	0.54	1.24 (0.33-4.62)	0.75
Postoperative tamsulosin	1.5 (0.54-4.22)	0.44	2.26 (0.49-10.4)	0.30
Postoperative phenazopyridine	1.38 (0.49-3.89)	0.54	4.64 (0.59-36.3)	0.14
Stone size, diameter	0.96 (0.87-1.06)	0.40	0.99 (0.88-1.11)	0.83
Operative time, minutes	1.0 (0.99-1.02)	0.81	1.00 (0.98-1.02)	0.87
Ureteral access sheath use	0.84 (0.36-1.96)	0.70	1.24 (0.40-3.83)	0.71
Laterality				
Left	1.47 (0.39-5.55)	0.57	1.45 (0.29-7.18)	0.65
Right	1.67 (0.43-6.48)	0.46	0.94 (0.16-5.41)	0.95
Preoperative ureteral stent	0.76 (0.30-1.91)	0.56	0.53 (0.14-1.96)	0.34
Postoperative ureteral stent duration, days				
1-3	0.4 (0.06-2.70)	0.35	UTC	UTC
4-6	0.38 (0.07-2.08)	0.27	UTC	UTC
7+	0.43 (0.07-2.74)	0.37	UTC	UTC

Due to small sample size of patients who required opiate prescriptions 60 days after ureteroscopy, several data points had frequency of zero and OR was unable to be calculated.

OR = odds ratio; UTC = unable to calculate.

Table 4. Multivariable Logistic Regression Models Days of Preoperative Opiate Prescriptions

Variable	Within 30 days		After 60 days	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Age	0.82 (0.69-0.98)	0.027	0.97 (0.79-1.20)	0.81
Sex	0.61 (0.22-1.70)	0.35	0.52 (0.15-1.84)	0.32
Race	1.21 (0.22-6.56)	0.82	3.8 (0.61-24.18)	0.15
BMI	0.98 (0.93-1.04)	0.58	0.89 (0.80-0.98)	0.022
ASA score	1.31 (0.74-2.31)	0.36	3.08 (1.25-7.61)	0.014
Laterality (left)	1.38 (0.32-5.90)	0.66	2.62 (0.46-14.97)	0.279
Laterality (right)	1.31 (0.29-5.81)	0.72	1.14 (0.17-7.64)	0.089
Preoperative opiate prescription, days	1.73 (1.14-2.61)	0.009	1.97 (1.16-3.36)	0.013

Multivariable logistic regression models were performed for number of preoperative opiate prescriptions, days of preoperative opiate prescriptions, and number of providers prescribing opiate prescriptions preoperatively. These variables were modeled separately, and additional models are provided in Supplementary Tables S16-18.

In our study, 12% of patients required an additional opioid prescription within a 30-day period after ureteroscopy. Risk factors identified included younger age and higher preoperative opiate use, which have been shown in prior studies to be associated with higher postoperative opiate use.^{12,64} A recent nationwide survey reported one in seven youths have prior inappropriate use of opiates, which may lead to increased tolerance to opiates and perception of pain in this patient cohort.^{15,67} Some patients in our study obtained more than three opiate prescriptions preoperatively and obtained opiate prescriptions from more than a single provider (Table 2). These findings mirror those in a study by Kappa and colleagues, who found that both younger age and higher preoperative opiate use were predictors of obtaining opiate pain medications from multiple providers after ureteroscopy.¹⁸ Interestingly, although preoperative opiate use was a predictor of increased postoperative opiate requirements, chronic opiate use, which we defined as having more than two opiate prescriptions >60 days preoperatively, only trended toward significance likely due to small sample size.

We found that ~7% of patients continued to have opiate prescriptions over 60 days after ureteroscopy, suggesting that a small but significant proportion of patients are at high risk for opioid dependence postoperatively. Only 1 of these 14 patients had this prescription obtained from our department. A study from the Centers for Disease Control found that increases in chronic opioid use were observed if patients received more than one opioid prescription, had longer duration of opioid use, or had higher quantities (MED) of opioid prescribed.¹⁹ Another study demonstrated ~6% of patients with new persistent opiate use after surgical procedures.⁶ In our study, risk factors for persistent opiate use included higher preoperative use, lower BMI, and greater ASA score.⁶ Notably, none of the patients with persistent opiate use in our study had a history of chronic opiate use. The rate of persistent opiate use in our study was notably higher than previously reported by Shah and colleagues, which estimated opioid dependence to be present in 0.09% of patients after urologic procedures based on diagnostic codes.⁴ For stone procedures, including percutaneous and endoscopic surgery, 0.10% and 0.15% of patients had hospital visits for opioid dependence or overdose within 90 and 365 days, respectively.⁴ The differences between our studies likely lies in the methods used to identify patients with persistent opiate use after ureteroscopy. Patients with higher ASA scores have a higher risk of developing postoperative complications,²⁰ and new or preexisting comorbidities may be unmasked or exacerbated by surgery, which may result in higher opioid consumption postoperatively. It is possible that patients with higher ASA scores continued to require opiate medications for management of other comorbidities and not post-ureteroscopy pain, but these data were not reviewable in the TN CSMD.

While this study was not designed to determine the optimal postoperative pain regimen for patients undergoing ureteroscopy, it is apparent that the analgesic needs for each patient are different. Preoperative counseling regarding expectations after surgery, and proper use of opiate medications postoperatively have been demonstrated to decrease postoperative opiate consumption while maintaining adequate pain control.^{21,22} Opiate-free protocols after ureteroscopy have been shown to be feasible.²³ Intraoperative use of nonopiate

medications, such as intravenous toradol and belladonna opium suppositories have been shown to help with postoperative pain control after ureteroscopy.^{11,24} Notably in this study, no surgical factors were identified as risk factors for persistent opiate use, including stent duration or sheath use. After surgery, providing a standardized MED per postoperative opiate prescription may help minimize the quantity of opiates prescribed and/or unused by patients.¹⁰ It is important to note that patients who have had prior exposure to opiate medications seem to be at greater risk for continued opiate use, and these at-risk patients may benefit most from targeted counseling regarding the risks and alternatives to opiate medications. Furthermore, providing prescriptions for non-opiate medications including nonsteroidal anti-inflammatory drugs, alpha-blockers, oxybutynin, and phenazopyridine may result in decreased opiate requirement.^{25,67}

Our findings should be interpreted within the context of the limitations of the retrospective design. The relatively low number of patients identified with persistent opiate use after ureteroscopy reduced the power to detect associations. The precise MED amount consumed by each patient is unknown. While we restricted this study to residents of the state, it is possible that prescriptions were given to patients outside of the state that would not be captured in the CSMD. Prescription data from TN CSMD does not list the indication for opioid prescription or specialty of the provider prescribing opiates, and additional prescriptions may have been prescribed for conditions related to stone disease or other diagnoses not related to kidney stone disease.

Conclusion

The majority of kidney stone patients receive opiate medications before surgery and often from multiple providers. Higher preoperative use of opiates was associated with receiving an additional opiate prescription within 30 days and persistent opiate use beyond 60 days after surgery. Younger age, lower BMI, and higher ASA score were also associated with persistent opiate after ureteroscopy.

Acknowledgments

The publication was supported by CTSA award No. UL1 TR002243 from the National Center for Advancing Translational Sciences. Its contents are solely the responsibility of the authors and do not necessarily represent official views of the National Center for Advancing Translational Sciences or the National Institutes of Health.

Author Disclosure Statement

The authors have no disclosures to declare.

Supplementary Material

Supplementary Table S1
Supplementary Table S2
Supplementary Table S3

References

1. Scales CD, Smith AC, Hanley JM, Saigal CS, Urologic Diseases in America Project. Prevalence of kidney stones in the United States. *Eur Urol* 2012;62:160-165.

2. Daniels B, Schoenfeld E, Taylor A, Weisenthal K, Singh D, Moore CL. Predictors of hospital admission and urological intervention in adult emergency department patients with computerized tomography confirmed ureteral stones. *J Urol* 2017;198:1359-1366.
3. Kiechle JE, Gonzalez CM. The opioid crisis and urology. *Urology* 2018;112:27-33.
4. Shah AS, Blackwell RH, Kuo PC, Gupta GN. Rates and risk factors for opioid dependence and overdose after urological surgery. *J Urol* 2017;198:1130-1136.
5. Mittakanti HR, Conti SL, Pao AC, Chertow GM, Liao JC, Leppert JT, Elliott CS. Unplanned emergency department visits and hospital admissions following ureteroscopy: Do ureteral stents make a difference? *Urology* 2018;117:44-50.
6. Brummett CM, Waljee JF, Goesling J, et al. New persistent opioid use after minor and major surgical procedures in us adults. *JAMA Surg* 2017;152:e170504.
7. Bicket MC, Long JJ, Pronovost PJ, Alexander GC, Wu CL. Prescription opioid analgesics commonly unused after surgery: A systematic review. *JAMA Surg* 2017;152:1066-1071.
8. Bates C, Laciak R, Southwick A, Bishoff J. Overprescription of postoperative narcotics: A look at postoperative pain medication delivery, consumption and disposal in urological practice. *J Urol* 2011;185:551-555.
9. Fujii MH, Hodges AC, Russell RL, et al. Post-Discharge Opioid Prescribing and Use after Common Surgical Procedure. *J Am Coll Surg* 2018;226:1004-1012.
10. Tan WH, Yu J, Feaman S, et al. Opioid Medication Use in the Surgical Patient: An Assessment of Prescribing Patterns and Use. *J Am Coll Surg* 2018;227:203-211.
11. Lee FC, Holt SK, Hsi RS, Haynes BM, Harper JD. Preoperative Belladonna and Opium Suppository for Ureteral Stent Pain: A Randomized, Double-blinded, Placebo-controlled Study. *Urology* 2017;100:27-32.
12. Hereford TE, Cryar KA, Edwards PK, Siegel ER, Barnes CL, Mears SC. Patients With Hip or Knee Arthritis Underreport Narcotic Usage. *J Arthroplast* 2018;33:3113-3117.
13. Pugely AJ, Bedard NA, Kalakoti P, et al. Opioid use following cervical spine surgery: Trends and factors associated with long-term use. *Spine J* 2018;18:S1529-S15430.
14. Willis-Gray MG, Husk KE, Brueseke TJ, et al. Predictors of Opioid Administration in the Acute Postoperative Period. *Female Pelvic Med Reconstr Surg* 2018;24:454.
15. Lavandhomme P, Steyaert A. Opioid-free anesthesia opioid side effects: Tolerance and hyperalgesia. *Best Pract Res Clin Anaesthesiol* 2017;31:487-508.
16. Joly V, Richebe P, Guignard B, Fletcher D, Maurette P, Sessler DI, Chauvin M. Remifentanyl-induced postoperative hyperalgesia and its prevention with small-dose ketamine. *Anesthesiology* 2005;103:147-155.
17. Reddy RR, Kent CK, Leahy MA, et al. Youth Risk Behavior Surveillance - United States 2017. *MMWR Surveill Summ* 2018;67:1-14.
18. Kappa SF, Green EA, Miller NL, Herrell SD, Mitchell CR, Mir HR, Resnick MJ. Narcotic Use and Postoperative Doctor Shopping by Patients with Nephrolithiasis Requiring Operative Intervention: Implications for Patient Safety. *J Urol* 2016;196:763-768.
19. Shah A, Hayes CJ, Martin BC. Characteristics of Initial Prescription Episodes and Likelihood of Long-Term Opioid Use - United States, 2006-2015. *MMWR Morb Mortal Wkly Rep* 2017;66:265-269.
20. Wolters U, Wolf T, Stöcker H, Schröder T. ASA classification and perioperative variables as predictors of postoperative outcome. *Br J Anaesth* 1996;77:217-222.
21. Schröder A, Campbell FA, Farhat WA, Salle JLP, Bagli DJ, Lorenzo AJ, Koyle MA. Postoperative pain and analgesia administration in children after urological outpatient procedures. *J Pediatr Urol* 2018;14:171.e1-171.e6.
22. Dwyer CL, Soong M, Hunter A, Dashe J, Tolo E, Kasparian NG. Prospective Evaluation of an Opioid Reduction Protocol in Hand Surgery. *J Hand Surg Am* 2018;43:516-522.e1.
23. Large T, Heiman J, Ross A, Anderson B, Krambeck A. Initial Experience with Narcotic Free Ureteroscopy: A Feasibility Analysis. *J Endourol* 2018;32:907-911.
24. Trowbridge ER, Kim D, Caldwell L, Franko D, Jackson JN, Shilling A, Hull KL. Use of ketorolac after outpatient urogynecologic surgery: A randomized control trial. *Female Pelvic Med Reconstr Surg* 2018;24:281-286.
25. Kwon JK, Cho KS, Oh CK, Kang DH, Lee H, Ham WS, Choi YD, Lee JY. The beneficial effect of alpha-blockers for ureteral stent-related discomfort: Systematic review and network meta-analysis for alfuzosin versus tamsulosin versus placebo. *BMC Urol* 2015;15:55.
26. Maldonado-Avila M, Garduno-Arteaga L, Jungfermann-Guzman R, et al. Efficacy of tamsulosin, oxybutynin, and their combination in the control of double-j stent-related lower urinary tract symptoms. *Int Braz J Urol* 2016;42:487-493.
27. Norris RD, Sur RL, Springhart WP, Marguet CG, Mathias BJ, Pietrow PK, Albala DM, Preminger GM. A prospective, randomized, double-blinded placebo-controlled comparison of extended release oxybutynin versus phenazopyridine for the management of postoperative ureteral stent discomfort. *Urology* 2008;71:792-795.

Address correspondence to:

Ryan S. Hsi, MD

Department of Urology

Vanderbilt University Medical Center

A-1302 Medical Center North

Nashville, TN 37232

E-mail: ryan.hsi@vumc.org

Abbreviations Used

ASA % American Society of Anesthesiologists

BMI % body mass index

MED % morphine equivalent dose

TN CSMD % Tennessee Controlled Substance Monitoring Database

UTC % unable to calculate