Thresholds, spectral geometry, and persistent radiation in mass-critical dispersive equations

Research continuation: analytic proofs and independent internal review

8 October 2026

Abstract

We develop five connected results for critical focusing dispersive equations. For homogeneous real-even elliptic symbols of degree $m > 1$, a sequence of negative-energy traveling waves minimizing mass is compact after scaling and translation, except for a finite splitting into zero-energy exact profiles; no diffuse remainder survives. Negative energy forces at least three negative directions in the full real action Hessian, and momentum response gives stronger branch-dependent bounds. A strictly convex density-norm argument unifies phase-kernel simplicity for local powers and positive-type Hartree attraction and yields a uniform nonlocal traveling-profile mass gap. For same-order dispersions interpolating between two homogeneous limits, we determine both variational thresholds, classify the rational model's upper endpoint, and prove a sharp lower-threshold selection law by a variational limit on the optimizer manifold. Finally, any bounded global zero-momentum negative-energy BNLS orbit must sustain exterior gradient at distances proportional to time. These are profile, variational, spectral-form, and necessary dynamical results; they do not constitute a general classification of near-critical time-dependent solutions.

Status. The proofs below have undergone analytic cross-checking within this research session. No new numerical certificate is used. External peer review and exhaustive literature-priority checks have not been completed. Conditional stability-index statements are explicitly separated from unconditional quadratic-form conclusions.

What has advanced

The original physical question concerns pure BNLS, $iu_t = \Delta^2 u - |u|^{8/N}u$, and the relationship between negative energy and collapse without radial symmetry. The present continuation studies the structures that distinguish persistent coherent motion from collapse, and the changes in those structures under modified dispersion or nonlocal attraction.

1. **The pure traveling-wave threshold.** When negative-energy exact profiles exist, define their least possible mass by M_{tr}. A normalized minimizing sequence either converges to a negative-energy minimizer or splits strongly into finitely many zero-energy profiles with common parameters. If $M_{\mathrm{tr}} < 2M_c$, the latter alternative has only one component. The theorem does not decide the endpoint sign or the value of M_{tr}.
2. **A spectral requirement at every mass.** Negative energy forces a three-dimensional negative plane in the full action Hessian. On a smooth isotropic branch with momentum parallel to velocity it forces at least $N + 2$ negative directions. The distinction between near-ground and excited branches is thus expressed by precise spectral conditions, rather than by an assumption that every coherent wave has the same sign of energy.
3. **A nonlocal extension with a common proof mechanism.** The simplicity of the phase kernel follows from equality in a strictly convex density norm and sharp critical interpolation. For positive-type Hartree interactions this gives the complete analytic input for a uniform

negative-energy profile mass gap, even with arbitrary admissible subhomogeneous dispersion additions.

4. **A complete variational model and selected scales.** For the rational same-order interpolation, the constrained energy has two exact mass thresholds and three different upper-endpoint behaviors. Near the lower threshold, the energy, kinetic scale, and frequency have explicit leading constants. The next low-frequency moment selects among all principal GN optimizers; uniqueness is unnecessary.
5. **A stronger necessary dynamical alternative.** A forward-global uniformly H^2-bounded pure BNLS solution with $E < 0$ and $P = 0$ must maintain nonzero time-averaged exterior gradient at a linear spatial scale. Gradient localization, including an averaged version, is incompatible with those assumptions. This does not exclude persistent splitting or radiation.

The common idea is to use the geometry of the density attraction to control phase directions, the response to dilation to identify energy thresholds, and the velocity–momentum response to distinguish coherent branches. The actual evolution introduces a separate transport question; a static profile theorem cannot supply that missing time-dependent step.

1 Analytic foundations and conventions

The Fourier transform is unitary, $D = -i\nabla$, and $P(u) = \operatorname{Im} \int \overline{u}\nabla u = \int \xi|\widehat{u}|^2$. The real Hilbert pairing is the real part of the complex L^2 pairing. Throughout the local-power arguments, $N \geq 1$, $m > 1$, $s = m/2$, $r = m/N$, and $p = 2 + 2r$. A principal symbol a is real, even, m-homogeneous, C^{N+2} off zero, and satisfies $c|\xi|^m \leq a(\xi) \leq C|\xi|^m$. Set $A(u) = \int a|\widehat{u}|^2$, $M(u) = \|u\|_2^2$, and $E(u) = A(u)/2 - \|u\|_p^p/p$. The sharp critical mass M_c is defined by

$$\|u\|_p^p \leq \frac{p}{2}\left(\frac{M(u)}{M_c}\right)^r A(u). \tag{1}$$

The traveling-wave convention $u(t,x) = e^{i\omega t}U(x - vt)$ gives $[a(D) + \omega + iv \cdot \nabla]U - |U|^{2r}U = 0$.

Lemma 1.1 (Normalized optimizer compactness)**.** *The inequality (1) is attained. If $M(f_n) \to M_c$, $A(f_n) \to 1$, and $\|f_n\|_p^p \to p/2$, then some translated subsequence converges strongly in H^s to an optimizer. The normalized optimizer set is compact modulo translations and phases.*

Proof. Sharp Sobolev interpolation gives a finite positive best constant, and amplitude and spatial scaling normalize a maximizing sequence as stated. Bounded energy-space sequences that vanish locally in mass converge to zero in L^p: use a bounded-overlap cover by unit balls, local Sobolev interpolation, and $2 < p < 2N/(N - m)$ when $N > m$ (all finite p otherwise). Thus translations give a nonzero weak limit Q and almost-everywhere convergence. Write $M_1 = M(Q) > 0$, $A_1 = A(Q) > 0$, $M_2 = M_c - M_1$, and $A_2 = 1 - A_1$. Quadratic splitting and Brezis–Lieb give

$$M_c^r \leq M_1^r A_1 + M_2^r A_2, \qquad A_1 + A_2 = 1.$$

If both masses are positive, the right side is strictly below M_c^r. Consequently $M_1 = M_c$ and then $A_1 = 1$. Weak convergence plus convergence of both quadratic norms gives strong H^s convergence. The same proof applies to a sequence of optimizers. Constant phase is proved by the density-norm lemma below; this compactness argument itself does not need a positivity-preserving kinetic operator. □

Lemma 1.2 (Weak-profile regularity and Pohozaev identities)**.** *Every weak H^s local-power profile belongs to $L^\infty \cap H^{m+1}$ and satisfies*

$$v \cdot P = 2mE, \qquad \omega M = rA + (2m - p)E. \tag{2}$$

No spatial moment is required.

Proof. Choose Λ so large that $a(\xi) - v \cdot \xi + \Lambda \asymp 1 + |\xi|^m$. The shifted resolvent maps L^q to $H^{m,q}$, $1 < q < \infty$, by the Mihlin multiplier theorem. For $N > m$, the starting exponent is $q_0 = 2N/(N-m)$. Splitting the nonlinearity into $|U| > 1$ and $|U| \le 1$, resolvent Sobolev improvement iterates the reciprocal exponent as $\beta_{j+1} = (1+2r)\beta_j - r$ with $\beta_0 = (1-r)/2$. The ideal iterates $\beta_j = 1/2 - (r/2)(1+2r)^j$ cross the boundedness threshold after finitely many steps. Inputs have exponent greater than one; intermediate exponents may be weakened to avoid endpoints. The linear contribution has the stronger improvement $\beta \mapsto \beta - r$ and is handled separately. When $N = m$ start at any sufficiently large finite exponent; when $N < m$ the initial space already embeds in L^∞. Now the nonlinearity is in L^2, giving $U \in H^m \subset H^1$. The C^1 map $z \mapsto |z|^{2r} z$, with bounded derivative on the range of U, sends U into H^1; one more resolvent application gives H^{m+1}.

Here are the cutoff details for the dilation calculation. Let $f_R(x) = x\chi(x/R)$ with χ compactly supported and equal to one near zero, and let $L_R = f_R \cdot \nabla + \operatorname{div} f_R/2$. The Fourier kernel of $[a(D), L_R]$ is, up to the transform constant,

$$\frac{i}{2}(a(\xi) - a(\eta))(\xi + \eta) \cdot \widehat{f}_R(\xi - \eta).$$

Splitting comparable and noncomparable frequencies gives

$$|a(\xi) - a(\eta)||\xi + \eta| \le C|\xi - \eta|(|\xi| + |\eta|)^m.$$

After division by $\langle \xi \rangle^s \langle \eta \rangle^s$, the convolution kernel is bounded by $C|k|\langle k \rangle^s R^{N+1} |\widehat{f}(Rk)|$, with a uniform L^1 norm for $R \ge 1$. Hence the commutators are uniformly bounded $H^s \to H^{-s}$. Schwartz approximation and the homogeneous symbol identity give $\langle U, [a(D), L_R]U \rangle \to mA(U)$. The drift commutator tends to $-v \cdot P$; its order is one and $s > 1/2$. In the nonlinear pairing integration by parts and dominated convergence give

$$\operatorname{Re} \int |U|^{p-2} \overline{U} L_R U \longrightarrow N(1/2 - 1/p)\|U\|_p^p = \frac{m}{p}\|U\|_p^p.$$

Testing the profile with $L_R U$ therefore gives $mE = v \cdot P/2$. Testing with U gives $A + \omega M - v \cdot P - \|U\|_p^p = 0$; combining these two equations proves (2). □

The same regularity proof holds for an individually fixed nonnegative order-m addition b with the stated Mihlin derivative bounds. Constants need not be uniform in that addition or in v. For nonhomogeneous minimizers, differentiation of their scalar mass-preserving dilation energy is legitimate directly by dominated convergence and minimality; differentiability of the dilation orbit as an H^s-valued map is not being assumed.

2 Compactification of the negative-energy traveling-wave threshold

Fix $N \ge 1$, $m > 1$, $s = m/2$, $r = m/N$, and $p = 2 + 2r$. Let a be a real, even, positive elliptic m-homogeneous symbol, of class C^{N+2} away from zero. Throughout this section,

$$M(U) = \|U\|_2^2, \quad A(U) = \int a(\xi)|\widehat{U}(\xi)|^2 \, d\xi, \quad E(U) = \tfrac{1}{2}A(U) - \tfrac{1}{p}\|U\|_p^p, \quad P(U) = \int \xi |\widehat{U}(\xi)|^2 \, d\xi.$$

Let M_c be the sharp mass in

$$\|f\|_p^p \le \frac{p}{2}\left(\frac{M(f)}{M_c}\right)^r A(f). \tag{3}$$

An exact traveling profile is a nonzero $U \in H^s$ satisfying

$$[a(D) + \omega + iv \cdot \nabla]U - |U|^{p-2}U = 0. \tag{4}$$

The cutoff Pohozaev identities established for this equation are

$$v \cdot P(U) = 2mE(U), \qquad \omega M(U) = rA(U) + (2m-p)E(U). \tag{5}$$

Assume the set of its negative-energy traveling profiles is nonempty, and define

$$M_{\mathrm{tr}} = \inf\{M(U) : U \text{ satisfies (4) for some } (\omega, v),\ E(U) < 0\}.$$

Then $M_c \le M_{\mathrm{tr}} < \infty$ by (3). The earlier phase argument strengthens this to $M_{\mathrm{tr}} > M_c$, but the compactification proof below does not need that improvement, optimizer uniqueness, or real nondegeneracy.

Lemma 2.1 (Group velocity without a spatial moment)**.** *Every profile (4) satisfies*

$$vM(U) = \int \nabla a(\xi) |\widehat{U}(\xi)|^2 \, d\xi. \tag{6}$$

In particular, with $C_a = \sup_{\xi \neq 0} |\nabla a(\xi)| / a(\xi)^{(m-1)/m}$,

$$|v| \le C_a \left(\frac{A(U)}{M(U)} \right)^{(m-1)/m}. \tag{7}$$

For $a(\xi) = |\xi|^4$, $C_a = 4$.

Proof. The profile is a critical point of $S = E + \omega M/2 - v \cdot P/2$. Choose real $\chi \in C_c^\infty$ equal to x_j near zero and put $\chi_R(x) = R\chi(x/R)$. Differentiating $S(e^{i\theta\chi_R}U)$ at $\theta = 0$ gives

$$2\,\mathrm{Re}\langle a(D)U, i\chi_R U\rangle = \int v \cdot \nabla\chi_R \, |U|^2.$$

The right side tends to $v_j M(U)$. The commutator form on the left has Fourier kernel proportional to

$$(a(\xi) - a(\eta)) R^{N+1} \widehat{\chi}(R(\xi - \eta)).$$

The mean-value bound $|a(\xi) - a(\eta)| \le C|\xi - \eta|(|\xi| + |\eta|)^{m-1}$, and a division into comparable and noncomparable frequency sizes, show that after division by $\langle\xi\rangle^s \langle\eta\rangle^s$ the kernel is bounded by

$$C|\xi - \eta| \langle \xi - \eta \rangle^s R^{N+1} |\widehat{\chi}(R(\xi - \eta))|.$$

Its convolution L^1 norm is uniform for $R \ge 1$ by the substitution $q = R(\xi - \eta)$. Thus the commutator forms are uniformly bounded on H^s. For Schwartz U, passage to the limit is justified by $\chi_R U \to x_j U$ in L^2, and Fourier differentiation gives $\int \partial_j a \, |\widehat{U}|^2$. Density and the uniform form bound extend this to every H^s profile. Hölder's inequality then gives

$$|v| M(U) \le C_a A(U)^{(m-1)/m} M(U)^{1/m}.$$

The proof never assumes differentiability of $\eta \mapsto e^{i\eta\cdot x}U$ in H^s. □

Theorem 2.2 (Finite zero-energy compactification)**.** *Let U_n be negative-energy exact traveling profiles with $M(U_n) \to M_{\mathrm{tr}}$. Apply mass-preserving scaling so that $A(U_n) = 1$, and scale their frequencies and velocities accordingly. After passage to a subsequence, precisely one of the following alternatives holds.*

1. *There are translations x_n and an exact traveling profile U_* such that $U_n(\cdot + x_n) \to U_*$ strongly in H^s, with*

$$M(U_*) = M_{\mathrm{tr}}, \qquad A(U_*) = 1, \qquad E(U_*) < 0.$$

The profile parameters converge to those of U_*.

2. $E(U_n) \to 0$. *There are an integer* $1 \le J \le \lfloor M_{\rm tr}/M_c \rfloor$, *nonzero profiles* $V_1, \dots, V_J$, *and pairwise separating translations* x_n^j *such that*

$$\left\| U_n - \sum_{j=1}^{J} V_j(\cdot - x_n^j) \right\|_{H^s} \longrightarrow 0. \tag{8}$$

All profiles solve (4) *with the same limiting parameters* (ω_*, v_*), *and*

$$E(V_j) = 0, \qquad M(V_j) \ge M_c, \qquad \sum_j M(V_j) = M_{\rm tr}, \qquad \sum_j A(V_j) = 1, \tag{9}$$

$$\omega_* = \frac{r}{M_{\rm tr}} > 0, \qquad v_* \cdot P(V_j) = 0, \qquad \omega_* M(V_j) = r A(V_j). \tag{10}$$

In particular, no diffuse mass or kinetic remainder survives at this threshold. The second alternative is a statement about a sequence of exact profiles, not a time-dependent soliton-resolution theorem.

Proof. The normalization is $U_\lambda(x) = \lambda^{N/2} U(\lambda x)$ with $\lambda = A(U)^{-1/m}$. The parameters become $\omega_\lambda = \lambda^m \omega$ and $v_\lambda = \lambda^{m-1} v$, and $E(U_\lambda) = \lambda^m E(U)$. By (3),

$$-\frac{1}{2}\left[\left(\frac{M(U_n)}{M_c}\right)^r - 1\right] \le E(U_n) < 0.$$

Lemma 2.1 bounds v_n, and (5) bounds ω_n. Take a subsequence with $E(U_n) \to e \le 0$, $v_n \to v_*$, and $\omega_n \to \omega_*$. Then

$$\omega_* M_{\rm tr} = r + (2m - p)e. \tag{11}$$

Apply the standard translation profile decomposition in H^s. The exponent p lies strictly between 2 and the Sobolev endpoint. The resulting nonzero profiles V_j have separating translation parameters, quadratic norm decoupling, and a remainder vanishing in L^p in the iterated limit: $\lim_{J\to\infty} \limsup_{n\to\infty} \|R_n^J\|_p = 0$. No small energy-space remainder is assumed at this stage. Therefore

$$\sum_j M(V_j) \le M_{\rm tr}, \qquad \sum_j A(V_j) \le 1, \qquad \lim_n \|U_n\|_p^p = \sum_j \|V_j\|_p^p. \tag{12}$$

Local compactness into L^p and convergence of the parameters allow passage to the equation for each translated weak limit. Hence each V_j is an exact profile for (ω_*, v_*). This step uses no positivity assumption on $a(\xi) + \omega_* - v_* \cdot \xi$.

If some $E(V_j) < 0$, the definition of $M_{\rm tr}$ and mass decoupling force $M(V_j) = M_{\rm tr}$ and strong L^2 convergence of $U_n(\cdot + x_n^j)$ to V_j. The differences w_n are bounded in H^s, so (3) gives $\|w_n\|_p \to 0$. Interpolation, using $m > 1$, gives $\|w_n\|_{H^{1/2}} \to 0$, hence $P(U_n) \to P(V_j)$. The first identity in (5) now yields $E(U_n) \to E(V_j)$. Convergence of the nonlinear norms then gives $A(U_n) \to A(V_j)$. Weak convergence and convergence of the Hilbert norms prove strong H^s convergence, establishing the first alternative.

If $e < 0$, a negative-energy profile must occur. Indeed, from (12),

$$e = \sum_j E(V_j) + \frac{1}{2}\left(1 - \sum_j A(V_j)\right). \tag{13}$$

The series converges because the quadratic and nonlinear norm series converge. If every $E(V_j) \ge 0$, its right side is nonnegative.

It remains to consider $e = 0$ with no negative-energy profile. All terms on the right of (13) are nonnegative. Therefore every nonzero profile has zero energy and $\sum_j A(V_j) = 1$. Sharp GN gives $M(V_j) \ge M_c$, so there are only finitely many such profiles, with the asserted bound on J. Their

kinetic energies sum to one, so $J \geq 1$. Equation (11) gives $\omega_* = r/M_{\mathrm{tr}} > 0$. Apply (5) to each zero-energy profile and sum:

$$\omega_* \sum_j M(V_j) = r \sum_j A(V_j) = r = \omega_* M_{\mathrm{tr}}.$$

Thus their masses also sum to M_{tr}. Both quadratic remainders vanish, proving (8). The remaining identities are the individual Pohozaev identities. □

Corollary 2.3 (Attainment criteria)**.** *If a threshold-minimizing sequence satisfies $-E(U_n)/A(U_n) \geq \eta > 0$, then M_{tr} is attained by a negative-energy traveling profile. If $M_{\mathrm{tr}} < 2M_c$, every normalized minimizing sequence has a strongly convergent translated subsequence to a single profile of mass M_{tr} and energy at most zero. The latter assertion does not decide whether the endpoint energy is strictly negative.*

Proposition 2.4 (Mass response along a traveling branch)**.** *Suppose $z \mapsto U_z$ is a C^2 branch of exact profiles at fixed frequency $\omega > 0$ and velocity ze, with $|e| = 1$ and real $U_0 = Q$. Put $\Pi(z) = e \cdot P(U_z)$ and $\kappa = \Pi'(0)$. Then*

$$\omega \frac{d}{dz} M(U_z) = \frac{(m-1)z\Pi'(z) - \Pi(z)}{m}, \tag{14}$$

and

$$E(U_z) = \frac{\kappa z^2}{2m} + o(z^2), \qquad M(U_z) = M(Q) + \frac{(m-2)\kappa z^2}{2m\omega} + o(z^2).$$

If $m > 2$ and $\kappa < 0$, the nearby nonzero branch therefore has both negative energy and mass below $M(Q)$; in particular $M_{\mathrm{tr}} < M(Q)$.

Proof. For $S_z = E + \omega M/2 - ze \cdot P/2$, criticality gives $\frac{d}{dz} S_z(U_z) = -\Pi(z)/2$. Substitute $E(U_z) = z\Pi(z)/(2m)$ from Pohozaev and rearrange to obtain (14). Since Q is real, $\Pi(0) = 0$. Taylor expansion and integration give the conclusions. In the validated four-dimensional cubic BNLS branch, $m = 4$, $\omega = 1$, and $\Pi(z) = 2Iz + O(z^3)$ with $I < 0$, so $M(U_z) = M(Q) + Iz^2/2 + O(z^4)$ and $M(U_z) - M(Q) = 2E(U_z) + O(z^4)$. □

3 Negative energy, Morse index, and the momentum response matrix

This section distinguishes two statements. The negative-index obstruction below holds for every exact profile and requires no nondegeneracy. Statements about derivatives with respect to velocity apply only when a differentiable family of profiles has been supplied. Such a family is not presumed to exist at every profile.

Let a be a real-even positive elliptic homogeneous symbol of degree $m > 1$, let $X = H^{m/2}(\mathbb{R}^N; \mathbb{C})$, and set

$$\alpha = \frac{2m}{N}, \qquad E(U) = \frac{1}{2} \int a(\xi) |\widehat{U}(\xi)|^2 \, d\xi - \frac{1}{\alpha + 2} \int |U|^{\alpha+2}, \qquad P(U) = \operatorname{Im} \int \overline{U} \nabla U.$$

Suppose that a nonzero $U \in X$ solves

$$a(D)U + \omega U + iv \cdot \nabla U - |U|^\alpha U = 0. \tag{15}$$

The mass-critical Pohozaev identity is $v \cdot P(U) = 2mE(U)$. All pairings below are form pairings on X and X^*; in particular the derivative pairings are defined since $m > 1$. Our complex pairing is antilinear in its first argument, and the real Hilbert pairing is its real part.

Define the real scalar operator and the complex-linear operator

$$A_U = a(D) + \omega - |U|^\alpha, \qquad K_U = A_U + iv \cdot \nabla.$$

Both are selfadjoint as forms on the complex space, and A_U commutes with complex conjugation. The Hessian of $S_{\omega,v} = E + \omega M/2 - v \cdot P/2$ is a *real-linear* operator $\mathcal{L}_U$, whose quadratic form is

$$q_{\mathcal{L}_U}(\eta) = \mathrm{Re}\langle \eta, K_U \eta \rangle - \alpha \int |U|^{\alpha-2} (\mathrm{Re}(\overline{U}\eta))^2. \tag{16}$$

At $U = 0$ the last integrand is defined to be zero. It is bounded by $|U|^\alpha |\eta|^2$ and defines a continuous form even when $\alpha < 2$. The notation $n_-(\mathcal{L})$ denotes the maximal dimension of a real subspace on which its quadratic form is strictly negative. It does not denote the number of exponentially growing modes of the evolution.

3.1 An unconditional negative-index obstruction

Theorem 3.1 (Two scalar directions and three action directions)**.** *If $E(U) < 0$, then*

$$n_-(A_U|_{X_\mathbb{R}}) \geq 2, \qquad n_-(\mathcal{L}_U) \geq 3. \tag{17}$$

More generally these conclusions require only $v \cdot P(U) < 0$ and the profile equation; homogeneity and criticality are used solely to infer that sign from the energy.

Consequently, a profile whose full action Hessian has negative index at most two cannot have negative energy. These statements are uniform in frequency and speed and do not require proximity to a ground state.

Proof. Write $U = h + ib$ with real h, b. The profile equation gives

$$A_U h = v \cdot \nabla b, \qquad A_U b = -v \cdot \nabla h.$$

Antisymmetry of the derivative and $P(U) = -2\langle b, \nabla h \rangle$ imply

$$\langle h, A_U h \rangle = \langle b, A_U b \rangle = \frac{1}{2} v \cdot P(U), \qquad \langle h, A_U b \rangle = 0. \tag{18}$$

The functions h, b are linearly independent: dependence would make U a constant phase times a real function and would force $P(U) = 0$. Thus their real span is a two-dimensional negative subspace.

For the full Hessian it is useful to retain complex notation. Since $K_U U = 0$, conjugation of the profile equation gives

$$K_U \overline{U} = 2iv \cdot \nabla \overline{U}, \qquad \langle \overline{U}, K_U \overline{U} \rangle = 2v \cdot P(U).$$

The latter pairing is real. Selfadjointness and $K_U U = 0$ show that all K_U form pairings between U and $\overline{U}$ vanish. Moreover, U and $\overline{U}$ are linearly independent over $\mathbb{C}$, for the same constant-phase reason as above.

For $t \in \mathbb{R}$ and $z \in \mathbb{C}$ put $\eta = tU + z\overline{U}$. Equation (16) becomes

$$q_{\mathcal{L}_U}(tU + z\overline{U}) = 2v \cdot P(U)|z|^2 - \alpha \int |U|^{\alpha-2} \left(t|U|^2 + \mathrm{Re}(z\overline{U}^2) \right)^2. \tag{19}$$

If $z \neq 0$ the first term is strictly negative. If $z = 0$ and $t \neq 0$, the second term equals $-\alpha t^2 \|U\|_{\alpha+2}^{\alpha+2} < 0$. The real span of $U, \overline{U}, i\overline{U}$ is therefore a three-dimensional negative subspace. On the full four-dimensional real space $\mathrm{span}_\mathbb{C}\{U, \overline{U}\}$, the restricted form has precisely three negative directions and the one-dimensional kernel spanned by iU. □

Remark 3.2 (What the obstruction does and does not say). The proof is an explicit finite-dimensional compression of the Hessian, not a computation of its full spectrum. If the essential spectrum reaches negative values, the variational index may be infinite; three discrete negative eigenvalues are not being asserted. In particular it does not by itself prove spectral or nonlinear instability. The same argument applies to a real dispersion operator and a gauge-invariant nonlinearity $f(|u|^2)u$ whenever $f' \geq 0$ on the range of $|U|^2$ and $\int f'(|U|^2)|U|^4 > 0$: the last term in (16) is then replaced by $-2\int f'(|U|^2)(\mathrm{Re}(\overline{U}\eta))^2$. The implication from negative energy to $v \cdot P < 0$ must be checked separately for such a model. No such monotonicity is asserted for every nonlinear correction in the preceding dilation cone.

Remark 3.3 (Positive-type Hartree interaction). The full index bound also extends to the critical Hartree setting with a positive interaction form D, under the Hartree form and regularity assumptions used in this manuscript. There the frozen complex-linear operator still obeys $K_U U = 0$ and $K_U\overline{U} = 2iv \cdot \nabla\overline{U}$, while

$$q_{\mathcal{L}_U}(\eta) = \langle \eta, K_U\eta\rangle_{\mathbb{R}} - 2D(\mathrm{Re}(\overline{U}\eta), \mathrm{Re}(\overline{U}\eta)).$$

Positivity of D makes the correction nonpositive, and its value on the amplitude direction is strictly negative when $D(|U|^2, |U|^2) > 0$. The proof on $\mathrm{span}_{\mathbb{R}}\{U, \overline{U}, i\overline{U}\}$ is unchanged. A nonnegative subhomogeneous dispersion addition preserves the implication from $E < 0$ because its dilation identity is $v \cdot P = 2mE - D_b < 0$. Thus $n_-(\mathcal{L}_U) \geq 3$ also holds in that class. The conclusion uses positivity of the interaction form, not merely pointwise nonnegativity of its kernel.

Corollary 3.4 (A continuation barrier from a ground state). *A real sharp mass-critical ground state has full action Hessian negative index one. A continuous Fredholm family of profiles starting at such a state cannot reach a negative-energy profile while retaining only its phase and translation kernel and keeping all other spectral values uniformly separated from zero. If the essential spectral gap stays open, reaching a negative-energy profile requires a net gain of at least two negative directions through additional zero eigenvalues, counted with multiplicity.*

Proof. At a real optimizer Q, the phase block is nonnegative. On the real mass tangent space $\{h : \langle Q, h\rangle = 0\}$ the real block is nonnegative as well: E is nonnegative on the mass sphere $M = M_c$ and vanishes at Q, so its constrained second variation is nonnegative. On Q itself the real block has quadratic form $-\alpha\|Q\|_{\alpha+2}^{\alpha+2} < 0$. Hence the full index is one. The index is locally constant for continuous selfadjoint Fredholm forms after quotienting a kernel of fixed dimension, as long as the remaining spectrum is separated from zero. Apply Theorem 3.1 at the proposed endpoint. □

3.2 Additional negative directions from momentum response

Assume here that $v \mapsto U_v$ is a C^1 family in X of exact profiles at fixed ω, and that differentiating the profile equation is justified in X^*. Define

$$B = D_v P(U_v).$$

The matrix is symmetric: if $d(v) = S_{\omega,v}(U_v)$, then $d_v = -P(U_v)/2$ and $B = -2d_{vv}$.

Theorem 3.5 (A response-refined index bound). *At a negative-energy member of this family,*

$$n_-(\mathcal{L}_{U_v}) \geq 3 + n_-(B|_{P(U_v)^\perp}). \tag{20}$$

For an isotropic, rotationally equivariant family with $P(U_v) = p(c)v/c$, $c = |v| > 0$, this gives

$$n_-(\mathcal{L}_{U_v}) \geq N + 2. \tag{21}$$

Consequently such a wave is not a local energy minimum on the joint mass and momentum level set, provided that level set is a C^2 manifold near the wave.

Proof. Let $w_\zeta = D_v U_v[\zeta]$. Profile differentiation gives

$$\mathcal{L}_{U_v} w_\zeta = -i\zeta \cdot \nabla U_v, \qquad \langle w_\eta, \mathcal{L}_{U_v} w_\zeta \rangle_{\mathbb{R}} = \tfrac{1}{2} \eta^T B \zeta.$$

The second formula uses $DP(U)[w] = 2\,\mathrm{Re}\langle w, -i\nabla U\rangle$. The pairings of $\mathcal{L}w_\zeta$ with $\overline{U}$ and $i\overline{U}$ vanish because $\int U \partial_j U = 0$. Its pairing with U is $\zeta \cdot P(U)$, which vanishes for $\zeta \perp P(U)$. Thus the image of a negative subspace of $B|_{P^\perp}$ is negative and form-orthogonal to the three-dimensional space in (19). The map is injective on that subspace, since its quadratic form is strictly negative. The two negative spaces have trivial intersection, proving (20).

In the isotropic case, with $e = v/c$,

$$B = p'(c) e \otimes e + \frac{p(c)}{c}(I - e \otimes e).$$

Negative energy gives $cp(c) = v \cdot P < 0$, so $p(c) < 0$ and $B|_{P^\perp} = (p(c)/c)I$ has $N-1$ negative directions. This proves (21). The joint mass and momentum tangent space has codimension at most $N+1$, so it intersects the $(N+2)$-dimensional negative subspace nontrivially. At a constrained critical point the corresponding second variation of E is the restriction of $\mathcal{L}$. If the joint level set is a C^2 manifold, this negative tangent direction is realized by a constraint-preserving curve and excludes a local minimum. □

The last level-set assumption is not a hidden spectral hypothesis. For a sufficiently regular nonzero profile its charge differentials are independent: a real relation between them would give $(\tau - i\zeta \cdot \nabla)U = 0$, forcing its L^2 Fourier transform onto a hyperplane (or forcing $U = 0$ if $\zeta = 0$). The finite-dimensional implicit-function theorem then supplies the local charge manifold in the energy space.

3.3 An exact reduction of the charge Hessian

Suppose now that $(\omega, v) \mapsto U_{\omega,v}$ is a C^1 profile family on an open parameter set, that $d(\omega, v) = S_{\omega,v}(U_{\omega,v})$ is C^2, and that its charges obey the critical scaling covariance

$$\begin{aligned} M(U_{\lambda^m \omega, \lambda^{m-1} v}) &= M(U_{\omega,v}), \\ P(U_{\lambda^m \omega, \lambda^{m-1} v}) &= \lambda P(U_{\omega,v}) \end{aligned}$$

whenever the parameters are in the domain. This follows, in particular, if the family is closed under the usual critical dilation, up to phase and translation. Let $H = d''(\omega, v)$ and $B = D_v P(U_{\omega,v})$.

Theorem 3.6 (Bordered momentum formula)**.** *If $\omega \neq 0$, the symmetric matrices H and*

$$-\frac{1}{2}\begin{pmatrix} (m-1)v \cdot P & P^T \\ P & B \end{pmatrix} \tag{22}$$

are congruent. If B is invertible, define

$$\kappa = P^T B^{-1} P - (m-1) v \cdot P. \tag{23}$$

Then, writing $p(H)$ for the number of positive eigenvalues,

$$p(H) = n_-(B) + \mathbf{1}_{\{\kappa > 0\}}, \qquad \dim\ker H = \mathbf{1}_{\{\kappa = 0\}}. \tag{24}$$

The Schur complement of the v block in H is exactly $\kappa/(2m^2\omega^2)$.

Proof. Stationarity gives $d_\omega = M/2$ and $d_v = -P/2$. Put $D = (m\omega, (m-1)v)^T$. Differentiating the two scaling laws gives the exact column identity

$$HD = \begin{pmatrix} 0 \\ -P/2 \end{pmatrix}, \qquad H_{vv} = -B/2.$$

The invertible matrix

$$R = \begin{pmatrix} m\omega & 0 \\ (m-1)v & I \end{pmatrix}$$

therefore satisfies $R^T H R =$ (22). Sylvester's law of inertia and the Schur complement of B prove (24). For the final assertion, the mixed block is

$$H_{v\omega} = -\frac{P - (m-1)Bv}{2m\omega},$$

and the upper-left entry follows from $HD = (0, -P/2)^T$. Substitution and cancellation give $H_{\omega\omega} - H_{\omega v} H_{vv}^{-1} H_{v\omega} = \kappa/(2m^2\omega^2)$. □

For an isotropic family $P = p(c)e$ with $p(c)p'(c) \neq 0$,

$$\kappa = \frac{p(c)^2}{p'(c)} - (m-1)cp(c). \tag{25}$$

Thus charge-Hessian degeneracy occurs precisely at $cp'(c)/p(c) = 1/(m-1)$, under these nonvanishing assumptions. This is a degeneracy criterion, not an assertion that every such point changes dynamical stability. The momentum and its derivative at a fixed frequency suffice; a separate numerical differentiation in frequency is unnecessary.

3.4 The small-speed transition at dispersion order two

Assume a C^1 branch at $\omega = 1$ starts at a real standing profile Q. Assume $L_- = a(D) + 1 - |Q|^\alpha$ has kernel $\mathbb{R}Q$ and a bounded inverse on its orthogonal complement. Assume also that the real-part derivative of the branch at zero vanishes after the chosen translation gauge. One sufficient condition for this last assertion is $\ker L_+ = \operatorname{span}\{\partial_j Q\}$ together with the translation-orthogonal slice; neither this nondegeneracy nor branch existence is asserted for all ground states. Define

$$T_{ij} = \langle \partial_i Q, L_-^{-1} \partial_j Q \rangle.$$

If T has k negative eigenvalues, then

$$n_-(L_-) \geq k, \qquad n_-\big(\operatorname{diag}(L_+, L_-)\big) \geq 2k + 1. \tag{26}$$

Indeed, on the inverse-response vectors $L_-^{-1}(\zeta \cdot \nabla Q)$ the L_- form is exactly $\zeta^T T \zeta$. Let V be the resulting k-dimensional negative subspace. The form of L_- is negative on V and vanishes on Q, with no cross term. Since $L_+ = L_- - \alpha|Q|^\alpha$, it is strictly negative on $V \oplus \mathbb{R}Q$: for a nonzero V component the first form is already negative, and on a nonzero multiple of Q the subtracted potential term is strictly positive. The two diagonal blocks then give (26). Finite strictly negative subspaces persist under form-continuous perturbations, so this lower bound persists on sufficiently nearby traveling profiles. In particular an isotropic negative response $T = I_0 I_N$, $I_0 < 0$, requires at least $2N + 1$ negative action directions on the standing state and its nearby traveling branch.

Differentiating the imaginary equation gives $D_{v_j} U|_{v=0} = -iL_-^{-1}\partial_j Q$, and hence

$$P(U_{1,v}) = 2Tv + o(|v|), \qquad B(1, v) = 2T + o(1). \tag{27}$$

For a nonsingular T and directions on which $v^T T v$ stays away from zero relative to $|v|^2$, the preceding exact formulas yield

$$E(U_{1,v}) = \frac{1}{m} v^T T v + o(|v|^2), \tag{28}$$

$$M(U_{1,v}) = M(Q) + \frac{m-2}{m} v^T T v + o(|v|^2), \tag{29}$$

$$\kappa(1, v) = 2(2-m) v^T T v + o(|v|^2). \tag{30}$$

For (29), integrate $d_v = -P/2$ to get $d(1, v) = M(Q)/2 - v^T T v/2 + o(|v|^2)$ and use weighted homogeneity, $d = d_\omega + (m-1) v \cdot d_v/m$ at $\omega = 1$. Only the stated C^1 branch and C^2 action regularity are used; no fourth derivative of a noninteger power nonlinearity is assumed.

At a sharp ground state T is positive definite by phase coercivity. For sufficiently small nonzero velocity the charge Hessian is consequently negative definite if $m > 2$, whereas it has exactly one positive eigenvalue if $1 < m < 2$. At $m = 2$ this leading-order test vanishes. General homogeneous symbols of degree two need not be quadratic forms, so no exact Galilean degeneracy is inferred for all such symbols.

These are charge and index statements. Under the additional hypotheses of a Hamiltonian index theorem—a closed Hamiltonian linearization, finite negative index, a coercive complementary subspace, exactly the phase and translation kernel, independent charges, and an invertible charge Hessian—the standard index formula reads

$$k_r + 2k_c + 2k_i^- = n_-(\mathcal{L}) - p(H) = n_-(\mathcal{L}) - n_-(B) - \mathbf{1}_{\{\kappa > 0\}}. \tag{31}$$

Here the terms on the left count real unstable eigenvalues, complex unstable pairs, and the negative index on nonzero imaginary modes, respectively, with the conventions of the chosen index theorem. This last formula invokes the established Hamiltonian index theory of Lin and Zeng [5]; it is not a replacement for checking its operator hypotheses. For example, if a nondegenerate small-speed ground-state branch retains $n_-(\mathcal{L}) = 1$ and all those hypotheses hold, the $m > 2$ case has index one and therefore one positive real eigenvalue. No unconditional nonlinear stability or instability conclusion is made for arbitrary branches.

4 A density-norm principle and nonlocal critical attraction

The phase-kernel argument has a first-order formulation that applies to both local powers and nonlocal Hartree interactions. It avoids second derivatives of the nonlinear density at nodal sets.

4.1 The common variational lemma

Let A be a positive quadratic form on a complex function space, with $A(X + iY) = A(X) + A(Y)$ for real X, Y, and let $M(u) = \|u\|_2^2$. Let $\mathcal{B}$ be a strictly convex real normed space containing the squared densities under consideration. Fix $\kappa > 1$ and set

$$\mathcal{F}(u) = \mathfrak{f}(|u|^2), \qquad \mathfrak{f}(\rho) = \|\rho\|_{\mathcal{B}}^{\kappa}.$$

Assume that the sharp inequality

$$\mathcal{F}(u) \le \kappa \left(\frac{M(u)}{M_c} \right)^{\kappa - 1} A(u) \tag{32}$$

has a nonzero optimizer. Assume the relevant density norm is Gateaux differentiable at the squared density of each optimizer. For a real optimizer Q of mass M_c, write its derivative as

$$D\mathfrak{f}(Q^2)[h] = \kappa \langle V_Q, h \rangle.$$

The notation allows either a multiplication potential or a bounded linear functional on the density space; in particular $\langle V_Q, b^2 \rangle$ must be a well-defined quadratic form.

Lemma 4.1 (Simple phase kernel from density convexity)**.** *Under these assumptions every complex optimizer is a constant phase times a real function. For a real mass-M_c optimizer Q, the form*

$$\mathcal{L}_-(b) = A(b) + (\kappa - 1)\frac{A(Q)}{M_c}M(b) - \langle V_Q, b^2\rangle$$

is nonnegative on real functions and has kernel $\operatorname{span}_{\mathbb{R}}\{Q\}$.

Proof. For $u = X + iY$, the density triangle inequality, sharp interpolation, and two-term Hölder give

$$\begin{aligned}\mathcal{F}(u)^{1/\kappa} &\le \mathcal{F}(X)^{1/\kappa} + \mathcal{F}(Y)^{1/\kappa} \\ &\le \kappa^{1/\kappa} M_c^{-(\kappa-1)/\kappa}\left[A(X)^{1/\kappa}M(X)^{(\kappa-1)/\kappa} + A(Y)^{1/\kappa}M(Y)^{(\kappa-1)/\kappa}\right] \\ &\le \kappa^{1/\kappa} M_c^{-(\kappa-1)/\kappa} A(u)^{1/\kappa}M(u)^{(\kappa-1)/\kappa}.\end{aligned}$$

All inequalities are equalities for an optimizer. Strict convexity in the density space forces $Y^2 = cX^2$ unless a component vanishes. Apply the same assertion after multiplication by $e^{i\pi/4}$. If both signs of Y/X occurred on sets of positive measure, the rotated squared components would fail to be proportional; when $c = 1$, they would instead have disjoint nonzero supports. Thus one relative sign holds almost everywhere, proving constant phase. Here, as usual for a normed space of densities, equality in $\mathcal{B}$ identifies densities almost everywhere.

Put $M = M_c$, $A = A(Q)$, $B = M(b)$ and $C = A(b)$. For real b, (32) gives

$$\mathfrak{f}(Q^2 + tb^2) \le \kappa(1 + tB/M)^{\kappa-1}(A + tC), \qquad t \ge 0.$$

Differentiation at zero proves $\mathcal{L}_-(b) \ge 0$. The homogeneity of $\mathfrak{f}$ gives $\langle V_Q, Q^2\rangle = \mathfrak{f}(Q^2) = \kappa A$, so $\mathcal{L}_-(Q) = 0$.

If $b \ne 0$ belongs to the form kernel, the first-derivative bound is an equality. The supporting-functional bound for a norm power is

$$D\mathfrak{f}(Q^2)[b^2] \le \kappa\mathfrak{f}(Q^2)^{(\kappa-1)/\kappa}\mathfrak{f}(b^2)^{1/\kappa}.$$

Use $\mathfrak{f}(Q^2) = \kappa A$ and (32) for b. With $x = C/A$ and $y = B/M$, the kernel equality implies

$$x + (\kappa - 1)y \le \kappa x^{1/\kappa}y^{(\kappa-1)/\kappa} \le x + (\kappa - 1)y.$$

The last inequality is weighted arithmetic–geometric mean. Hence $x = y$ and equality holds in the supporting-functional bound. Strict convexity makes the supporting hyperplane meet the unit sphere at one point, giving $b^2 = yQ^2$. Since a nonnegative quadratic form has a linear kernel, apply this conclusion also to $b + Q$. Subtracting the squared identities proves $b = cQ$ with one real constant c. □

For the local critical power, take $\mathcal{B} = L^\kappa$ and $\kappa = 1 + m/N$. For a positive-type Hartree interaction, take the Hilbert space of densities in its interaction norm and $\kappa = 2$. Lemma 4.1 itself asserts neither compactness of near-optimizers nor a dynamical theorem. Those require additional analytic information, supplied for the Hartree model below.

Corollary 4.2 (The local-power profile gap)**.** *For the homogeneous local-power class of Section 1, there is $\delta > 0$ such that every negative-energy traveling profile has mass at least $M_c+\delta$. Consequently, when the negative-energy profile set is nonempty, $M_{\mathrm{tr}} > M_c$.*

Proof. For a real normalized optimizer Q, the preceding lemma with $\kappa = 1+r$ gives the nonnegative phase form $L_Q = a(D)+r/M_c-|Q|^{2r}$ with kernel $\mathbb{R}Q$. Multiplication by $|Q|^{2r}$ is compact $H^s \to H^{-s}$: Hölder with two L^p factors controls its tails, and local Rellich compactness controls its restriction to a ball. Hence $\langle h, L_Q h\rangle \ge c_Q\|h\|_{H^s}^2$ for real $h \perp Q$. If negative-energy profiles had masses

tending to M_c, normalize $A(U_n) = 1$. Sharp GN then gives $E(U_n) \to 0$ and convergence to a normalized optimizer modulo translations and phase. Make its limit Q real and choose phases so $Y_n = \operatorname{Im} U_n \perp Q$. Pohozaev gives $\omega_n \to r/M_c$, and strong H^s convergence gives convergence of $|U_n|^{2r}$ in potential form norm. The imaginary equation and $P(U_n) = -2\int Y_n \nabla \operatorname{Re} U_n$ give the exact identity

$$mE(U_n) = \langle Y_n, [a(D) + \omega_n - |U_n|^{2r}]Y_n\rangle \geq \tfrac{1}{2}c_Q\|Y_n\|_{H^s}^2 \geq 0$$

for large n, a contradiction. □

4.2 The critical Hartree class

Fix $1 < m < N$, set $s = m/2$, and let a be real, even, positive elliptic, C^{N+2} away from zero, and homogeneous of degree m:

$$c|\xi|^m \leq a(\xi) \leq C|\xi|^m.$$

Let K be real, even, C^1 away from zero, homogeneous of degree $-m$, and satisfy

$$|K(x)| + |x||\nabla K(x)| \leq C_K|x|^{-m}. \tag{33}$$

Assume its interaction form

$$\mathcal{D}(f,g) = \iint K(x-y)f(x)g(y)\,dx\,dy$$

is positive definite on the real density space in use. More explicitly, it suffices that the Fourier transform of K is a locally integrable, strictly positive function almost everywhere; then the Fourier representation of $\mathcal{D}$ gives the required inner product and identifies zero-norm densities with zero functions. Examples include $K(x) = |x|^{-m}$ and positive finite sums of $|B_jx|^{-m}$ for invertible real matrices B_j. Pointwise positivity of K is not needed for the proof.

Write

$$H(u) = \mathcal{D}(|u|^2, |u|^2), \qquad A(u) = \int a(\xi)|\widehat{u}(\xi)|^2\,d\xi.$$

Hardy–Littlewood–Sobolev and Sobolev interpolation give the finite sharp inequality

$$H(u) \leq \frac{2}{M_c}M(u)A(u). \tag{34}$$

Indeed, with $q = 4N/(2N-m)$, the interaction is bounded by $C\|u\|_q^4$, and interpolation gives $\|u\|_q^4 \leq CM(u)A(u)$.

Allow in addition any real-even symbol $b \geq 0$ satisfying

$$\xi\cdot\nabla b \leq mb, \qquad |\partial^\beta b(\xi)| \leq C_{b,\beta}|\xi|^{-|\beta|}(1+|\xi|^m), \quad |\beta| \leq N+2, \tag{35}$$

with $b \in C^{N+2}(\mathbb{R}^N \setminus \{0\})$. There is no common bound on the individual constants. Define

$$B_b(u) = \int b(\xi)|\widehat{u}(\xi)|^2\,d\xi, \qquad E_b(u) = \tfrac{1}{2}[A(u) + B_b(u)] - \tfrac{1}{4}H(u).$$

Theorem 4.3 (Nonlocal mass gap over the dispersion cone)**.** *There exists $\delta > 0$, depending only on N, m, a, K, such that for every admissible b, every $\omega \in \mathbb{R}$ and $v \in \mathbb{R}^N$, a nonzero weak $H^{m/2}$ solution of*

$$[a(D) + b(D) + \omega + iv\cdot\nabla]U - (K * |U|^2)U = 0 \tag{36}$$

with $E_b(U) < 0$ satisfies $M(U) \geq M_c + \delta$. The gap is uniform over the entire addition class, frequencies, and velocities. No radial symmetry, spatial moment, positivity or uniqueness of a ground state, or real-sector nondegeneracy is assumed.

4.3 Compactness and the phase form

Normalized maximizing sequences for (34) are compact modulo translation and constant phase. Here are the details needed for this nonlocal statement. Normalize $M(u_n) = M_c$, $A(u_n) = 1$, and $H(u_n) \to 2$. Since $H(u_n) \le C\|u_n\|_q^4$, local vanishing in mass is impossible by the localized Sobolev inequality. A translated subsequence has a nonzero weak H^s limit Q, with local strong L^q convergence. Set $r_n = u_n - Q$. The product Qr_n tends to zero in $L^{q/2}$ by local convergence and the small L^q tail of Q. Moreover,

$$\mathcal{D}(|Q|^2, |r_n|^2) \longrightarrow 0. \tag{37}$$

To see this, approximate Q in L^q by a smooth compactly supported function. Its potential is bounded and tends to zero at infinity, so multiplication by it is compact as a form $H^s \to H^{-s}$. The Hardy–Littlewood–Sobolev estimate controls the approximation error, proving (37). It also proves the nonlocal splitting

$$H(u_n) = H(Q) + H(r_n) + o(1).$$

If M_j, A_j are the limiting mass and kinetic contributions of the two pieces, (34) implies

$$M_c \le M_1 A_1 + M_2 A_2, \qquad M_1 + M_2 = M_c, \quad A_1 + A_2 = 1, \quad M_1 > 0.$$

Both masses cannot be positive. Thus $M_1 = M_c$, $A_1 = 1$, proving strong convergence and attainment. The same argument covers sequences with these normalization limits and all sequences of normalized optimizers.

Take $\mathcal{B}$ to be the completion of real densities under $\|\rho\|_{\mathcal{B}}^2 = \mathcal{D}(\rho, \rho)$. Lemma 4.1 shows that every optimizer has constant phase. A real optimizer with $M(Q) = M_c$, $A(Q) = 1$ satisfies

$$a(D)Q + \omega_* Q - (K * Q^2)Q = 0, \qquad \omega_* = 1/M_c.$$

The real phase form

$$L_Q = a(D) + \omega_* - K * Q^2$$

is nonnegative with kernel $\operatorname{span}_{\mathbb{R}}\{Q\}$. The potential is a compact form by the same approximation used in (37). Consequently

$$\langle h, L_Q h\rangle \ge c_Q \|h\|_{H^s}^2 \quad \text{for real } h \perp Q. \tag{38}$$

The constants can be taken uniform over normalized optimizers: otherwise translate a convergent sequence of optimizers and use form continuity to contradict (38) for its limit. Form continuity follows from HLS, since $u \mapsto |u|^2$ is continuous $H^s \to L^{q/2}$.

4.4 Weak-profile regularity and identities

Every weak profile in (36) belongs to $L^\infty \cap H^{m+1}$. For the fixed b, v, choose Λ large so that

$$a(\xi) + b(\xi) - v \cdot \xi + \Lambda \asymp 1 + |\xi|^m.$$

The resolvent maps L^p to $H^{m,p}$ for $1 < p < \infty$ by the individual Mihlin bounds. Split K into its parts inside and outside a fixed ball. The outside potential applied to $|U|^2$ is bounded. The inside kernel lies in L^j for $1 \le j < N/m$, while $|U|^2 \in L^{N/(N-m)}$. Young's inequality therefore puts the inside potential in arbitrarily large finite L^t spaces. Fix sufficiently large $t > N/m$. Multiplication by this potential, followed by the resolvent, improves a reciprocal integrability exponent by $m/N - 1/t > 0$. Starting from the energy Sobolev exponent, finitely many iterations give $U \in L^\infty$; intermediate exponents may be weakened to avoid endpoints. Bounded potential and linear terms improve at least as strongly and may be combined using interpolation with L^2.

Now $K * |U|^2$ is bounded, giving $U \in H^m \subset H^1$. Also $\nabla|U|^2 \in L^1 \cap L^2$, and the near/far kernel split gives $\nabla(K * |U|^2) \in L^2 + L^\infty$. Thus $(K * |U|^2)U \in H^1$, and the resolvent gives H^{m+1}.

Put $\rho = |U|^2$, $f_R(x) = x\chi(x/R)$ with smooth compactly supported $\chi = 1$ near zero, and $L_R = f_R \cdot \nabla + (\operatorname{div} f_R)/2$. Symmetrized integration by parts gives

$$\begin{aligned} &\operatorname{Re} \int (K * \rho) U \, \overline{L_R U} \, dx \\ &\quad = -\frac{1}{4} \iint \rho(x)\rho(y)[f_R(x) - f_R(y)] \cdot \nabla K(x-y) \, dx \, dy \longrightarrow \frac{m}{4} H(U). \end{aligned} \tag{39}$$

One can first regularize or exclude the diagonal. The omitted boundary term vanishes because $m < N$ after symmetrization. The uniform Lipschitz bound for f_R and (33) dominate the integrand by $C\rho(x)\rho(y)|x-y|^{-m}$, integrable by HLS, and justify the limit without a spatial moment.

For clarity, the kinetic commutator used here has kernel, up to Fourier normalization,

$$\frac{i}{2}[c(\xi) - c(\eta)](\xi + \eta) \cdot \widehat{f}_R(\xi - \eta), \qquad c = a + b.$$

The symbol assumptions give

$$|c(\xi) - c(\eta)||\xi + \eta| \le C|\xi - \eta|[1 + (|\xi| + |\eta|)^m].$$

Divide by the two energy weights and use $\widehat{f}_R(k) = R^{N+1}\widehat{f}(Rk)$ to get an integrable kernel majorant with a bound independent of R. The limiting form for Schwartz functions is $\int \xi \cdot \nabla c \, |\widehat{U}|^2$; density extends it to the energy space. The degree-one drift term has limit $-v \cdot P(U)$, where $P(U) = \int \xi |\widehat{U}|^2$ is finite because $m > 1$.

Abbreviate $A = A(U)$, $B = B_b(U)$, $M = M(U)$ and $E = E_b(U)$, and set

$$D_b = \int [mb(\xi) - \xi \cdot \nabla b(\xi)] |\widehat{U}(\xi)|^2 \, d\xi \ge 0.$$

Testing by $L_R U$, taking the limit, and testing separately by U gives

$$v \cdot P = 2mE - D_b, \qquad \omega M = A + B + (2m-4)E - D_b. \tag{40}$$

Write $U = X + iY$ with real components and put $\widetilde{\omega} = \omega + D_b/M$. The imaginary profile equation, together with $P(U) = -2\int Y \nabla X$, yields the exact shifted identity

$$\begin{aligned} mE &= \langle Y, [a(D) + \widetilde{\omega} - K * |U|^2] Y \rangle + B_b(Y) \\ &\quad + D_b \left(\frac{1}{2} - \frac{M(Y)}{M} \right), \qquad \widetilde{\omega} M = A + B + (2m-4)E. \end{aligned} \tag{41}$$

4.5 Proof of the uniform nonlocal gap

Proof of Theorem 4.3. Negative energy and (34) require $M > M_c$. If the gap failed, take profiles with $M(U_n) \to M_c$ and negative energies, allowing arbitrary admissible b_n, ω_n, v_n. The critical scaling

$$U_\lambda(x) = \lambda^{N/2} U(\lambda x), \quad b_\lambda(\xi) = \lambda^m b(\xi/\lambda), \quad \omega_\lambda = \lambda^m \omega, \quad v_\lambda = \lambda^{m-1} v$$

preserves the class, mass, and profile equation, and multiplies energy by λ^m. Normalize $A(U_n) = 1$. Writing $d_n = M(U_n)/M_c - 1$, the sharp bound gives

$$0 \le B_{b_n}(U_n) < d_n, \qquad -d_n/2 \le E_n < 0, \qquad H(U_n) \longrightarrow 2.$$

Compactness yields, after translations and phases, strong H^s convergence to a real normalized optimizer Q. A further phase makes $Y_n = \operatorname{Im} U_n$ orthogonal to Q. Equation (41) gives $\widetilde{\omega}_n \to 1/M_c$ even if D_{b_n} is unbounded. HLS gives convergence of the Hartree potentials in form norm. By (38), for large n,

$$\langle Y_n, [a(D) + \widetilde{\omega}_n - K * |U_n|^2] Y_n \rangle \ge c \|Y_n\|_{H^s}^2.$$

The other terms in (41) are nonnegative, since $M(Y_n)/M(U_n) \to 0$. Its right side is therefore nonnegative, contradicting $mE_n < 0$. □

The conclusion extends a local attraction mechanism to a nonlocal density interaction. It is a profile theorem, not a classification of arbitrary global trajectories. The positive-density geometry is an explicit hypothesis; arbitrary matrix couplings, non-even symbols, and indefinite interaction kernels are not covered.

5 Two dispersion thresholds and the selection of critical profiles

The loss of a profile gap under a small nonhomogeneous perturbation has a more precise variational description. We prove two distinct mass thresholds, classify the upper endpoint for a full interpolation family, and determine the scale and selected profiles as the lower threshold is approached. The selection argument requires neither uniqueness of the homogeneous optimizers nor nondegeneracy of their real linearization. These are constrained variational statements; they do not classify arbitrary solutions of the homogeneous evolution.

5.1 Symbols and the two thresholds

Fix $N \geq 1$, $m > 1$, $s = m/2$, $r = m/N$, and $p = 2 + 2r$. Let a_0 and a_∞ be real, even, positive elliptic symbols homogeneous of degree m, continuous on the sphere and C^{N+2} off zero. Suppose $a_\infty - a_0$ is also positive elliptic. Use the notation

$$M(f) = \|f\|_2^2, \qquad A_j(f) = \int a_j(\xi)|\widehat{f}(\xi)|^2\,d\xi, \qquad E_j(f) = \tfrac{1}{2}A_j(f) - \tfrac{1}{p}\|f\|_p^p \quad (j = 0, \infty).$$

The sharp homogeneous masses M_j are defined by

$$\frac{1}{p}\|f\|_p^p \leq \frac{1}{2}\left(\frac{M(f)}{M_j}\right)^r A_j(f). \tag{42}$$

In particular, $M_\infty > M_0$: if $a_\infty \geq (1+\eta)a_0$, sharpness gives $M_\infty \geq (1+\eta)^{1/r}M_0$.

Write the full symbol as $k = a_0 + b$, with b real, even, and continuous off zero. Assume that, for some $q > 0$ and constants $c, C > 0$,

$$c\min\{|\xi|^{m+q}, |\xi|^m\} \leq b(\xi) \leq C\min\{|\xi|^{m+q}, |\xi|^m\}, \tag{43}$$

and that, uniformly for $\theta \in S^{N-1}$,

$$R^{-m}k(R\theta) \longrightarrow a_\infty(\theta) \qquad (R \longrightarrow \infty). \tag{44}$$

Thus k is comparable with $|\xi|^m$. Define

$$B(f) = \int b(\xi)|\widehat{f}(\xi)|^2\,d\xi, \qquad E_k(f) = \tfrac{1}{2}[A_0(f) + B(f)] - \tfrac{1}{p}\|f\|_p^p, \qquad I(M) = \inf_{\substack{f\in H^s \\ M(f)=M}} E_k(f).$$

Theorem 5.1 (Two homogeneous limits determine two mass thresholds)**.** *Under (43)–(44), the sharp GN mass for the full form $A_0 + B$ is exactly M_0, and its sharp inequality has no nonzero optimizer. The constrained infima satisfy*

1. *$I(M) = 0$, without attainment, for $0 < M \leq M_0$;*
2. *$-\infty < I(M) < 0$ for $M_0 < M < M_\infty$. Every minimizing sequence has, after translations and extraction, a strong H^s limit attaining $I(M)$. A real minimizer can be chosen;*
3. *$I(M) = -\infty$ for $M > M_\infty$.*

The symbols a_0 and a_∞ need not have the same anisotropy.

Proof. Since $A_0 + B \geq A_0$, the full GN constant is at most the principal one. If Q is a principal optimizer and $Q_\lambda(x) = \lambda^{N/2} Q(\lambda x)$, then

$$\lambda^{-m} B(Q_\lambda) = \int \lambda^{-m} b(\lambda\xi) |\widehat{Q}(\xi)|^2 \, d\xi \longrightarrow 0 \qquad (\lambda \downarrow 0)$$

by dominated convergence and (43). Thus the sharp constants coincide. Since $B(f) > 0$ for every nonzero f, equality in the full inequality is impossible. At mass $M \leq M_0$, the sharp principal inequality gives $E_k(f) > 0$. Dilating any smooth mass-M trial to low frequency proves that the infimum is zero.

For every $\eta > 0$, (44) gives a finite C_η such that $k \geq (1-\eta) a_\infty - C_\eta$. Therefore, at fixed mass M,

$$E_k(f) \geq \frac{1}{2} \left[1 - \eta - (M/M_\infty)^r \right] A_\infty(f) - \frac{1}{2} C_\eta M. \tag{45}$$

If $M < M_\infty$, choose η to make the bracket positive. This proves a finite lower bound and boundedness of minimizing sequences. If also $M > M_0$, sharpness and amplitude scaling give a real smooth mass-M trial f with $E_0(f) < 0$. Its low-frequency dilation satisfies

$$E_k(f_\lambda) = \lambda^m E_0(f) + O(\lambda^{m+q}) < 0$$

for sufficiently small λ, proving $I(M) < 0$.

We record the compactness argument. Amplitude scaling gives

$$E_k(\sqrt{\vartheta} f) = \vartheta^{p/2} E_k(f) + \tfrac{1}{2}(\vartheta - \vartheta^{p/2})[A_0(f) + B(f)] \qquad (\vartheta > 1).$$

Thus $I(\vartheta L) \leq \vartheta I(L)$, with $I(\vartheta L) \leq \vartheta^{p/2} I(L) < \vartheta I(L)$ if $I(L) < 0$, whenever the masses are below M_∞. For a mass M with $I(M) < 0$, it follows that $I(M)/M < I(L)/L$ for every $0 < L < M$. Consequently

$$I(M) < I(L) + I(M - L) \qquad (0 < L < M). \tag{46}$$

A bounded minimizing sequence cannot vanish locally in mass: the localized GN inequality would then imply $\|f_n\|_p \to 0$, contradicting $I(M) < 0$. After translations there is a nonzero weak H^s limit Q and almost-everywhere convergence. The quadratic forms split by weak Hilbert convergence, and the nonlinear integral by the Brezis–Lieb lemma. If $0 < M(Q) < M$, normalize the remainder to its limiting mass; boundedness of its kinetic and nonlinear terms makes the energy change $o(1)$. Splitting then contradicts (46). Hence $M(Q) = M$. For the remainder $w_n = f_n - Q$, the principal GN inequality gives

$$\|w_n\|_p^p \leq C M(w_n)^r A_0(w_n) \longrightarrow 0.$$

Energy splitting now forces $A_0(w_n) + B(w_n) \to 0$, giving strong H^s convergence and attainment.

Realness does not require a modulus inequality for a higher-order operator. Write $f = X + iY$, put $\vartheta = M(X)/M(f)$, and, when both components are nonzero, set $g = X/\sqrt{\vartheta}$, $h = Y/\sqrt{1-\vartheta}$. These real functions have the same mass as f. Evenness of k splits its form over real and imaginary parts, while convexity gives

$$|f|^p \leq \vartheta |g|^p + (1-\vartheta)|h|^p, \qquad E_k(f) \geq \vartheta E_k(g) + (1-\vartheta) E_k(h).$$

Therefore a real minimizing sequence exists and its strong limit is real.

Finally, for $M > M_\infty$ choose a smooth mass-M function with $E_\infty(f) < 0$. High-frequency dilation and (44) give

$$\lambda^{-m} E_k(f_\lambda) \longrightarrow E_\infty(f) < 0 \qquad (\lambda \longrightarrow \infty),$$

using $k \leq C|\xi|^m$ for domination. Hence $I(M) = -\infty$. □

For each minimizer U in the open window, the multiplier equation and its scalar identity are

$$k(D)U + \omega U - |U|^{2r}U = 0, \qquad \omega M = r[A_0(U) + B(U)] - pI(M) > 0. \tag{47}$$

If additionally k is C^{N+2} off zero and satisfies the individual order-m derivative estimates

$$|\partial^\beta k(\xi)| \le C_\beta |\xi|^{m-|\beta|}, \qquad |\beta| \le N+2,$$

the elliptic bootstrap gives $U \in L^\infty \cap H^{m+1}$. A real minimizer has zero momentum and its standing-wave formula gives a global solution directly.

5.2 The complete isotropic upper-endpoint classification

Consider the interpolation

$$h = a_\infty - a_0, \qquad k_q(\xi) = a_0(\xi) + h(\xi)\frac{|\xi|^q}{1+|\xi|^q} = a_\infty(\xi) - d_q(\xi), \qquad d_q(\xi) = \frac{h(\xi)}{1+|\xi|^q}. \tag{48}$$

Write $D_q(f) = \int d_q |\widehat{f}|^2$. At mass M_∞,

$$E_{k_q}(f) = E_\infty(f) - \tfrac{1}{2}D_q(f), \qquad E_\infty(f) \ge 0. \tag{49}$$

Theorem 5.2 (Upper-endpoint trichotomy)**.** *For (48), $q < m$ implies $I(M_\infty) = -\infty$, whereas $q > m$ implies $-\infty < I(M_\infty) < 0$, attainment by a real minimizer, and precompactness of every minimizing sequence modulo translations. Both assertions allow anisotropic a_0, a_∞.*

If $a_0(\xi) = |\xi|^m$ and $a_\infty(\xi) = (1+\varepsilon)|\xi|^m$, then at $q = m$,

$$I(M_\infty) = -\tfrac{1}{2}\varepsilon M_\infty, \qquad M_\infty = (1+\varepsilon)^{N/m} M_0, \tag{50}$$

and the infimum is not attained.

Proof. Suppose $q < m$ and take an a_∞ optimizer Q of mass M_∞, so $E_\infty(Q) = 0$. Then

$$\lambda^{-(m-q)} D_q(Q_\lambda) = \int \frac{h(\xi)}{\lambda^{-q} + |\xi|^q} |\widehat{Q}(\xi)|^2 \, d\xi \longrightarrow \int h(\xi)|\xi|^{-q}|\widehat{Q}(\xi)|^2 \, d\xi > 0.$$

The limit is finite: $0 < m - q < m$, and the integrands are dominated by $C|\xi|^{m-q}|\widehat{Q}|^2$. Thus (49) tends to $-\infty$ along this sequence.

Suppose $q > m$. Then d_q is bounded and tends to zero at infinity, so (49) supplies the finite lower bound $-\|d_q\|_\infty M_\infty/2$. Negativity follows from the low-frequency trial since $M_\infty > M_0$. We must prove boundedness of minimizing sequences because (45) degenerates at this endpoint. If instead a subsequence satisfies $A_\infty(f_n) = t_n^m \to \infty$, write $f_n(x) = t_n^{N/2} V_n(t_n x)$. Then

$$M(V_n) = M_\infty, \quad A_\infty(V_n) = 1, \quad E_\infty(V_n) = t_n^{-m}[E_{k_q}(f_n) + D_q(f_n)/2] \longrightarrow 0.$$

Normalized sharp-GN compactness gives, after translations, $V_n \to Q$ strongly in H^s, where Q is an a_∞ optimizer. The bounded multiplier and strong L^2 convergence permit replacement of V_n by Q in

$$D_q(f_n) = \int d_q(t_n\xi)|\widehat{V}_n(\xi)|^2 \, d\xi.$$

The expression for Q tends to zero by dominated convergence. Thus $D_q(f_n) \to 0$ and (49) implies $\liminf E_{k_q}(f_n) \ge 0$, contradicting the negative infimum. Minimizing sequences are therefore bounded. Amplitude scaling proves (46) also at $M = M_\infty$, since both positive submasses lie strictly below it. The compactness and realness arguments of Theorem 5.1 now apply unchanged.

In the isotropic case $q = m$, $d_m(\xi) = \varepsilon|\xi|^m/(1+|\xi|^m) < \varepsilon$. Every admissible nonzero function consequently has $E_{k_m}(f) > -\varepsilon M_\infty/2$. Conversely, high-frequency dilations of an a_∞ optimizer have zero homogeneous energy and $D_m(Q_\lambda) \to \varepsilon M_\infty$ by dominated convergence. This proves the exact infimum and its nonattainment. □

For completeness, the anisotropic borderline $q = m$ cannot be replaced by the isotropic formula. Let $\mathcal{G}_\infty$ be the equality set normalized by $M = M_\infty$, $A_\infty = 1$. Define

$$J_{\mathrm{esc}} = -\frac{1}{2} \sup_{Q \in \mathcal{G}_\infty} \int h(\xi/|\xi|) |\widehat{Q}(\xi)|^2 \, d\xi.$$

The supremum is attained by normalized GN compactness and continuity of this bounded multiplier form. High-frequency optimizer trials give $I(M_\infty) \le J_{\mathrm{esc}}$. If this inequality is strict, minimizing sequences are bounded and attainment follows. Indeed an unbounded sequence, normalized exactly as in the preceding proof, has a limiting homogeneous optimizer and energy liminf at least J_{esc}. We make no assertion about attainment in the general anisotropic equality case.

Every minimizer of (48), including the attained upper endpoint $q > m$, obeys the exact identity

$$E_{k_q}(U) = -\frac{q}{2m} \int h(\xi) \frac{|\xi|^q}{(1+|\xi|^q)^2} |\widehat{U}(\xi)|^2 \, d\xi < 0. \tag{51}$$

To prove it, differentiate the explicit Fourier energy of the mass-preserving dilation at its scalar minimum $\lambda = 1$. The derivative is dominated by $C|\xi|^m$. Subtract the resulting identity divided by m from the energy. This uses no spatial moment.

5.3 Sharp behavior at the lower threshold

Return to the general symbol and assume additionally, uniformly on S^{N-1},

$$R^{-(m+q)} b(R\theta) \longrightarrow c_*(\theta) \qquad (R \downarrow 0), \tag{52}$$

where c_* is positive elliptic and homogeneous of degree $m + q$. Let

$$\mathcal{G}_0 = \{Q : M(Q) = M_0,\ A_0(Q) = 1,\ \|Q\|_p^p = p/2\}, \qquad C_*(f) = \int c_*(\xi) |\widehat{f}(\xi)|^2 \, d\xi.$$

Assume that at least one $Q \in \mathcal{G}_0$ has $C_*(Q) < \infty$. This finite-moment assumption is automatic for $0 < q \le m + 2$ under the stated regularity of a_0: its optimizer equation and the elliptic bootstrap give $Q \in H^{m+1}$. For larger q it is an explicit hypothesis; no unlimited smoothness of a non-polynomial nonlinearity is presumed.

Define

$$C_{\min} = \min_{Q \in \mathcal{G}_0} C_*(Q), \qquad \beta_* = \left[\frac{m}{(m+q) C_{\min}}\right]^{1/q}, \qquad K_* = \frac{q}{2(m+q)} \beta_*^m, \tag{53}$$
$$\mathcal{G}_{\mathrm{sel}} = \{Q \in \mathcal{G}_0 : C_*(Q) = C_{\min}\}.$$

These definitions are well posed. The assumed finite trial gives finiteness; positivity follows from Jensen's inequality,

$$\int |\xi|^{m+q} |\widehat{Q}|^2 \ge M_0^{-q/m} \left(\int |\xi|^m |\widehat{Q}|^2 \right)^{1+q/m},$$

and $A_0(Q) = 1$. A minimizing sequence in $\mathcal{G}_0$ is compact in H^s modulo translations, and the nonnegative C_* form is lower semicontinuous. Hence the minimum is attained. The selected set is compact in H^s modulo translations and constant phase.

Theorem 5.3 (Scale law and optimizer selection)**.** *Let $M \downarrow M_0$ and let U_M be any minimizer of $I(M)$. Set*

$$\alpha = \frac{M}{M_0}, \qquad d = \alpha^r - 1, \qquad \lambda_M = \left[\frac{A_0(U_M)}{\alpha}\right]^{1/m}, \qquad U_M(x) = \sqrt{\alpha}\, \lambda_M^{N/2} V_M(\lambda_M x). \tag{54}$$

Then $M(V_M) = M_0$, $A_0(V_M) = 1$, *and*

$$I(M) = -\alpha K_* d^{1+m/q}(1 + o(1)), \tag{55}$$

$$\lambda_M / d^{1/q} \longrightarrow \beta_*, \tag{56}$$

$$\operatorname{dist}_{H^s}(V_M, \mathcal{G}_{\mathrm{sel}}) \longrightarrow 0, \tag{57}$$

$$1 - \frac{2}{p}\|V_M\|_p^p = o(d), \tag{58}$$

$$\frac{B(U_M)}{A_0(U_M)} = \frac{m}{m+q} d + o(d). \tag{59}$$

The distance includes translations and constant phases, already contained in $\mathcal{G}_{\mathrm{sel}}$. *For the multiplier in* (47),

$$\omega_M = \frac{r}{M_0} \beta_*^m d^{m/q}(1 + o(1)). \tag{60}$$

Proof. For the normalization $M(V) = M_0$, $A_0(V) = 1$, let $H(V) = 1 - (2/p)\|V\|_p^p \geq 0$. The exact energy formula is

$$I(M) = \frac{\alpha \lambda_M^m}{2}[-d + (1+d)H(V_M)] + \frac{\alpha}{2} \int b(\lambda_M \xi) |\widehat{V}_M(\xi)|^2 \, d\xi. \tag{61}$$

Negativity implies $0 \leq H(V_M) < d/(1+d)$. Therefore normalized sharp-GN compactness applies, giving subsequential convergence, modulo translation and phase, to some $Q \in \mathcal{G}_0$ strongly in H^s.

First $\lambda_M \to 0$. Otherwise along a subsequence $\lambda_M \geq \ell > 0$, and (43) gives

$$\frac{B(U_M)}{A_0(U_M)} \geq c \int \min\{\lambda_M^q |\xi|^{m+q}, |\xi|^m\} |\widehat{V}_M(\xi)|^2 \, d\xi.$$

Choose a compact annulus carrying positive Fourier mass of the limiting Q. The right side is then bounded below by a fixed positive number. But (61) implies $B/A_0 < d \to 0$, a contradiction.

Define the rescaled correction form

$$D_\lambda(V) = \int \frac{b(\lambda \xi)}{\lambda^{m+q}} |\widehat{V}(\xi)|^2 \, d\xi.$$

Whenever $\lambda_n \downarrow 0$ and $V_n \to Q$ strongly in H^s,

$$\liminf_n D_{\lambda_n}(V_n) \geq C_*(Q). \tag{62}$$

Indeed on a compact annulus, (52) gives uniform multiplier convergence and strong Fourier L^2 convergence gives convergence of the integral. Discard the complement by nonnegativity, then exhaust all frequencies. In particular the liminf is at least $C_{\min}$ for the normalized minimizing profiles. This step does not assume uniform higher moments of those profiles.

For a fixed $Q \in \mathcal{G}_{\mathrm{sel}}$, (43) bounds the rescaled multiplier by $C|\xi|^{m+q}$. Dominated convergence gives $D_\lambda(Q) \to C_{\min}$. Testing with $\sqrt{\alpha}(\beta_* d^{1/q})^{N/2} Q(\beta_* d^{1/q} x)$ proves

$$\limsup_{M \downarrow M_0} \frac{I(M)}{\alpha d^{1+m/q}} \leq -K_* < 0. \tag{63}$$

On the other hand, negativity in (61) gives $\lambda_M^q D_{\lambda_M}(V_M) < d$. Equation (62) consequently yields $\lambda_M^q \leq 2d/C_{\min}$ for small d. The lower bound $I(M) \geq -\alpha d \lambda_M^m/2$, combined with (63), yields $\lambda_M^m \geq K_* d^{m/q}$ for small d. Thus

$$0 < c \leq \beta_M := \lambda_M / d^{1/q} \leq C < \infty. \tag{64}$$

The same inequalities bound $D_{\lambda_M}(V_M)$ and $H(V_M)/d$.

Extract a subsequence with $\beta_M \to \beta \in (0,\infty)$, $V_M \to Q \in \mathcal{G}_0$ strongly in H^s modulo symmetries, and convergent bounded scalar quantities $D_{\lambda_M}(V_M)$ and $H(V_M)/d$. Divide (61) by $\alpha d^{1+m/q}$:

$$\frac{I(M)}{\alpha d^{1+m/q}} = \frac{1}{2}\left[-\beta_M^m + (1+d)\beta_M^m \frac{H(V_M)}{d} + \beta_M^{m+q} D_{\lambda_M}(V_M)\right]. \tag{65}$$

Its liminf is at least

$$\tfrac{1}{2}[-\beta^m + C_*(Q)\beta^{m+q}] \geq \min_{t>0} \tfrac{1}{2}[-t^m + C_{\min} t^{m+q}] = -K_*.$$

Equality with (63) forces $\beta = \beta_*$, the unique scalar minimizer, and $C_*(Q) = C_{\min}$. It also forces $H(V_M)/d \to 0$ and $D_{\lambda_M}(V_M) \to C_{\min}$. Every extracted limit has these properties. This proves (55)–(58), including the distance statement. Finally,

$$\frac{B(U_M)}{A_0(U_M)} = \lambda_M^q D_{\lambda_M}(V_M) = d\beta_*^q C_{\min} + o(d) = \frac{m}{m+q} d + o(d).$$

The multiplier identity (47), together with $A_0(U_M) = \alpha\lambda_M^m$, $B/A_0 = O(d)$, and $I/A_0 = O(d)$, gives $\omega_M/\lambda_M^m \to r/M_0$, proving (60). □

The proof also gives the normalized frequency correction

$$\frac{\omega_M}{\lambda_M^m} = \frac{r}{M_0} + \frac{1}{M_0}\left(r + \frac{q}{m+q}\right) d + o(d).$$

This does not assert an additional-order expansion of λ_M.

5.4 The limiting functional

The selection has a precise variational interpretation. For $d > 0$ set $\alpha_d = (1+d)^{1/r}$ and $\ell = d^{1/q}$, and define, on H^s,

$$\mathcal{F}_d(W) = \frac{E_k(\sqrt{\alpha_d}\,\ell^{N/2} W(\ell\,\cdot))}{\alpha_d d^{1+m/q}} \quad \text{if } M(W) = M_0, \qquad \mathcal{F}_d(W) = +\infty \quad \text{otherwise.}$$

The exact expression on the mass constraint is

$$\mathcal{F}_d(W) = \frac{E_0(W)}{d} - \frac{1}{p}\|W\|_p^p + \frac{1}{2}\int \frac{b(\ell\xi)}{\ell^{m+q}} |\widehat{W}(\xi)|^2\, d\xi. \tag{66}$$

Proposition 5.4 (Variational selection on the equality manifold)**.** *In the strong H^s topology, $\mathcal{F}_d$ Γ-converges as $d \downarrow 0$ to*

$$\mathcal{F}_0(W) = \begin{cases} \tfrac{1}{2}[C_*(W) - A_0(W)], & M(W) = M_0,\ E_0(W) = 0, \\ +\infty, & \textit{otherwise.} \end{cases}$$

Its minimizers are exactly the dilates $\beta_*^{N/2} Q(\beta_*\,\cdot)$ with $Q \in \mathcal{G}_{\mathrm{sel}}$, including their symmetries.

Proof. On the fixed mass constraint $E_0 \geq 0$. If $W_n \to W$ strongly in H^s and $E_0(W) > 0$, continuity of the homogeneous energy and the nonlinear norm makes (66) tend to $+\infty$. If $E_0(W) = 0$, discard its nonnegative first term and apply the annular liminf argument (62). Since equality in the GN inequality gives $\|W\|_p^p/p = A_0(W)/2$, this is the required liminf. A limit of the fixed mass constraint has mass M_0 by strong L^2 convergence. For finite $\mathcal{F}_0(W)$, the constant sequence W is a recovery sequence, using domination by $C|\xi|^{m+q}|\widehat{W}|^2$. This proves the Γ-limit. Every nonzero equality profile is a dilation of an element of $\mathcal{G}_0$; minimizing its scalar dilation gives exactly (53). □

Translation invariance prevents global equicoercivity in H^s; none is asserted. Theorem 5.3 separately proves compactness of the rescaled minimizers modulo translations. The limiting principle says that the first nonzero low-frequency correction selects a subset of the entire homogeneous optimizer manifold.

5.5 A joint small-perturbation and mass-excess limit

There is a complementary selection limit that needs no higher-moment hypothesis. Fix a positive elliptic homogeneous symbol a as above and let

$$b_0(\xi) = a(\xi)\frac{|\xi|^q}{1+|\xi|^q}, \qquad k_\varepsilon = a + \varepsilon b_0, \qquad M_c = M_{c,a}.$$

Write $A = A_a$, $B_0(f) = \int b_0|\widehat{f}|^2$, and let I_ε be the constrained infimum for k_ε. The entire homogeneous equality manifold, with no kinetic normalization, is

$$\mathcal{H} = \{Q : M(Q) = M_c,\ E_a(Q) = 0\}.$$

For $0 < \tau < 1$ define

$$J(\tau) = \inf_{Q\in\mathcal{H}} \tfrac{1}{2}[B_0(Q) - \tau A(Q)], \qquad \mathcal{S}_\tau = \operatorname*{argmin}_{Q\in\mathcal{H}} \tfrac{1}{2}[B_0(Q) - \tau A(Q)].$$

Proposition 5.5 (Selection by an arbitrarily weak perturbation)**.** *For every $0 < \tau < 1$, $J(\tau)$ is finite, negative, and attained, and $\mathcal{S}_\tau$ is compact in H^s modulo translations and phase. Let $\varepsilon \downarrow 0$, set*

$$\alpha_\varepsilon = (1+\varepsilon\tau)^{1/r}, \qquad M_\varepsilon = \alpha_\varepsilon M_c,$$

and let U_ε be any minimizer of $I_\varepsilon(M_\varepsilon)$. Then

$$\frac{I_\varepsilon(M_\varepsilon)}{\alpha_\varepsilon\varepsilon} \longrightarrow J(\tau), \qquad \operatorname{dist}_{H^s}(U_\varepsilon/\sqrt{\alpha_\varepsilon}, \mathcal{S}_\tau) \longrightarrow 0.$$

No spatial rescaling is used. The same assertions hold along $\tau_\varepsilon \to \tau \in (0,1)$, with $\alpha_\varepsilon = (1+\varepsilon\tau_\varepsilon)^{1/r}$ and limiting set $\mathcal{S}_\tau$; the compactness estimates are uniform for τ_ε in a compact subinterval of $(0,1)$.

Proof. For every $\eta > 0$, $B_0(f) \geq (1-\eta)A(f) - C_\eta M(f)$. Thus the objective defining $J(\tau)$ is coercive in A on $\mathcal{H}$ if $\eta < 1 - \tau$. Low-frequency dilation of any homogeneous optimizer makes the objective negative. Its minimizing sequences consequently have A bounded above and bounded away from zero, the latter because the objective is at least $-\tau A/2$. Normalize A to one and apply sharp-GN compactness. The normalizing scales stay in a compact subset of $(0,\infty)$, giving strong H^s convergence modulo translations. Continuity of the order-m form B_0 proves attainment and compactness of the selected set.

On the mass constraint $M(f) = M_c$, the exact identity is

$$\frac{E_{k_\varepsilon}(\sqrt{\alpha_\varepsilon}f)}{\alpha_\varepsilon\varepsilon} = \frac{E_a(f)}{\varepsilon} + \frac{1}{2}B_0(f) - \frac{\tau}{p}\|f\|_p^p. \tag{67}$$

It Γ-converges in the strong H^s topology to $[B_0 - \tau A]/2$ on $\mathcal{H}$ and $+\infty$ off $\mathcal{H}$: use $E_a \geq 0$, continuity of the last two terms, and constant equality profiles as recovery sequences.

We prove compactness of actual minimizers separately. The sharp GN inequality at mass $\alpha_\varepsilon M_c$ gives

$$E_{k_\varepsilon}(U) \geq \frac{\varepsilon}{2}[(1-\eta-\tau)A(U) - C_\eta M(U)].$$

Testing (67) with a selected profile gives the exact bound $I_\varepsilon/(\alpha_\varepsilon\varepsilon) \leq J(\tau) < 0$. Hence $A(U_\varepsilon)$ is uniformly bounded. The other lower bound $E_{k_\varepsilon}(U) \geq -\varepsilon\tau A(U)/2$ forces it to be bounded away from zero. With $f_\varepsilon = U_\varepsilon/\sqrt{\alpha_\varepsilon}$, identity (67) now gives $E_a(f_\varepsilon) = O(\varepsilon)$. Normalize the kinetic form and use sharp-GN compactness to obtain a strong H^s limit $Q \in \mathcal{H}$ modulo translation, without any escaping spatial scale. The Γ-liminf is at least $J(\tau)$ and the trial upper bound is at most $J(\tau)$, proving convergence of the energies and selection of $Q \in \mathcal{S}_\tau$.

For $\tau_\varepsilon \to \tau$, the same proof has uniform constants: choose η below the distance of the compact parameter interval to one, and a single sufficiently low-frequency optimizer trial that is strictly negative for the smallest parameter in that interval. Continuity of the bounded-energy objective gives the stated limit. □

Along every convergent subsequence in this proposition, the multiplier satisfies $\omega_\varepsilon \to rA(Q)/M_c$ for $Q \in \mathcal{S}_\tau$. Thus a perturbation tending to zero can select a finite spatial scale and frequency from the critical homogeneous family when the mass excess has the same order as the perturbation. The selection functional J is concave and strictly decreasing. The first property follows by taking an infimum of affine functions; for the second, test the objective at $\tau_2 > \tau_1$ with a minimizer Q_1 at τ_1 to obtain

$$J(\tau_2) \le J(\tau_1) - \tfrac{1}{2}(\tau_2 - \tau_1)A(Q_1) < J(\tau_1).$$

5.6 A quartic example with explicit exponents

For $a_0(\xi) = |\xi|^4$ and $b(\xi) = \varepsilon|\xi|^6/(1+|\xi|^2)$, let

$$S_6 = \min_{Q\in\mathcal{G}_0} \int |\xi|^6 |\widehat{Q}(\xi)|^2 \, d\xi, \qquad d = (M/M_0)^{4/N} - 1, \qquad \alpha = M/M_0.$$

The higher-moment hypothesis holds, and $C_{\min} = \varepsilon S_6$. Theorem 5.3 gives

$$I(M) \sim -\alpha \frac{2}{27\varepsilon^2 S_6^2} d^3, \qquad \lambda_M^2 \sim \frac{2}{3\varepsilon S_6} d, \qquad \omega_M \sim \frac{4}{NM_0}\frac{4}{9\varepsilon^2 S_6^2} d^2.$$

Thus the spatial dilation length grows as $d^{-1/2}$, the principal kinetic energy vanishes as d^2, and the negative constrained energy vanishes as d^3. The selected critical profiles minimize their sixth Fourier moment. At the other end, $q = 2 < m = 4$ implies $I(M_\infty) = -\infty$.

For the isotropic family as a whole the result is a complete constrained energy classification: zero nonattained infimum up to M_0, negative attained infimum in (M_0, M_∞), and $-\infty$ above M_∞, with the upper endpoint determined by the three cases in Theorem 5.2. The new asymptotic concerns a class of nonhomogeneous equations; it does not resolve the unrestricted dynamics of the original homogeneous BNLS.

6 Ballistic exterior-gradient escape and weaker localization rigidity

We consider the pure mass-critical equation

$$iu_t = \Delta^2 u - |u|^{8/N} u, \qquad N \ge 1, \tag{68}$$

with the conventions

$$p = 2 + 8/N, \quad M = \|u\|_2^2, \quad P = \operatorname{Im} \int \overline{u}\nabla u, \quad E = \tfrac{1}{2}\|\Delta u\|_2^2 - \tfrac{1}{p}\|u\|_p^p.$$

Throughout this section, a solution is a standard strong H^2 solution for which these quantities are conserved and the localized virial identity holds in its integrated energy-space form. Suppose it is forward global and

$$\sup_{t\ge 0} \|u(t)\|_{H^2} \le B < \infty. \tag{69}$$

No radial symmetry or finite spatial moment is imposed. Our main result is a necessary spatial-spreading condition for a hypothetical bounded global negative-energy solution with zero momentum. It does not establish that such solutions exist, and it does not exclude all such solutions.

6.1 A tail estimate and a moving-center identity

For a center $c(t)$, write

$$D_R^c(t) = \int_{|x-c(t)|>R} |\nabla u(t,x)|^2 \, dx,$$
$$q_N = \min(4, 16/N), \quad \kappa_N = \min(1/2, 8/N), \quad \beta_N = \min(2, q_N).$$

The distinction between gradient localization and mass localization is useful: the former allows noncompact L^2 tails whose gradients vanish.

Lemma 6.1 (Gradient control of the nonlinear tail)**.** *For every center $c(t)$ and $R \geq 1$,*

$$\int_{|x-c(t)|>2R} |u(t,x)|^p \, dx \leq C_{N,M,B}\big(D_R^c(t)^{1/2} + R^{-1}\big)^{q_N}. \tag{70}$$

If $E < 0$, there is also $g_* > 0$, depending only on $N, M, B, |E|$, such that $\|\nabla u(t)\|_2 \geq g_*$ for all $t \geq 0$.

Proof. For $f \in H^2$, homogeneous Sobolev embedding and interpolation give

$$\|f\|_p^p \leq \begin{cases} C_N \|f\|_2^{2(4-N)/N} \|\nabla f\|_2^4, & N \leq 4, \\ C_N \|\nabla f\|_2^{16/N} \|\Delta f\|_2^{2-8/N}, & N > 4. \end{cases} \tag{71}$$

Indeed, the derivative order in the homogeneous embedding into L^p is $\sigma = N(1/2 - 1/p) = 2N/(N+4)$. Interpolate between orders zero and one when $\sigma \leq 1$, and between orders one and two when $\sigma > 1$. This includes the usual $\dot{H}^1 \hookrightarrow L^4$ endpoint at $N = 4$.

Choose a smooth cutoff η equal to zero on the unit ball and to one outside the ball of radius two. For $f(x) = \eta((x - c(t))/R)u(t,x)$, product differentiation gives

$$\|f\|_2 \leq \sqrt{M}, \quad \|\nabla f\|_2 \leq D_R^c(t)^{1/2} + C\sqrt{M}/R, \quad \|\Delta f\|_2 \leq C_B.$$

Substitution in (71) proves (70). Finally, negative energy implies $\|u(t)\|_p^p = p\|\Delta u(t)\|_2^2/2 + p|E| \geq p|E|$. The global inequalities (71), mass conservation, and (69) therefore give the positive gradient lower bound. □

Choose a smooth radial function ϕ_R which equals $|x|^2/2$ for $|x| \leq 4R$, is constant for $|x| \geq 8R$, and satisfies

$$D^2\phi_R \leq I, \qquad \|D^j \phi_R\|_\infty \leq C_j R^{2-j} \quad (1 \leq j \leq 6).$$

For a Lipschitz center a with $\|a'\|_\infty \leq L$, define

$$J_R(t) = 2\,\mathrm{Im} \int \nabla\phi_R(x - a(t)) \cdot \overline{u}(t,x) \nabla u(t,x) \, dx. \tag{72}$$

Lemma 6.2 (Moving virial with gradient-tail error)**.** *There are constants $C_0 = C_0(N, M, B) > 0$ and $C_1 = C_1(N, M, B, L) > 0$ such that*

$$|J_R(t)| \leq C_0 R, \tag{73}$$

$$J_R'(t) \leq 16E - 2a'(t) \cdot P + C_1\big(R^{-\beta_N} + D_R^a(t)^{\kappa_N}\big). \tag{74}$$

The derivative inequality is understood in the integrated sense.

Proof. The endpoint estimate follows directly from Cauchy–Schwarz. Put $j = \mathrm{Im}(\overline{u}\nabla u)$ and $\alpha = 8/N$. At a fixed center, differentiation and integration by parts give

$$\begin{aligned} J_R' = {} & 8\,\mathrm{Re} \int \partial_{jk}\phi_R \, \partial_{ij}\overline{u} \, \partial_{ik} u - 4\,\mathrm{Re} \int \partial_{jk}\Delta\phi_R \, \partial_j \overline{u} \, \partial_k u \\ & - 2\int \Delta^2\phi_R |\nabla u|^2 + \int \Delta^3 \phi_R |u|^2 - \frac{2\alpha}{p} \int \Delta\phi_R |u|^p. \end{aligned}$$

Repeated indices are summed, and the cutoff is evaluated at $x - a(t)$. Since $D^2\phi_R \leq I$, the leading term is at most $8\|D^2 u\|_2^2 = 8\|\Delta u\|_2^2$. The higher cutoff derivatives contribute $O_B(R^{-2})$. The nonlinear term equals $-16\|u\|_p^p/p$ up to an upper error bounded by $C\int_{|x-a(t)|>4R} |u|^p$. These principal terms sum to $16E$.

Motion of the center adds exactly

$$-2a'(t)\cdot\int D^2\phi_R(x-a(t))j(t,x)\,dx.$$

Replacing the Hessian by I gives $-2a'\cdot P$. Its error is at most

$$CL\int_{|x-a(t)|>4R}|u||\nabla u|\le CL\sqrt{M}\,D^a_{4R}(t)^{1/2}.$$

Apply Lemma 6.1, using $D^a_{2R},D^a_{4R}\le D^a_R$ and $0\le D^a_R\le B^2$. The nonlinear-tail exponent is $q_N/2=\min(2,8/N)$; the center error has exponent $1/2$. Reducing both to κ_N on this bounded interval proves (74). The calculation passes from smooth solutions to the integrated H^2 identity using the displayed bounds. A Lipschitz center has an almost-everywhere derivative; alternatively, smooth the center and pass to the integrated identity. □

6.2 A ballistic necessary alternative

Theorem 6.3 (Persistent exterior gradient at a linear distance)**.** *Suppose* (69) *holds,* $E=-e<0$*, and* $P=0$*. For every Lipschitz center* $a:[0,\infty)\to\mathbb{R}^N$ *with* $\|a'\|_\infty\le L$*, there are constants* $\nu,d>0$*, depending only on* N,M,B,e,L*, such that*

$$\liminf_{T\to\infty}\inf_{S\ge0}\frac{1}{T}\int_S^{S+T}\int_{|x-a(t)|>\nu T}|\nabla u(t,x)|^2\,dx\,dt\ge d. \tag{75}$$

In particular no choice of bounded-speed center can localize the gradient inside a sublinearly growing spatial radius in this averaged sense.

Proof. Integrate (74) on $[S,S+T]$, use $P=0$, and apply (73). Uniformly in S,

$$C_1\frac{1}{T}\int_S^{S+T}D^a_R(t)^{\kappa_N}\,dt\ge16e-C_1R^{-\beta_N}-2C_0R/T. \tag{76}$$

Choose a fixed $\nu>0$ so small that $2C_0\nu\le4e$, and put $R=\nu T$, held constant during each time integration. For all sufficiently large T, the right side is at least $8e$. Since $0<\kappa_N\le1$, Jensen's inequality gives

$$\left(\frac{1}{T}\int_S^{S+T}D^a_{\nu T}(t)\,dt\right)^{\kappa_N}\ge\frac{1}{T}\int_S^{S+T}D^a_{\nu T}(t)^{\kappa_N}\,dt\ge8e/C_1.$$

This proves (75) with $d=(8e/C_1)^{1/\kappa_N}$. The argument also proves the assertion uniformly over all centers with the specified speed bound. □

Remark 6.4. The theorem asserts persistent exterior gradient in a time-averaged sense, not at every time. Since $D^a_R(t)\le B^2$, the lower bound also forces a positive fraction of times in every sufficiently long window to have a nonzero exterior-gradient amount. The radius νT is proportional to the length of that window. No uniform positive ν or d is claimed as $E\to0$ or $B\to\infty$. The exterior gradient is not a physical dissipation, and the theorem supplies no finite budget for its repeated transport or replenishment.

6.3 Uniform and averaged gradient-localization rigidity

Theorem 6.5 (Averaged localization forces counter-momentum drift)**.** *Suppose* (69) *holds and* $E<0$*. Let* a *be Lipschitz, and suppose*

$$\lim_{R\to\infty}\limsup_{T\to\infty}\frac{1}{T}\int_0^T D^a_R(t)\,dt=0. \tag{77}$$

Then

$$\limsup_{T\to\infty} \frac{P\cdot[a(T)-a(0)]}{T} \le 8E. \tag{78}$$

Consequently (77) *is impossible when* $P = 0$.

Proof. Integrating (74) and rearranging gives

$$2\frac{P\cdot[a(T)-a(0)]}{T} \le 16E + C_1R^{-\beta_N} + C_1\left(\frac{1}{T}\int_0^T D_R^a(t)\,dt\right)^{\kappa_N} + 2C_0R/T.$$

Here Jensen was again used. First take the upper limit as $T\to\infty$ for fixed R, and then let $R\to\infty$. This proves the result. □

Lemma 6.6 (Regularizing a gradient concentration center)**.** *Suppose* (69) *holds,* $E<0$*, and an arbitrary map* $c:[0,\infty)\to\mathbb{R}^N$ *satisfies*

$$\sup_{t\ge0} D_R^c(t) \longrightarrow 0 \qquad (R\to\infty). \tag{79}$$

Then there is a Lipschitz map a *with* $\sup_{t\ge0}|a(t)-c(t)|<\infty$*. Gradient localization also holds about* a*.*

Proof. The nonlinearity maps bounded H^2 sets into bounded H^{-2} sets: for a test function $h\in H^2$, Hölder and $H^2\hookrightarrow L^p$ give

$$\left|\int |u|^{p-2}u\overline{h}\right| \le \|u\|_p^{p-1}\|h\|_p \le C_B\|h\|_{H^2}.$$

The equation therefore implies a uniform bound on u_t in H^{-2}. Interpolation between H^{-2} and the bounded H^2 norm yields

$$\|u(t)-u(s)\|_{H^1} \le C_B|t-s|^{1/4}. \tag{80}$$

Let $g_*>0$ be the gradient lower bound from Lemma 6.1. Choose R_0 so that $\sup_t D_{R_0}^c(t)\le g_*^2/16$, and choose $\tau>0$ so that (80) is at most $g_*/4$ when $|t-s|\le\tau$.

If $|c(t)-c(s)|>2R_0$, the two radius-R_0 balls are disjoint. On $B(c(s),R_0)$, the gradient of $u(s)$ has L^2 norm at least $\sqrt{15}g_*/4$, whereas that of $u(t)$ has norm at most $g_*/4$. The reverse triangle inequality contradicts (80). Thus $|c(t)-c(s)|\le 2R_0$ whenever $|t-s|\le\tau$.

Let a interpolate linearly through the values $c(n\tau)$, $n=0,1,\dots$. Then

$$\|a'\|_\infty \le 2R_0/\tau, \qquad \sup_{t\ge0}|a(t)-c(t)| \le 4R_0.$$

The bounded displacement preserves the uniform tail condition. □

Corollary 6.7 (Uniform gradient tightness, with no center regularity)**.** *Under* (69)*,* $E<0$*, and* (79)*,*

$$\limsup_{T\to\infty} \frac{P\cdot[c(T)-c(0)]}{T} \le 8E.$$

In particular there is no forward-global uniformly H^2*-bounded negative-energy zero-momentum solution with uniformly localized gradient modulo translations. This holds at every mass.*

Proof. Apply Lemma 6.6, followed by Theorem 6.5. The bounded difference between the two centers does not change the limiting drift quotient. □

Uniform mass tightness and (69) imply uniform gradient tightness by cutoff interpolation. The converse is false for general bounded H^2 families: a fixed localized core can be accompanied by a fixed-mass packet whose width tends to infinity and whose gradient norm tends to zero. The result above consequently weakens the required spatial localization, although it does not weaken the uniform-in-time H^2 bound.

6.4 The unresolved near-critical step

These results identify an actual requirement on any prospective bounded global counterexample, but they do not remove the general dynamical gap. If M_c is the sharp GN critical mass and $\delta_M = (M/M_c)^{4/N} - 1$, the sharp inequality gives

$$-\tfrac{1}{2}\delta_M A(u) \le E(u) < 0, \qquad A(u) = \|\Delta u\|_2^2.$$

For fixed negative energy, $A \ge 2|E|/\delta_M$. Conversely, normalizing $A = 1$ along a sequence $M \downarrow M_c$ forces $E \to 0$, so compactness can produce an allowed zero-energy optimizer. Normalizing $E = -1$ instead forces the kinetic norm to diverge. Near-optimizer compactness therefore does not, by itself, preserve a negative-energy contradiction at the limiting mass.

A near-critical orbit may have a small, nonzero exterior-gradient budget; $|E|/A$ can be arbitrarily smaller than that budget. The positive lower bounds in Theorem 6.3 can be compatible with this small tail. Closing the unrestricted problem requires a further mechanism that controls sustained radiation or its replenishment, or a construction showing that such behavior is possible. No such additional mechanism is asserted here.

7 Relation to established methods and remaining questions

Concentration compactness, translation profile decomposition, sharp interpolation, constrained minimization and localized virial identities are established tools. The claims requiring mathematical and historical assessment here are the particular threshold compactification, spectral bounds, uniform nonlocal gap, and dispersion-dependent selection statements in their precise stated classes.

Pausader–Shao [1] already use compact-orbit virial and center-motion arguments for the mass-critical fourth-order equation. Our gradient-tail and ballistic conclusions are consequences of that virial framework with the displayed localization estimates; we do not present the virial identity itself as new. The dynamical result allows noncompact mass tails, but it still assumes a uniform energy-space bound on the orbit.

Bonheure–Castéras–Gou–Jeanjean [2] analyze normalized solutions for mixed second- and fourth-order dispersion in a critical and supercritical regime. Liu–Romani–Su [3] study relations between optimizers and ground states in the mass-competition regime of that mixed-order equation. Those works supply relevant variational context. The same-order low/high frequency interpolation, its endpoint trichotomy and its optimizer-manifold selection laws are the specific objects analyzed here; a targeted comparison does not prove priority.

Cho–Hwang–Kwon–Lee [4], Theorems 1.1 and 1.2, prove negative-energy finite-time blowup for fractional/higher-order Hartree equations under radial, high-regularity and spatially weighted initial data assumptions. Their higher-order theorem has $2 < \alpha < 1 + N/2$. It is therefore not an unrestricted nonradial theorem for every symbol and interaction in Section 4. The Hartree result here is a traveling-profile exclusion theorem; no contradiction with that blowup theorem is asserted.

The central unfinished dynamical question in this continuation remains the existence or exclusion of a forward-global uniformly H^2-bounded pure BNLS orbit with $P = 0$, $E < 0$, and mass in a fixed sufficiently small interval above M_c. The new escape theorem identifies a necessary mechanism for such an orbit but does not make that mechanism impossible. Even excluding all global bounded orbits would not, by itself, exclude infinite-time growth of the H^2 norm or prove finite-time blowup.

For the static threshold, the next precise questions are whether the endpoint has negative or zero energy, whether more than one component can occur, and which spectral degeneracies permit the endpoint to be approached. For modified dispersion the full isotropic constrained classification is proved, while general anisotropic borderline upper-endpoint equality and nonlinear stability of individual branches require additional analysis. These are separate mathematical questions; none is settled merely by increasing the number of parameters covered.

References

[1] B. Pausader and S. Shao, *The mass-critical fourth-order Schrödinger equation in high dimensions*, J. Hyperbolic Differ. Equ. **7** (2010), 651–705, arXiv:0906.1547.

[2] D. Bonheure, J.-B. Castéras, T. Gou and L. Jeanjean, *Normalized solutions to the mixed dispersion nonlinear Schrödinger equation in the mass critical and supercritical regime*, arXiv:1802.09217v2.

[3] Z. Liu, G. Romani and Y. Su, *Mixed-dispersion Schrödinger equations and Gagliardo–Nirenberg inequalities: equivalence between ground states and optimizers*, arXiv:2603.26493v1 (2026).

[4] Y. Cho, G. Hwang, S. Kwon and S. Lee, *On finite time blowup for the mass-critical Hartree equations*, arXiv:1208.2302v3 (2014).

[5] Z. Lin and C. Zeng, *Instability, index theorem, and exponential trichotomy for linear Hamiltonian PDEs*, Mem. Amer. Math. Soc. **275** (2022), no. 1347, arXiv:1703.04016v4.