

PORTFOLIO

CHARAN TEJA REDDY A M

May 19 2026

General Problems in Mathematics

1. Problem in a Plane mirror Field-of-view solved by using complex numbers.

Consider a plane mirror of fixed length a . Suppose a point source O is placed in front of the mirror as shown in fig 1.1. It is also given that the ends of the mirror make angle α at O when subtended to meet. There is a straight screen which is placed to exactly fit the field of view of O w.r.t. the mirror. Furthermore, the ends of the screen when subtended to meet at the virtual image have equal lengths. Show that:

$$\text{Screen length} \propto \left| \sin \frac{\alpha}{2} \right|$$




Figure 1.1: Geometric configuration of the mirror and the point source.

2.

We know that in the Binomial theorem for positive integral powers,

$$(1+x)^n = {}^nC_0 + {}^nC_1x + \cdots + {}^nC_nx^n, \quad x \in \mathbb{R}, \quad n \in \mathbb{N}.$$

Similarly, let $W_n(x)$ be a function such that

$$W_n(x) = f^{(0)}(x) + {}^nC_1f^{(1)}(x) + {}^nC_2f^{(2)}(x) + \cdots + {}^nC_nf^{(n)}(x).$$

In a cleaner way,

$$W_n(x) = {}^nC_0 f^{(0)}(x) + {}^nC_1 f^{(1)}(x) + {}^nC_2 f^{(2)}(x) + \cdots + {}^nC_n f^{(n)}(x)$$

where $f(x)$ is an infinitely differentiable function over $x \in \mathbb{R}$ [i.e. differentiable of any order] and $f^{(r)}(x)$ denotes the r th derivative of $f(x)$; $f^{(0)}(x) = f(x)$.

For $W_n(x) = 0$, answer the following questions:

- (i) Solve for $f(x)$.
- (ii) What can be the degree of $f(x)$, if applicable?
- (iii) How many roots can $f(x)$ have at most, and at least?
- (iv) Suppose $S_i(x)$ is a function which is differentiable over $x \in \mathbb{R}$. Find the number of such $S_i(x)$ for which

$$\int_a^b f(x) dx + \int_{f(a)}^{f(b)} S_i(x) dx = b f(b) - a f(a); \quad a, b \in \mathbb{R}, \quad a \neq b.$$

Assume that $f(x)$ is fixed w.r.t. $S_i(x)$.

3. Basic

Consider the following statements. For two differentiable functions $f, g : \mathbb{R} \rightarrow \mathbb{R}$,

S_1 : There exists a c such that $c \in (a, b)$ and

$$\frac{g'(c)}{f'(c)} = \frac{g(b) - g(a)}{f(b) - f(a)}; \quad f(b) \neq f(a).$$

S_2 : There exists a $c \in (a, b)$ such that $g'(c) = \frac{g(a) - g(b)}{a - b}$.

S_3 : There exists a $c \in (a, b)$ such that $f'(c) = \frac{f(a) - f(b)}{a - b}$.

S_4 : If for $x \in [a, b]$, $f(a) = f(b)$ or $g(a) = g(b)$, then there exists a $c \in (a, b)$ such that $f'(c) = 0$ or $g'(c) = 0$ respectively.

[In all the above statements, $a, b, c \in \mathbb{R}$.]

Which of the following is/are correct?

- (A) $S_1 \Rightarrow S_2$ or $S_1 \Rightarrow S_3$
- (B) ' S_1 and S_2 ' $\Rightarrow S_4$
- (C) S_2 and $S_3 \Rightarrow S_1$
- (D) ' S_4 ' is derived from S_1, S_2, S_3 (any one of them).

4.

Let $I(x) = \int \frac{a(x)}{(b(x))^2} dx$, where $a(x)$ and $b(x)$ are real valued functions, $b(x) \neq 0$. Show that there exists a function $f(x)$, which is real valued, such that

$$\frac{f(x)}{I(x)} = b(x); \quad I(x) \neq 0,$$

if not, then disprove it.

5. Consider a function $f(x) = {}^nC_x$ defined for $n \neq 0$, $0 \leq x \leq n$; $x, n \in \mathbb{Z} - \mathbb{Z}^-$ where ${}^nC_r = \frac{n!}{(n-r)!r!}$. In the XY -plane, if $f(x)$ is plotted on it then find the area under the path traced by connecting its points with a single straight lines (that is, connecting every two consecutive points with a single straight line).