


Hardening the transient→sustained finding against pseudoreplication
tracked-unit collapse (BG_046) · session random-intercept (all mice) · per-session sign test



[A tracked-collapse BG_046] 24 responsive cell-sessions -> 23 unique consensus units (mean 1.0 sess/unit); classes: {'transient': 14, 'sustained': 6, 'intermediate': 3}
[change_on] RAW $p=3.30e-07$ | MIXED(session RE) $p=1.81e-07$ ($\beta=+0.98\text{Hz}$) | TRACKED $p=7.79e-01$ ($t=0.3978042679837923/s=0.7482093663911868$, n 14/6)
[hit_ramp] RAW $p=2.09e-07$ | MIXED(session RE) $p=1.54e-08$ ($\beta=+3.49\text{Hz}$) | TRACKED $p=6.22e-02$ ($t=1.5571051423324178/s=5.76953125$, n 14/6)
[fa_ramp] RAW $p=4.42e-11$ | MIXED(session RE) $p=3.35e-13$ ($\beta=+3.88\text{Hz}$) | TRACKED $p=5.07e-02$ ($t=1.72078043665506/s=5.620954239583665$, n 14/6)
[C per-session Δ] change_on: median $\Delta(\text{sus-tra})=+0.98\text{Hz}$ over 24 sessions, Wilcoxon $p=3.90e-03$
[C per-session Δ] hit_ramp: median $\Delta(\text{sus-tra})=+1.92\text{Hz}$ over 24 sessions, Wilcoxon $p=5.68e-04$
[C per-session Δ] fa_ramp: median $\Delta(\text{sus-tra})=+2.73\text{Hz}$ over 24 sessions, Wilcoxon $p=5.13e-06$