

# Computational Physics HW2

Siyuan Chen

September 20, 2023

P1: Numpy.float32 represents a real number with 32-digit of binary numbers. It approximates the number to be the form of  $\pm(1 + \sum_{i=0}^{22} m_i 2^{-(i+23)})2^{e-127}$  then represents it with the nearest mantissa  $m_i$  and exponent  $e$  from the format. In the 32-digit of binary numbers, the first digit on the left represents the positive or negative sign; the following 8 digit is the binary form of the exponent  $e$ ; the last 23 digit are the mantissa  $ms$ . In this method, 100.98763 is represented by 01000010110010011111100110101011. The round-off error of such representation is about  $2.7514648479609605 \times 10^{-6}$ .

P2: From the equation given in Exercise 2.9, we can cancel the physical constants on both sides and get  $M = \sum_{i,j,k=-L}^L (-1)^{i+j+k} \frac{1}{\sqrt{i^2+j^2+k^2}}$ , not  $i=j=k=0$ . In the first method, we write a for loop to add the potentials from atoms at different positions on the grid. In the second method, we write coordinates of the grid and calculate potentials as an array-like object. Notice here we have to use floating point data type to record coordinates such that certain calculations could be allowed. For  $L = 100$ , the time to execute for loop is about 40s whereas for grid method, it's only 0.25s. Both methods agree the Madelung constant for sodium in sodium chloride to be -1.7418.




Figure 1: Mandelbrot

P3: