

+++

date = '2025-03-31T22:24:23+08:00'

draft = true

title = 'Divisibility and number theory'

summary = "A comprehensive guide on divisibility rules, prime and composite numbers, GCD, and LCM, with manual number conversions using the Euclidean method for number system representation."

tags = ["Discrete Mathematics"]

[params]

math = true

+++

Number Theory (A.K.A. Higher Arithmetic)

A branch of mathematics concerned with the properties of integers.

Divisibility

Divisibility means dividing one number evenly by another, without leaving a remainder.



- $d \mid n$ means that d divides n
 - $d \nmid n$ means that d does not divide n
 - Conditions:
 - $d \neq 0$
 - $n \neq 0$
 - If $d \times q = n$, then $q = \frac{n}{d}$
-

Divisibility Rules

Rule for 2

A number is divisible by 2 if its **last digit** is even (i.e. 0, 2, 4, 6, or 8).


Examples:

- $2 \mid 128$ → 8 is even ⇒ 
 - $2 \nmid 129$ → 9 is odd ⇒ 
-

Rule for 3

A number is divisible by 3 if the **sum of its digits** is divisible by 3.

Examples:

- $3 \mid 381$:
 $3 + 8 + 1 = 12$ → $3 \mid 12$ ⇒ 

- $3 \nmid 217$:
 $2 + 1 + 7 = 10 \rightarrow 3 \nmid 10 \Rightarrow \text{X}$

Rule for 4

A number is divisible by 4 if the **last two digits** form a number divisible by 4.

Examples:

- $4 \mid 1312$:
 Last two digits: 12 $\rightarrow 4 \mid 12 \Rightarrow \checkmark$
- $4 \nmid 7019$:
 Last two digits: 19 $\rightarrow 4 \nmid 19 \Rightarrow \text{X}$

Rule for 5

A number is divisible by 5 if its **last digit** is either 0 or 5.

Examples:

- $5 \mid 205$ \rightarrow Last digit is 5 $\Rightarrow \checkmark$
- $5 \nmid 128$ \rightarrow Last digit is 8 $\Rightarrow \text{X}$

Rule for 6

A number is divisible by 6 if it is divisible by **both 2 and 3**.

Examples:

- $6 \mid 114$:
 - Divisible by 2: last digit 4 is even $\Rightarrow \checkmark$
 - Divisible by 3: $1 + 1 + 4 = 6 \rightarrow 3 \mid 6 \Rightarrow \checkmark$
- $6 \nmid 308$:
 - Divisible by 2: 8 is even $\Rightarrow \checkmark$
 - Not divisible by 3: $3 + 0 + 8 = 11 \Rightarrow \text{X}$

Rule for 7

Take the last digit, double it, and subtract it from the rest of the number. If the result is divisible by 7, then the number is divisible by 7.

Examples:

- $7 \mid 672$:
 $67 - 2 \times 2 = 63 \rightarrow 7 \mid 63 \Rightarrow \checkmark$
- $7 \nmid 905$:
 $90 - 2 \times 5 = 80 \rightarrow 7 \nmid 80 \Rightarrow \text{X}$

Rule for 8

A number is divisible by 8 if the **last three digits** form a number divisible by 8.

Examples:

- $8 \mid 109816$:
Last three digits: $816 \rightarrow 8 \mid 816 \Rightarrow \checkmark$
 - $8 \nmid 216302$:
Last three digits: $302 \rightarrow 8 \nmid 302 \Rightarrow \times$
-

Rule for 9

A number is divisible by 9 if the **sum of its digits** is divisible by 9.

Example:

- $9 \mid 1629$:
 $1 + 6 + 2 + 9 = 18 \rightarrow 9 \mid 18 \Rightarrow \checkmark$
-

Rule for 10

A number is divisible by 10 if it ends in **0**.

Examples:

- $10 \mid 130$ → ends in 0 $\Rightarrow \checkmark$
 - $10 \nmid 131$ → ends in 1 $\Rightarrow \times$
-

Rule for 11

A number is divisible by 11 if the **alternating sum of its digits** (i.e., subtracting every other digit) is divisible by 11 or equals 0.



Examples:

- $11 \mid 3729$:
 $(3 + 2) - (7 + 9) = 5 - 16 = -11 \Rightarrow 11 \mid 11 \Rightarrow \checkmark$
 - $11 \nmid 987$:
 $(9 + 7) - 8 = 16 - 8 = 8 \Rightarrow 11 \nmid 8 \Rightarrow \times$
-

Rule for 12

A number is divisible by 12 if it is divisible by **both 3 and 4**.

Examples:

- $12 \mid 648$:
 - $6 + 4 + 8 = 18 \Rightarrow 3 \mid 18$
 - Last two digits: $48 \Rightarrow 4 \mid 48$
 \Rightarrow 
- $12 \nmid 524$:
 - $5 + 2 + 4 = 11 \Rightarrow 3 \nmid 11$
 \Rightarrow 

Prime Numbers

These are **positive integers greater than 1** that are only divisible by **1 and themselves**.

Examples: 2, 3, 5, 7, 11, 13, ...

Composite Numbers

Composite numbers are **positive integers greater than 1** that are **not prime**.
 They have more than two distinct positive divisors.

Examples:

- $4 \rightarrow 2 \times 2$
 - $9 \rightarrow 3 \times 3$
 - $12 \rightarrow 2 \times 2 \times 3$
-

Greatest Common Divisor (GCD)

The **GCD** of two or more integers is the largest positive integer that divides each of the numbers.

- Denoted as $\gcd(a, b)$
- Can be found using **prime factorization** or the **Euclidean algorithm**

Example: GCD of 375 and 525

Prime factorization method:

- $375 = 3 \times 5^3$
- $525 = 3 \times 5^2 \times 7$

Common factors: $3 \times 5^2 = 75$
 $\Rightarrow \gcd(375, 525) = 75$

Least Common Multiple (LCM)

The **LCM** of two or more integers is the smallest number that is a multiple of all the given numbers.

- Denoted as $\text{lcm}(a, b)$

- Often found using prime factorization or the formula:
$$\text{lcm}(a, b) = \frac{a \times b}{\text{gcd}(a, b)}$$

Example:

LCM of 12 and 18

- $\text{gcd}(12, 18) = 6$
- $\text{lcm}(12, 18) = \frac{12 \times 18}{6} = 36$

Number Systems

Decimal (Base 10)

Digits: 0–9

Example: 123, 45

Binary (Base 2)

Digits: 0, 1

Prefix: 0b or 0x in some notations

Example: 0b1010 (10 in decimal)

Octal (Base 8)

Digits: 0–7

Prefix: 0o

Example: 0o17 (15 in decimal)

Hexadecimal (Base 16)

Digits: 0–9 and A–F (A=10, B=11, ..., F=15)

Prefix: 0x

Example: 0x1F (31 in decimal)

PROFI

Conversion Between Number Systems Using the Euclidean Method

The Euclidean algorithm is used for converting between different bases manually. Below is the step-by-step process of converting a decimal number to any base using the Euclidean method.

Example 1: Convert 156 to Binary (Base 2)

1. Divide 156 by 2:
 $156 \div 2 = 78$ remainder 0
2. Divide 78 by 2:
 $78 \div 2 = 39$ remainder 0

3. Divide 39 by 2:

$$39 \div 2 = 19 \text{ remainder } 1$$

4. Divide 19 by 2:

$$19 \div 2 = 9 \text{ remainder } 1$$

5. Divide 9 by 2:

$$9 \div 2 = 4 \text{ remainder } 1$$

6. Divide 4 by 2:

$$4 \div 2 = 2 \text{ remainder } 0$$

7. Divide 2 by 2:

$$2 \div 2 = 1 \text{ remainder } 0$$

8. Divide 1 by 2:

$$1 \div 2 = 0 \text{ remainder } 1$$

Now, read the remainders from bottom to top:

$$156_{10} = 10011100_2$$

Example 2: Convert 156 to Hexadecimal (Base 16)

1. Divide 156 by 16:

$$156 \div 16 = 9 \text{ remainder } 12 \text{ (12 in hexadecimal is C)}$$

2. Divide 9 by 16:

$$9 \div 16 = 0 \text{ remainder } 9$$

Now, read the remainders from bottom to top:

$$156_{10} = 9C_{16}$$

This method can be applied to convert between any number systems by following similar steps for division and taking remainders.
