

Track: AM

Low-Cost Biomodified Magnetite Nanoparticles for the Removal of Methylene Blue and Pb(II) ion from Simulated Wastewater

Chanchal Das^{1*}, Goutam Biswas¹ and Snehasis Bhakta²

¹Department of Chemistry, Cooch Behar Panchanan Barma University, Cooch Behar, India.

²Department of Chemistry, Coochbehar College, Cooch Behar, India.

Co-author, Email: chanchaldas453@gmail.com, snehasis.bhakta1989@gmail.com

Corresponding author, Email: goutam@cbpbu.ac.in

Industrialization and increasing population in this era making serious impact on aquatic life as well as making 'water stress' to the human. To enhance the resources for drinkable water there is need to develop efficient and cheaper purification technology so that anyone can afford it. Though current purification technology is now in much advanced level, high cost and energy is the preferable thinking to avoid the use of them. In this context, a cheaper and efficient adsorbent for inorganic and organic pollutant is framed (Das et al., 2021). It has magnetite nanoparticles in its core and the surface is modified by the natural products (polyphenols, steroids, flavonoids etc.) coming from the *Terminalia arjuna* (TA) bark extract (aqua). The synthetic procedure is simple greener and done under room temperature, via co-precipitation. Production cost for the nanoparticles (NPs) was \$19.48/kg per day. It has saturation magnetization value of 50 emu g⁻¹, having sizes 43 nm [from high resolution transmission electron microscopy (HRTEM) analysis] and had 43.2% coating (from thermogravimetric analysis). The batch adsorption technique revealed that, the Langmuir maximum adsorption capacity (Q_m) for the removal of methylene blue (MB) and Pb(II) (as organic and inorganic pollutant) onto *Terminalia arjuna* bark extract coated magnetite nanoparticles (TA@MNPs), was 294.1 and 210 mg g⁻¹ (**Table 1**). Adsorption process by the NPs obey pseudo-second-order kinetics and the physisorption behavior was also proven by the thermodynamic study. Desorption of the MB and Pb(II) by various eluents was also observed. Effect of various conditions was also applied to fine best adsorption result. From such low production cost and high efficiency to adsorb MB and Pb(II) ions, we can conclude that our nanoparticles might be used in industrial level to purify polluted surface water.

The results from the adsorption experiment can be summarized below:

- ✚ Highest adsorption occurred at pH 9, for MB and pH 3, for Pb(II).
- ✚ Maximum percentage of adsorption (equilibrium) achieved after 120 min (95%) for MB and after 514 min (61.2%) for Pb(II).
- ✚ Percentage of removal was increased with the increasing amount TA@MNPs for both the adsorbates.
- ✚ Percentage of adsorption was decreased with every 10 K rise of temperature.
- ✚ $\Delta G^\circ = -7.75$ to -9.74 kJ mol⁻¹ for MB and -0.3 to -1.62 kJ mol⁻¹ for Pb(II) adsorption, which lies in the range, $0 > \Delta G^\circ < -20$ kJ mol⁻¹, hence the process is physisorption (Húmpola et al., 2013).
- ✚ Our low-cost nanoparticles are advantageous over other, as it is reproducible, thermally stable over 400 °C and superparamagnetic (absence of hysteresis loop in the alternating gradient magnetometry study), hence can be separable after wastewater treatment by applying magnetic field.

A comparison for the maximum adsorption capacity (Q_m) of MB and Pb(II) on various magnetite nanoparticles (MNPs) based adsorbents.

	Adsorbents	Q_m (mg g ⁻¹)	References
For Pb(II)	TA@MNPs	210.5	Our study
	<i>Cnidium monnieri</i> (L.)Cuss-MNPs	105.6	(Lingamdinne et al., 2017)
	MDA-Fe ₃ O ₄	333.3	(Jiryaei Sharahi and Shahbazi, 2017)
	Fe ₃ O ₄ nanoparticles	36.0	(Nassar, 2010)
For MB	Fe ₃ O ₄ –multi wall carbon nanotubes	48.06	(Ai et al., 2011)
	Fe ₃ O ₄ @AC nanoparticles	138	(Joshi et al., 2019)
	Fe ₃ O ₄ nanoparticles	70.4	(Giri et al., 2011)
	TA@MNPs	294.1	Our study




Fig. a. Solid TA@MNPs, b. HRTEM image, c. FTIR spectra of i. TA and ii. TA@MNPs.

References

- Das, C., Singh, S., Bhakta, S., Mishra, P., Biswas, G., 2021. Bio-modified magnetic nanoparticles with Terminalia arjuna bark extract for the removal of methylene blue and lead (II) from simulated wastewater. *Chemosphere* 132673. <https://doi.org/10.1016/j.chemosphere.2021.132673>
- Húmpola, P.D., Odetti, H.S., Fertitta, A.E., Vicente, J.L., 2013. Thermodynamic analysis of adsorption models of phenol in liquid phase on different activated carbons. *J. Chil. Chem. Soc.* 58, 1541–1544. <https://doi.org/10.4067/S0717-97072013000100009>
- Ai, L., Zhang, C., Liao, F., Wang, Y., Li, M., Meng, L., Jiang, J., 2011. Removal of methylene blue from aqueous solution with magnetite loaded multi-wall carbon nanotube: Kinetic, isotherm and mechanism analysis. *J. Hazard. Mater.* 198, 282–290. <https://doi.org/10.1016/j.jhazmat.2011.10.041>
- Chang, J., Ma, J., Ma, Q., Zhang, D., Qiao, N., Hu, M., Ma, H., 2016. Adsorption of methylene blue onto Fe₃O₄/activated montmorillonite nanocomposite. *Appl. Clay Sci.* 119, 132–140. <https://doi.org/10.1016/j.clay.2015.06.038>
- Giri, S.K., Das, N.N., Pradhan, G.C., 2011. Synthesis and characterization of magnetite nanoparticles using waste iron ore tailings for adsorptive removal of dyes from aqueous solution. *Colloids Surf. Physicochem. Eng. Asp.* 389, 43–49. <https://doi.org/10.1016/j.colsurfa.2011.08.052>
- Jiryaei Sharahi, F., Shahbazi, A., 2017. Melamine-based dendrimer amine-modified magnetic nanoparticles as an efficient Pb(II) adsorbent for wastewater treatment: Adsorption optimization by response surface methodology. *Chemosphere* 189, 291–300. <https://doi.org/10.1016/j.chemosphere.2017.09.050>
- Joshi, S., Garg, V.K., Kataria, N., Kadirvelu, K., 2019. Applications of Fe₃O₄@AC nanoparticles for dye removal from simulated wastewater. *Chemosphere* 236, 124280. <https://doi.org/10.1016/j.chemosphere.2019.07.011>
- Lingamdinne, L.P., Chang, Y.-Y., Yang, J.-K., Singh, J., Choi, E.-H., Shiratani, M., Koduru, J.R., Attri, P., 2017. Biogenic reductive preparation of magnetic inverse spinel iron oxide nanoparticles for the adsorption removal of heavy metals. *Chem. Eng. J.* 307, 74–84. <https://doi.org/10.1016/j.cej.2016.08.067>
- Nassar, N.N., 2010. Rapid removal and recovery of Pb(II) from wastewater by magnetic nano-adsorbents. *J. Hazard. Mater.* 184, 538–546. <https://doi.org/10.1016/j.jhazmat.2010.08.069>