

Immunoinformatics Mediated Screening of Rotavirus A epitopes for Potential Peptide Vaccine Development

Pooja Rani Kuri¹ and Pranab Goswami^{1*}

¹Department of Biosciences and Bioengineering, Indian Institute of Technology Guwahati, Assam, India.

pooja176106028@iitg.ac.in

*Corresponding author, Email: pgoswami@iitg.ac.in

Rotavirus is a leading cause of severe infantile gastroenteritis worldwide, with about 199,000 childhood deaths in 2015, of which 90% cases occurred in low-income countries. India alone accounts for 22% of the global rotavirus gastroenteritis related deaths among children below 5 years of age. The World Health Organization recommends introduction of rotavirus vaccines as a priority in developing countries where high rates of rotavirus gastroenteritis are observed. There are however gaps in molecular data on rotavirus epidemiology in India. We therefore reviewed molecular data on rotavirus A (most common etiological agent of infantile gastroenteritis) isolated from India and also analyzed viral capsid protein genes (vp7, vp4 and vp6) to assess any particular trend with reference to different genotypes by phylogenetic analysis. Phylogenetic analysis revealed G1P[8] to be the most prevalently circulating strain of RVA in India. Therefore, capsid proteins VP4 P[8] and VP7 G[1] were subjected to antigenic analysis to determine epitopes for potential development of peptide vaccines. In this work, we identified highly antigenic, immunogenic, non-toxic and non-allergenic CTL and HTL epitopes of RVA G1P[8] capsid proteins VP4 and VP7. A total of 8 and 7 CTL epitopes were selected for RVA VP4 and VP7 respectively, while a total of 2 and 3 HTL epitopes were selected for RVA VP4 and VP7 respectively. The obtained epitopes were linked using suitable linkers followed by addition of a suitable adjuvant. The designed vaccine candidate was characterized using bioinformatics tools.

Results of the present work:

1. Molecular diversity of Indian Rotavirus A strains was determined by phylogenetic analysis and RVA G1P8 was found to be the most prevalently circulating strain.
2. Immunoinformatics approach was used to identify potential B and T cell epitopes using VP4 and VP7.
3. Selected epitopes were used to design a vaccine Sequence, followed by its computational physiochemical characterization.

References

- GBD 2015 Mortality and Causes of Death Collaborators. Global, regional, and national life expectancy, all-cause and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016. 34
- GBD Diarrhoeal Disease Collaborators. Estimates of global, regional, and national morbidity, mortality, and aetiologies of diarrhoeal diseases: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Infect. Dis*. 2017
- Jacqueline E. Tate et al. Global, Regional, and National Estimates of Rotavirus Mortality in Children <5 Years of Age, 2000–2013, *Clinical Infectious Diseases*. 2016.
- Estes, M. K. & Greenberg, H. B. in *Field's Virology*. 2013.
- Crawford et al. Rotavirus infection. *Disease Primers*. 2017.
- Jayaram et al. Emerging themes in rotavirus cell entry, genome organization, transcription and replication. *Virus Res*. 2004
- McClain et al. X-ray crystal structure of the rotavirus inner capsid particle at 3.8 Å resolution. *J. Mol. Biol.* 2010.
- Charpilienne et al. Identification of rotavirus VP6 residues located at the interface with VP2 that are essential for capsid assembly and transcriptase activity. *J. Virol.* 2002.
- Ulrich Desselberger. Rotaviruses. *Virus Res*. 2014.
- Matthijnsens et al. VP6-sequence-based cut off values as a criterion for rotavirus species demarcation. *Arch. Virol.* 2012.

Matthijnsens, J. et al. Uniformity of rotavirus strain nomenclature proposed by the Rotavirus Classification Working Group (RCWG). *Arch. Virol.* 2011.

Estes et al. Rotaviruses. In: Knipe, D.M., Howley, P.M., et al.(Eds.), *Fields Virology*. 2013.

Manual of Rotavirus detection and characterization methods. Immunization, Vaccination and Biologicals. World Health Organization. WHO/IVB/08.17




Figure 1 Molecular Diversity of Indian RVA strains. Genotypes Distribution of RVA VP4 (A) and RVA VP7 (B) determined by Phylogenetic Analysis.




Figure 2 A. Physiochemical characterization of the designed vaccine. B.Vaccine Secondary Structure Prediction by Phyre2.