
FULL MOLECULAR HAMILTONIAN

From Classical to Quantum

Chih-En Shen

Contents

1	Classical Cartesian Hamiltonian	4
2	Transformation to the Nuclear Center-of-Mass	4
2.1	New coordinates	4
2.2	Canonical momenta	4
2.3	Hamiltonian after the transformation	5
2.4	Eliminating the Cross-Coupling via a Classical Canonical Transformation . .	5
2.4.1	Generator	6
2.4.2	Induced canonical transformation	6
2.4.3	Canonicity check	7
2.4.4	Transformed Hamiltonian	7
2.4.5	Geometric interpretation	8
3	BF frame for both nuclei and electrons	9
3.1	Setup	9
3.2	From the internal Hamiltonian to a working Routhian	9
3.3	Body-fixed Routhian	10
3.4	Expanding (3.7)	10
3.4.1	Generic squared-term identity	10
3.4.2	Application to the nuclear term	11
3.4.3	Assembled Routhian	11
3.5	Canonical momenta and the total angular momentum	12
3.5.1	Nuclear angular momentum $\mathbf{J}_N = \partial\mathcal{R}/\partial\boldsymbol{\omega}$	12
3.5.2	Total angular momentum $\mathbf{J} = \mathbf{J}_N + \mathbf{L}_{\text{elec}}$	12
3.5.3	Vibrational momentum $P_b = \partial\mathcal{R}/\partial\dot{Q}_b$	12
3.6	Key identity	13
3.6.1	The clean split	13
3.6.2	Cross-check: the $\mathbf{I}_e/\mathbf{I}_X$ cancellation of the velocity-form route	13
3.6.3	Inversion: the Watson reciprocal inertia	14
3.7	Hamiltonian in canonical variables	14
3.7.1	Nuclear rotation–vibration block	14
3.7.2	Electronic block	15
3.7.3	Final assembly: (3.22)	15
3.8	Complete coupling enumeration	16
4	Quantization	17
4.1	Why curvilinear coordinates require care	17
4.2	Body-frame kinematics: the Euler-angle derivatives, explicitly	18
4.3	The ro-vibrational metric tensor	19
4.4	The electron sector: flat, constant, and decoupled from the measure	20
4.5	Determinant and the volume element	21
4.6	The inverse metric	21
4.7	The rotational derivative operators (defined, not postulated)	22
4.8	The Podolsky operator, block by block	23

4.9	Rescaling to the flat measure and the Watson pseudopotential	24
4.10	Electronic kinetic operator	25
4.11	Complete quantum molecular Hamiltonian	26
4.12	Born–Huang expansion	27
4.12.1	Clamped-nuclei electronic Hamiltonian	27
4.12.2	The exact Born–Huang expansion	27
4.12.3	Coupled-channel equations	28
4.12.4	Born–Oppenheimer approximation as truncation	28
4.12.5	Summary	28

1 Classical Cartesian Hamiltonian

We have N_{nuc} nuclei labeled by j with masses M_j , charges $Z_j e$, positions $\mathbf{R}_j$, and conjugate momenta $\mathbf{P}_j$; and N_{elec} electrons labeled by i with mass m_e , charge $-e$, positions $\mathbf{r}_i$, and momenta $\mathbf{p}_i$. All vectors live in a space-fixed (SF) lab frame. Using SI units and writing $R_{ij} = |\mathbf{R}_i - \mathbf{R}_j|$ etc.,

$$H = \underbrace{\sum_{j=1}^{N_{\text{nuc}}} \frac{\mathbf{P}_j^2}{2M_j}}_{T_N} + \underbrace{\sum_{i=1}^{N_{\text{elec}}} \frac{\mathbf{p}_i^2}{2m_e}}_{T_e} + V, \quad (1.1)$$

$$V = \frac{1}{4\pi\epsilon_0} \left(\frac{1}{2} \sum_{i \neq j} \frac{Z_i Z_j e^2}{R_{ij}} - \sum_{i,j} \frac{Z_j e^2}{|\mathbf{R}_j - \mathbf{r}_i|} + \frac{1}{2} \sum_{i \neq j} \frac{e^2}{r_{ij}} \right). \quad (1.2)$$

Canonical Poisson brackets $\{R_{j\alpha}, P_{k\beta}\} = \delta_{jk} \delta_{\alpha\beta}$, $\{r_{i\alpha}, p_{j\beta}\} = \delta_{ij} \delta_{\alpha\beta}$.

2 Transformation to the Nuclear Center-of-Mass

2.1 New coordinates

Define the nuclear COM, $3(N-1)$ internal nuclear vectors, and electron positions in the nuclear COM frame:

$$\mathbf{R}_{\text{cm}} = \frac{1}{M_N} \sum_j M_j \mathbf{R}_j, \quad M_N = \sum_j M_j, \quad (2.1)$$

$$\mathbf{R}'_j = \mathbf{R}_j - \mathbf{R}_{\text{cm}}, \quad \mathbf{r}'_i = \mathbf{r}_i - \mathbf{R}_{\text{cm}}. \quad (2.2)$$

The $\mathbf{R}'_j$ are linearly dependent:

$$\sum_j M_j \mathbf{R}'_j = 0. \quad (2.3)$$

2.2 Canonical momenta

This is a point transformation $\mathbf{Q}'_{\text{new}} = \mathbf{Q}'_{\text{new}}(\mathbf{q}_{\text{old}})$ with $\mathbf{q}_{\text{old}} = (\mathbf{R}_j, \mathbf{r}_i)$ and $\mathbf{Q}'_{\text{new}} = (\mathbf{R}_{\text{cm}}, \mathbf{R}'_j, \mathbf{r}'_i)$. Use the generating function: $F_2 = F_2(\mathbf{q}_{\text{old}}, \mathbf{P}, t) = \mathbf{P} \cdot \mathbf{Q}'_{\text{new}}(\mathbf{q}_{\text{old}})$

$$F_2 = \mathbf{P}_{\text{cm}} \cdot \mathbf{R}_{\text{cm}}(\{\mathbf{R}_j\}) + \sum_j \mathbf{P}'_j \cdot \mathbf{R}'_j(\{\mathbf{R}_j\}) + \sum_i \mathbf{p}'_i \cdot \mathbf{r}'_i(\mathbf{R}_j, \mathbf{r}_i). \quad (2.4)$$

Then $\mathbf{P}_j = \partial F_2 / \partial \mathbf{R}_j$, $\mathbf{p}_i = \partial F_2 / \partial \mathbf{r}_i$. Using

$$\frac{\partial \mathbf{R}_{\text{cm}}}{\partial \mathbf{R}_j} = \frac{M_j}{M_N} \mathbb{1}, \quad \frac{\partial \mathbf{R}'_k}{\partial \mathbf{R}_j} = (\delta_{jk} - \frac{M_j}{M_N}) \mathbb{1}, \quad \frac{\partial \mathbf{r}'_i}{\partial \mathbf{R}_j} = -\frac{M_j}{M_N} \mathbb{1}, \quad \frac{\partial \mathbf{r}'_i}{\partial \mathbf{r}_j} = \delta_{ij} \mathbb{1}, \quad (2.5)$$

one obtains

$$\mathbf{p}_i = \mathbf{p}'_i, \quad \mathbf{P}_j = \frac{M_j}{M_N} (\mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}'_i - \sum_k \mathbf{P}'_k) + \mathbf{P}'_j. \quad (2.6)$$

The constraint dual to (2.3) is $\sum_k \mathbf{P}'_k = 0$, so

$$\boxed{\mathbf{P}_j = \frac{M_j}{M_N}(\mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}_i) + \mathbf{P}'_j, \quad \mathbf{p}_i = \mathbf{p}'_i} \quad (2.7)$$

Total momentum: $\sum_j \mathbf{P}_j + \sum_i \mathbf{p}_i = \mathbf{P}_{\text{cm}}$. So $\mathbf{P}_{\text{cm}}$ is the *total* (nuclei+electrons) momentum.

2.3 Hamiltonian after the transformation

Insert (2.7) into T_N in (1.1):

$$T_N = \sum_j \frac{\mathbf{P}_j^2}{2M_j} = \sum_j \frac{1}{2M_j} \left[\frac{M_j}{M_N}(\mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}_i) + \mathbf{P}'_j \right]^2. \quad (2.8)$$

Expanding the square and summing,

$$T_N = \frac{(\mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}_i)^2}{2M_N} + \frac{1}{M_N}(\mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}_i) \cdot \underbrace{\sum_j \mathbf{P}'_j}_{=0} + \sum_j \frac{\mathbf{P}'_j^2}{2M_j}. \quad (2.9)$$

Combining with T_e and V :

$$H = \frac{\mathbf{P}_{\text{cm}}^2}{2M_N} - \underbrace{\frac{\mathbf{P}_{\text{cm}} \cdot \sum_i \mathbf{p}_i}{M_N}}_{\text{artifact}} + \underbrace{\frac{(\sum_i \mathbf{p}_i)^2}{2M_N}}_{\text{mass-polarization}} + \sum_j \frac{\mathbf{P}'_j^2}{2M_j} + \sum_i \frac{\mathbf{p}_i^2}{2m_e} + V(\mathbf{R}'_j, \mathbf{r}'_i). \quad (2.10)$$

Because H is independent of $\mathbf{R}_{\text{cm}}$, $\mathbf{P}_{\text{cm}}$ is conserved; pick the eigenvalue $\mathbf{P}_{\text{cm}} = 0$ (center-of-mass at rest) and drop the trivial $\mathbf{P}_{\text{cm}}$ -pieces:

$$\boxed{H_{\text{int}} = \sum_j \frac{\mathbf{P}'_j^2}{2M_j} + \sum_i \frac{\mathbf{p}_i^2}{2m_e} + \underbrace{\frac{1}{2M_N} \left(\sum_i \mathbf{p}_i \right)^2}_{\text{electron mass polarization}} + V(\mathbf{R}'_j, \mathbf{r}'_i).} \quad (2.11)$$

The potential is unchanged in form because $\mathbf{R}'_j - \mathbf{R}'_i = \mathbf{R}_j - \mathbf{R}_i$, $\mathbf{R}'_j - \mathbf{r}'_i = \mathbf{R}_j - \mathbf{r}_i$, $\mathbf{r}'_i - \mathbf{r}'_j = \mathbf{r}_i - \mathbf{r}_j$. The mass-polarization term, $(M_N)^{-1} \sum_{i < j} \mathbf{p}_i \cdot \mathbf{p}_j + (2M_N)^{-1} \sum_i \mathbf{p}_i^2$, is the price of referring electrons to the nuclear (rather than the total) COM. This rest-frame choice is not essential: Section 2.4 shows that the *same* internal Hamiltonian follows for any $\mathbf{P}_{\text{cm}}$ from a classical canonical transformation, and it is that frame-independent form we carry forward.

2.4 Eliminating the Cross-Coupling via a Classical Canonical Transformation

The rest-frame choice $\mathbf{P}_{\text{cm}} = 0$ made in Section 2.3 is convenient but stronger than necessary, and we now lift it. For $\mathbf{P}_{\text{cm}} \neq 0$ the Hamiltonian (2.10) carries two artifacts inherited from

having referred electrons to the *nuclear* (rather than the *total*) center of mass:

$$H = \underbrace{\frac{\mathbf{P}_{\text{cm}}^2}{2M_N}}_{(A) \text{ wrong total mass}} - \underbrace{\frac{\mathbf{P}_{\text{cm}} \cdot \sum_i \mathbf{p}'_i}{M_N}}_{(B) \text{ cross-coupling}} + \frac{(\sum_i \mathbf{p}'_i)^2}{2M_N} + \sum_j \frac{\mathbf{p}'_j{}^2}{2M_j} + \sum_i \frac{\mathbf{p}'_i{}^2}{2m_e} + V. \quad (2.12)$$

The wrong-mass term (A) has denominator M_N instead of the proper $M_{\text{tot}} = M_N + N_{\text{elec}}m_e$, and (B) couples the COM motion to the internal electronic momenta. We eliminate both *exactly, for any* $\mathbf{P}_{\text{cm}}$, by a single classical canonical transformation generated on phase space — so that the internal Hamiltonian carried into the body-fixed treatment of Section 3 is the frame-independent object, not a rest-frame special case. This transformation is the classical preimage of the Galilean-boost unitary that one would apply in the quantum theory.

2.4.1 Generator

Define on phase space the function

$$G \equiv \frac{m_e}{M_{\text{tot}}} \mathbf{P}_{\text{cm}} \cdot \sum_i \mathbf{r}'_i, \quad M_{\text{tot}} = M_N + N_{\text{elec}}m_e. \quad (2.13)$$

G depends only on $\mathbf{P}_{\text{cm}}$ and $\{\mathbf{r}'_i\}$; in particular it is independent of $\mathbf{R}_{\text{cm}}$, $\mathbf{R}'_j$, $\mathbf{P}'_j$, and $\mathbf{p}'_i$.

2.4.2 Induced canonical transformation

The infinitesimal canonical transformation generated by G acts on any phase-space function f by $\delta f = \{f, G\}$. Because G is linear in $\mathbf{P}_{\text{cm}}$ and $\{\mathbf{r}'_i\}$, the bracket $\{\cdot, G\}$ applied twice annihilates the canonical variables, so the finite transformation at unit parameter is given by a single application:

$$\tilde{f} = f + \{f, G\}. \quad (2.14)$$

Using $\{f, g\} = \sum (\partial_q f \partial_p g - \partial_p f \partial_q g)$, compute the action on each canonical pair.

Electron pair $(\mathbf{r}'_i, \mathbf{p}'_i)$:

$$\begin{aligned} \{r'_{i,a}, G\} &= \frac{\partial G}{\partial p'_{i,a}} = 0, \\ \{p'_{i,a}, G\} &= -\frac{\partial G}{\partial r'_{i,a}} = -\frac{m_e}{M_{\text{tot}}} P_{\text{cm},a}. \end{aligned}$$

Nuclear-COM pair $(\mathbf{R}_{\text{cm}}, \mathbf{P}_{\text{cm}})$:

$$\begin{aligned} \{R_{\text{cm},a}, G\} &= \frac{\partial G}{\partial P_{\text{cm},a}} = \frac{m_e}{M_{\text{tot}}} \sum_i r'_{i,a}, \\ \{P_{\text{cm},a}, G\} &= -\frac{\partial G}{\partial R_{\text{cm},a}} = 0. \end{aligned}$$

Internal nuclear pair $(\mathbf{R}'_j, \mathbf{P}'_j)$: G depends on neither, so $\{\mathbf{R}'_j, G\} = \{\mathbf{P}'_j, G\} = 0$. Hence the finite canonical transformation reads

$$\boxed{\begin{aligned} \tilde{\mathbf{r}}'_i &= \mathbf{r}'_i, & \tilde{\mathbf{p}}'_i &= \mathbf{p}'_i - \frac{m_e}{M_{\text{tot}}} \mathbf{P}_{\text{cm}}, \\ \tilde{\mathbf{R}}_{\text{cm}} &= \mathbf{R}_{\text{cm}} + \frac{m_e}{M_{\text{tot}}} \sum_i \mathbf{r}'_i, & \tilde{\mathbf{P}}_{\text{cm}} &= \mathbf{P}_{\text{cm}}, \\ \tilde{\mathbf{R}}'_j &= \mathbf{R}'_j, & \tilde{\mathbf{P}}'_j &= \mathbf{P}'_j. \end{aligned}} \quad (2.15)$$

2.4.3 Canonicity check

The only non-obvious Poisson bracket is between the two simultaneously shifted variables:

$$\begin{aligned}
\{\tilde{p}'_{i,a}, \tilde{R}_{\text{cm},b}\} &= \{\mathbf{p}'_{i,a} - \frac{m_e}{M_{\text{tot}}} P_{\text{cm},a}, R_{\text{cm},b} + \frac{m_e}{M_{\text{tot}}} \sum_l r'_{l,b}\} \\
&= \underbrace{\{p'_{i,a}, R_{\text{cm},b}\}}_0 + \frac{m_e}{M_{\text{tot}}} \sum_l \underbrace{\{p'_{i,a}, r'_{l,b}\}}_{-\delta_{il}\delta_{ab}} \\
&\quad - \frac{m_e}{M_{\text{tot}}} \underbrace{\{P_{\text{cm},a}, R_{\text{cm},b}\}}_{-\delta_{ab}} - \frac{m_e^2}{M_{\text{tot}}^2} \sum_l \underbrace{\{P_{\text{cm},a}, r'_{l,b}\}}_0 \\
&= -\frac{m_e}{M_{\text{tot}}} \delta_{ab} + \frac{m_e}{M_{\text{tot}}} \delta_{ab} = 0.
\end{aligned} \tag{2.16}$$

All other elementary brackets are preserved by inspection (each is either unchanged or trivially zero), so (2.15) is canonical.

2.4.4 Transformed Hamiltonian

Invert (2.15) for substitution into (2.12): $\mathbf{p}'_i = \tilde{\mathbf{p}}'_i + (m_e/M_{\text{tot}})\mathbf{P}_{\text{cm}}$. The potential V depends only on $\mathbf{R}'_j$ and $\mathbf{r}'_i$, which are unchanged, so V is form-invariant. Working term by term.

Cross-coupling (B):

$$\begin{aligned}
-\frac{\mathbf{P}_{\text{cm}} \cdot \sum_i \mathbf{p}'_i}{M_N} &= -\frac{\mathbf{P}_{\text{cm}} \cdot (\sum_i \tilde{\mathbf{p}}'_i + N_{\text{elec}} \frac{m_e}{M_{\text{tot}}} \mathbf{P}_{\text{cm}})}{M_N} \\
&= -\frac{\mathbf{P}_{\text{cm}} \cdot \sum_i \tilde{\mathbf{p}}'_i}{M_N} - \frac{N_{\text{elec}} m_e \mathbf{P}_{\text{cm}}^2}{M_N M_{\text{tot}}}.
\end{aligned} \tag{2.17}$$

Mass-polarization quadratic:

$$\begin{aligned}
\frac{(\sum_i \mathbf{p}'_i)^2}{2M_N} &= \frac{(\sum_i \tilde{\mathbf{p}}'_i + N_{\text{elec}} \frac{m_e}{M_{\text{tot}}} \mathbf{P}_{\text{cm}})^2}{2M_N} \\
&= \frac{(\sum_i \tilde{\mathbf{p}}'_i)^2}{2M_N} + \frac{N_{\text{elec}} m_e \mathbf{P}_{\text{cm}} \cdot \sum_i \tilde{\mathbf{p}}'_i}{M_N M_{\text{tot}}} + \frac{N_{\text{elec}}^2 m_e^2 \mathbf{P}_{\text{cm}}^2}{2M_N M_{\text{tot}}^2}.
\end{aligned} \tag{2.18}$$

Electron kinetic energy:

$$\begin{aligned}
\sum_i \frac{\mathbf{p}'_i{}^2}{2m_e} &= \sum_i \frac{(\tilde{\mathbf{p}}'_i + \frac{m_e}{M_{\text{tot}}} \mathbf{P}_{\text{cm}})^2}{2m_e} \\
&= \sum_i \frac{\tilde{\mathbf{p}}'_i{}^2}{2m_e} + \frac{\mathbf{P}_{\text{cm}} \cdot \sum_i \tilde{\mathbf{p}}'_i}{M_{\text{tot}}} + \frac{N_{\text{elec}} m_e \mathbf{P}_{\text{cm}}^2}{2M_{\text{tot}}^2}.
\end{aligned} \tag{2.19}$$

Coefficient of $\mathbf{P}_{\text{cm}} \cdot \sum_i \tilde{\mathbf{p}}'_i$: Summing (2.17), (2.18), (2.19),

$$-\frac{1}{M_N} + \frac{N_{\text{elec}} m_e}{M_N M_{\text{tot}}} + \frac{1}{M_{\text{tot}}} = \frac{-M_{\text{tot}} + N_{\text{elec}} m_e + M_N}{M_N M_{\text{tot}}} = 0, \tag{2.20}$$

using $M_{\text{tot}} = M_N + N_{\text{elec}} m_e$. *The cross-coupling (B) is killed identically.*

Coefficient of $\mathbf{P}_{\text{cm}}^2$: Adding the $\mathbf{P}_{\text{cm}}^2/(2M_N)$ from (A) plus the residual $\mathbf{P}_{\text{cm}}^2$ -pieces from

(2.17)–(2.19),

$$\begin{aligned}
\frac{1}{2M_N} - \frac{N_{\text{elec}}m_e}{M_N M_{\text{tot}}} + \frac{N_{\text{elec}}^2 m_e^2}{2M_N M_{\text{tot}}^2} + \frac{N_{\text{elec}}m_e}{2M_{\text{tot}}^2} &= \frac{1}{2M_N} - \frac{N_{\text{elec}}m_e}{M_N M_{\text{tot}}} + \frac{N_{\text{elec}}m_e}{2M_{\text{tot}}^2} \left(\frac{N_{\text{elec}}m_e}{M_N} + 1 \right) \\
&= \frac{1}{2M_N} - \frac{N_{\text{elec}}m_e}{M_N M_{\text{tot}}} + \frac{N_{\text{elec}}m_e}{2M_{\text{tot}}^2} \cdot \frac{M_{\text{tot}}}{M_N} \\
&= \frac{1}{2M_N} - \frac{N_{\text{elec}}m_e}{M_N M_{\text{tot}}} + \frac{N_{\text{elec}}m_e}{2M_N M_{\text{tot}}} \\
&= \frac{1}{2M_N} - \frac{N_{\text{elec}}m_e}{2M_N M_{\text{tot}}} = \frac{M_{\text{tot}} - N_{\text{elec}}m_e}{2M_N M_{\text{tot}}} = \frac{M_N}{2M_N M_{\text{tot}}} = \frac{1}{2M_{\text{tot}}}.
\end{aligned} \tag{2.21}$$

The wrong-mass (A) is corrected to M_{tot} .

Assembling everything,

$$\boxed{\tilde{H} = \frac{\mathbf{P}_{\text{cm}}^2}{2M_{\text{tot}}} + \sum_j \frac{\mathbf{P}_j'^2}{2M_j} + \sum_i \frac{\tilde{\mathbf{p}}_i'^2}{2m_e} + \frac{1}{2M_N} \left(\sum_i \tilde{\mathbf{p}}_i' \right)^2 + V(\mathbf{R}_j', \mathbf{r}_i').} \tag{2.22}$$

The COM kinetic energy carries the *correct* total mass M_{tot} . The internal piece is structurally identical to (2.11): the same mass-polarization $(\sum_i \tilde{\mathbf{p}}_i')^2/(2M_N)$, the same nuclear kinetic energy, the same potential. Since $\tilde{H}$ is independent of $\tilde{\mathbf{R}}_{\text{cm}}$, $\tilde{\mathbf{P}}_{\text{cm}} = \mathbf{P}_{\text{cm}}$ is conserved, and one may proceed with the internal dynamics for *any* value of $\mathbf{P}_{\text{cm}}$ — not just zero — with no kinetic artifact.

2.4.5 Geometric interpretation

The new “COM coordinate” is

$$\tilde{\mathbf{R}}_{\text{cm}} = \mathbf{R}_{\text{cm}} + \frac{m_e}{M_{\text{tot}}} \sum_i \mathbf{r}_i' = \frac{M_N \mathbf{R}_{\text{cm}} + m_e \sum_i \mathbf{r}_i}{M_{\text{tot}}} = \mathbf{R}_{\text{tot}}^{\text{CoM}}, \tag{2.23}$$

using $\mathbf{r}_i = \mathbf{R}_{\text{cm}} + \mathbf{r}_i'$. The canonical transformation has *quietly relabeled* the COM coordinate from nuclear to total COM, while *leaving the relative coordinates $\mathbf{R}_j'$ and $\mathbf{r}_i'$ untouched*. This is precisely what makes the framework still “nuclear-COM-based”: the Born–Oppenheimer-friendly relative coordinates of Section 2.1 survive unchanged. The momentum shift $\tilde{\mathbf{p}}_i' = \mathbf{p}_i' - (m_e/M_{\text{tot}})\mathbf{P}_{\text{cm}}$ is the kinetic compensation required to keep the Poisson brackets canonical: in Galilean terms, each electron’s lab-frame momentum is reduced by $m_e \mathbf{V}_{\text{CoM}}$ with $\mathbf{V}_{\text{CoM}} = \mathbf{P}_{\text{cm}}/M_{\text{tot}}$.

The canonical transformation has thus cleanly separated the total COM motion *exactly and for any $\mathbf{P}_{\text{cm}}$* : in (2.22) the COM appears only through the decoupled free term $\mathbf{P}_{\text{cm}}^2/2M_{\text{tot}}$ — carrying the *correct* total mass M_{tot} and free of the cross-coupling (B) — while every other term depends only on the internal variables. It is this total Hamiltonian (2.22), exact for *any* $\mathbf{P}_{\text{cm}}$ and not merely the rest frame, that we carry into the body-fixed treatment of Section 3.

3 BF frame for both nuclei and electrons

3.1 Setup

With Eckart's frame [1] fixed by the *nuclei*,

$$\mathbf{R}'_j = \mathbf{S}(\phi, \theta, \chi) \bar{\mathbf{R}}_j(Q), \quad \mathbf{r}'_i = \mathbf{S}(\phi, \theta, \chi) \bar{\mathbf{r}}_i. \quad (3.1)$$

The Eckart conditions $\sum_j M_j \bar{\mathbf{R}}_j = 0$ and $\sum_j M_j \bar{\mathbf{R}}_j^{\text{eq}} \times \bar{\mathbf{R}}_j = 0$ fix the orientation of $\mathbf{S}$ from the *nuclear* positions; the same $\mathbf{S}$ is then used to define BF electron coordinates $\bar{\mathbf{r}}_i$. The nuclear shape is parametrized by rectilinear mass-weighted **normal coordinates** Q_b ,

$$\bar{\mathbf{R}}_j(Q) = \bar{\mathbf{R}}_j^{\text{eq}} + \frac{1}{\sqrt{M_j}} \sum_b \mathbf{l}_{jb} Q_b, \quad (3.1a)$$

with mass-weighted normal-mode vectors $\mathbf{l}_{jb}$ obeying orthonormality and the Eckart (Sayvetz) conditions [1, 2]

$$\sum_j \mathbf{l}_{ja} \cdot \mathbf{l}_{jb} = \delta_{ab}, \quad \sum_j \sqrt{M_j} \mathbf{l}_{jb} = 0, \quad \sum_j \sqrt{M_j} \bar{\mathbf{R}}_j^{\text{eq}} \times \mathbf{l}_{jb} = 0, \quad (3.1b)$$

so that $\partial \bar{\mathbf{R}}_j / \partial Q_b = \mathbf{l}_{jb} / \sqrt{M_j}$ is constant.

BF angular velocity:

$$(\mathbf{S}^T \dot{\mathbf{S}})_{\alpha\beta} = -\epsilon_{\alpha\beta\gamma} \omega_\gamma \implies \mathbf{S}^T \dot{\mathbf{S}} \mathbf{v} = \boldsymbol{\omega} \times \mathbf{v}. \quad (3.2)$$

Velocities (using $|\mathbf{S}\mathbf{v}|^2 = |\mathbf{v}|^2$):

$$\dot{\mathbf{R}}'_j = \mathbf{S}(\boldsymbol{\omega} \times \bar{\mathbf{R}}_j + \dot{\bar{\mathbf{R}}}_j), \quad \dot{\mathbf{r}}'_i = \mathbf{S}(\boldsymbol{\omega} \times \bar{\mathbf{r}}_i + \dot{\bar{\mathbf{r}}}_i). \quad (3.3)$$

3.2 From the internal Hamiltonian to a working Routhian

I need an action principle for the body-fixed *nuclear* coordinates. The electrons, however, are best left in the canonical (momentum) form they already carry in (2.22): their kinetic energy and mass polarization form a rotational scalar that passes through the body-fixed transformation unchanged. I therefore Legendre-transform *only* the COM and nuclear momenta back to velocities — keeping $\mathbf{P}_{\text{cm}}$ arbitrary, never setting it to zero — and keep the electron momenta $\tilde{\mathbf{p}}'_i$ as canonical variables. The relevant Hamilton equations from (2.22) are

$$\dot{\mathbf{R}}_{\text{cm}} = \mathbf{P}_{\text{cm}} / M_{\text{tot}}, \quad \dot{\mathbf{R}}'_j = \mathbf{P}'_j / M_j, \quad \dot{\mathbf{r}}'_i = \tilde{\mathbf{p}}'_i / m_e + (\sum_l \tilde{\mathbf{p}}'_l) / M_N, \quad (3.4)$$

the last of which inverts to the electron momentum–velocity relation [recorded for the velocity-form cross-check below, around (3.18)]

$$\tilde{\mathbf{p}}'_i = m_e \dot{\mathbf{r}}'_i - \frac{m_e^2}{M_{\text{tot}}} \sum_l \dot{\mathbf{r}}'_l, \quad \sum_l \tilde{\mathbf{p}}'_l = \frac{m_e M_N}{M_{\text{tot}}} \sum_i \dot{\mathbf{r}}'_i. \quad (3.5)$$

Legendre-transforming the COM and nuclear momenta only, $\mathcal{R} = \mathbf{P}_{\text{cm}} \dot{\tilde{\mathbf{R}}}_{\text{cm}} + \sum_j \mathbf{P}'_j \dot{\tilde{\mathbf{R}}}'_j - \tilde{H}$, yields the **Routhian**

$$\boxed{\mathcal{R} = \frac{1}{2} M_{\text{tot}} |\dot{\tilde{\mathbf{R}}}_{\text{cm}}|^2 + \frac{1}{2} \sum_j M_j |\dot{\tilde{\mathbf{R}}}'_j|^2 - H_e(\tilde{\mathbf{p}}'_i) - V, \quad H_e \equiv \sum_i \frac{\tilde{\mathbf{p}}'^2_i}{2m_e} + \frac{1}{2M_N} \left(\sum_i \tilde{\mathbf{p}}'_i \right)^2,} \quad (3.6)$$

which is a Lagrangian in $(\tilde{\mathbf{R}}_{\text{cm}}, \mathbf{R}'_j)$ and *minus* a Hamiltonian in $(\mathbf{r}'_i, \tilde{\mathbf{p}}'_i)$: extremizing $\int \mathcal{R} dt$ gives Euler–Lagrange equations for the COM and nuclei and Hamilton’s equations for the electrons. The electronic kinetic energy and mass polarization thus retain their (2.22) momentum form H_e — they are never converted to velocity form. (The overall minus sign on H_e is the Routhian’s partial-Legendre signature; the mass polarization keeps its physical positive coefficient $+1/2M_N$ inside H_e , and the whole H_e re-enters the Hamiltonian with a plus sign when it is reassembled in (3.22).)

3.3 Body-fixed Routhian

Substitute the body-fixed velocities (3.3) into the nuclear term of (3.6); the COM term is untouched by the rotation. The electron Hamiltonian H_e is a *rotational scalar* — it is built from the dot products $\tilde{\mathbf{p}}'_i \cdot \tilde{\mathbf{p}}'_i$ — so under the passive rotation to the body frame it is unchanged in form. Defining the body-fixed electron momenta $\bar{\mathbf{p}}_i \equiv \mathbf{S}^T \tilde{\mathbf{p}}'_i$ (a proper rotation, so $|\bar{\mathbf{p}}_i|^2 = |\tilde{\mathbf{p}}'_i|^2$ and $\sum_i \bar{\mathbf{p}}_i = \mathbf{S}^T \sum_i \tilde{\mathbf{p}}'_i$, whence $(\sum_i \bar{\mathbf{p}}_i)^2 = (\sum_i \tilde{\mathbf{p}}'_i)^2$),

$$H_e = \sum_i \frac{\tilde{\mathbf{p}}'^2_i}{2m_e} + \frac{1}{2M_N} \left(\sum_i \tilde{\mathbf{p}}'_i \right)^2 = \sum_i \frac{\bar{\mathbf{p}}_i^2}{2m_e} + \frac{1}{2M_N} \left(\sum_i \bar{\mathbf{p}}_i \right)^2. \quad (3.6a)$$

Defining $\bar{\mathbf{r}} \equiv \sum_i \bar{\mathbf{r}}_i$, the body-fixed Routhian is therefore

$$\mathcal{R} = \frac{1}{2} M_{\text{tot}} |\dot{\tilde{\mathbf{R}}}_{\text{cm}}|^2 + \frac{1}{2} \sum_j M_j |\boldsymbol{\omega} \times \bar{\mathbf{R}}_j + \dot{\tilde{\mathbf{R}}}'_j|^2 - \sum_i \frac{\bar{\mathbf{p}}_i^2}{2m_e} - \frac{1}{2M_N} \left(\sum_i \bar{\mathbf{p}}_i \right)^2 - V, \quad (3.7)$$

in which only the nuclear kinetic term carries the body-fixed angular velocity $\boldsymbol{\omega}$. The potential is now manifestly **S**-independent:

$$V = \frac{1}{4\pi\epsilon_0} \left(\frac{1}{2} \sum_{j \neq k} \frac{Z_j Z_k e^2}{R_{jk}} - \sum_{j,i} \frac{Z_j e^2}{|\bar{\mathbf{R}}_j - \bar{\mathbf{r}}_i|} + \frac{1}{2} \sum_{i \neq l} \frac{e^2}{\bar{r}_{il}} \right). \quad (3.8)$$

Rotational symmetry is rendered explicit; the BF electronic problem at fixed Q is self-contained.

3.4 Expanding (3.7)

3.4.1 Generic squared-term identity

For any vectors $\mathbf{a}, \dot{\mathbf{a}}, \boldsymbol{\omega} \in \mathbb{R}^3$, the BCH-style identity $\mathbf{a} \times (\boldsymbol{\omega} \times \mathbf{a}) = \boldsymbol{\omega} a^2 - \mathbf{a}(\mathbf{a} \cdot \boldsymbol{\omega})$ gives

$$\begin{aligned} |\boldsymbol{\omega} \times \mathbf{a}|^2 &= (\boldsymbol{\omega} \times \mathbf{a}) \cdot (\boldsymbol{\omega} \times \mathbf{a}) = \boldsymbol{\omega} \cdot [\mathbf{a} \times (\boldsymbol{\omega} \times \mathbf{a})] \\ &= \boldsymbol{\omega} \cdot [\boldsymbol{\omega} a^2 - \mathbf{a}(\mathbf{a} \cdot \boldsymbol{\omega})] = \boldsymbol{\omega}^T (a^2 \mathbb{1} - \mathbf{a} \mathbf{a}^T) \boldsymbol{\omega}, \end{aligned} \quad (3.9a)$$

while the scalar triple product gives

$$(\boldsymbol{\omega} \times \mathbf{a}) \cdot \dot{\mathbf{a}} = \boldsymbol{\omega} \cdot (\mathbf{a} \times \dot{\mathbf{a}}). \quad (3.9b)$$

Hence, for any constant mass-like coefficient m ,

$$\begin{aligned} \frac{m}{2} |\boldsymbol{\omega} \times \mathbf{a} + \dot{\mathbf{a}}|^2 &= \frac{m}{2} [|\boldsymbol{\omega} \times \mathbf{a}|^2 + 2(\boldsymbol{\omega} \times \mathbf{a}) \cdot \dot{\mathbf{a}} + |\dot{\mathbf{a}}|^2] \\ &= \underbrace{\frac{m}{2} \boldsymbol{\omega}^T (a^2 \mathbb{1} - \mathbf{a} \mathbf{a}^T) \boldsymbol{\omega}}_{\text{rotational}} + \underbrace{m \boldsymbol{\omega} \cdot (\mathbf{a} \times \dot{\mathbf{a}})}_{\text{Coriolis-type}} + \underbrace{\frac{m}{2} |\dot{\mathbf{a}}|^2}_{\text{vibrational}}. \end{aligned} \quad (3.9c)$$

3.4.2 Application to the nuclear term

Only the nuclear kinetic term of the Routhian (3.7) carries $\boldsymbol{\omega}$. Apply (3.9c) with $m \rightarrow M_j$, $\mathbf{a} \rightarrow \bar{\mathbf{R}}_j$ and sum over j :

$$\frac{1}{2} \sum_j M_j |\boldsymbol{\omega} \times \bar{\mathbf{R}}_j + \dot{\bar{\mathbf{R}}}_j|^2 = \frac{1}{2} \boldsymbol{\omega}^T \mathbf{I}_N \boldsymbol{\omega} + \boldsymbol{\omega} \cdot \mathbf{L}_N^{\text{vib}} + T_{\text{vib}}^N. \quad (3.9d)$$

The electronic kinetic energy is *not* expanded this way — it is carried as the rotational scalar H_e of (3.6a). Only for the velocity-form cross-check of § 3.6 do we record the decomposition that the electronic and mass-polarization terms *would* produce if the electrons were instead kept in velocity form,

$$\frac{m_e}{2} \sum_i |\boldsymbol{\omega} \times \bar{\mathbf{r}}_i + \dot{\bar{\mathbf{r}}}_i|^2 = \frac{1}{2} \boldsymbol{\omega}^T \mathbf{I}_e \boldsymbol{\omega} + \boldsymbol{\omega} \cdot \mathbf{L}_e + T_e^{\text{BF}}, \quad -\frac{m_e^2}{2M_{\text{tot}}} |\boldsymbol{\omega} \times \bar{\mathbf{r}} + \dot{\bar{\mathbf{r}}}|^2 = -\frac{1}{2} \boldsymbol{\omega}^T \mathbf{I}_X \boldsymbol{\omega} - \boldsymbol{\omega} \cdot \mathbf{L}_X - T_X^{\text{mp}}. \quad (3.9e)$$

The constituent inertia tensors are

$$\mathbf{I}_N = \sum_j M_j (\bar{R}_j^2 \mathbb{1} - \bar{\mathbf{R}}_j \bar{\mathbf{R}}_j^T), \quad \mathbf{I}_e = m_e \sum_i (\bar{r}_i^2 \mathbb{1} - \bar{\mathbf{r}}_i \bar{\mathbf{r}}_i^T), \quad (3.9)$$

$$\mathbf{I}_X = \frac{m_e^2}{M_{\text{tot}}} (\bar{r}^2 \mathbb{1} - \bar{\mathbf{r}} \bar{\mathbf{r}}^T), \quad (3.10)$$

the angular-momentum-like (velocity-form) quantities are

$$\mathbf{L}_N^{\text{vib}} = \sum_j M_j \bar{\mathbf{R}}_j \times \dot{\bar{\mathbf{R}}}_j, \quad \mathbf{L}_e = m_e \sum_i \bar{\mathbf{r}}_i \times \dot{\bar{\mathbf{r}}}_i, \quad \mathbf{L}_X = \frac{m_e^2}{M_{\text{tot}}} \bar{\mathbf{r}} \times \dot{\bar{\mathbf{r}}}, \quad (3.11)$$

and the “vibrational” kinetic pieces are

$$T_{\text{vib}}^N = \frac{1}{2} \sum_j M_j |\dot{\bar{\mathbf{R}}}_j|^2 = \frac{1}{2} \sum_b \dot{Q}_b^2, \quad T_e^{\text{BF}} = \frac{m_e}{2} \sum_i |\dot{\bar{\mathbf{r}}}_i|^2, \quad T_X^{\text{mp}} = \frac{m_e^2}{2M_{\text{tot}}} |\dot{\bar{\mathbf{r}}}|^2. \quad (3.12)$$

3.4.3 Assembled Routhian

Inserting the nuclear expansion (3.9d) into (3.7), together with the decoupled COM term, the electron scalar H_e , and the potential:

$$\boxed{\mathcal{R} = \frac{1}{2} M_{\text{tot}} |\dot{\bar{\mathbf{R}}}_{\text{cm}}|^2 + \underbrace{\frac{1}{2} \boldsymbol{\omega}^T \mathbf{I}_N \boldsymbol{\omega} + \boldsymbol{\omega} \cdot \mathbf{L}_N^{\text{vib}} + T_{\text{vib}}^N}_{\text{nuclear, velocity form}} - H_e - V.} \quad (3.13)$$

The rotational kernel is the **nuclear** inertia $\mathbf{I}_N$ *alone*: because the electrons are carried as the scalar H_e rather than expanded in $\boldsymbol{\omega}$, no electronic or mass-polarization inertia ($\mathbf{I}_e, \mathbf{I}_X$) enters the kernel. The electrons couple to rotation only *kinematically*, through their angular momentum, which we identify next. The body-fixed electron momenta and the **electronic angular momentum** are

$$\bar{\mathbf{p}}_i = \mathbf{S}^T \tilde{\mathbf{p}}'_i, \quad \mathbf{L}_{\text{elec}} \equiv \sum_i \bar{\mathbf{r}}_i \times \bar{\mathbf{p}}_i. \quad (3.14)$$

Remark. The electronic and mass-polarization inertia/angular-momentum quantities $\mathbf{I}_e, \mathbf{I}_X, \mathbf{L}_e, \mathbf{L}_X$ of (3.9)–(3.11), and the velocity-form decomposition (3.9e), are needed *only* in the cross-check of §3.6; they play no role in the Routhian (3.13) itself.

3.5 Canonical momenta and the total angular momentum

Compute the conjugate momenta from the Routhian $\mathcal{R}$. The cyclic COM coordinate gives the conserved total momentum $\mathbf{P}_{\text{cm}} = \partial\mathcal{R}/\partial\dot{\mathbf{R}}_{\text{cm}} = M_{\text{tot}}\dot{\mathbf{R}}_{\text{cm}}$; being decoupled, it contributes nothing to the internal momenta below. The electron momenta $\bar{\mathbf{p}}_i$ are already canonical [eq. (3.14)]; it remains to find the rotational and vibrational momenta, most transparently by differentiating the unexpanded nuclear term of (3.7).

3.5.1 Nuclear angular momentum $\mathbf{J}_N = \partial\mathcal{R}/\partial\boldsymbol{\omega}$

In the Routhian (3.13) only the nuclear rotational kernel and the nuclear Coriolis piece carry $\boldsymbol{\omega}$ — the electron scalar H_e and the COM term do not. The momentum conjugate to the body-fixed angular velocity is therefore the *nuclear* angular momentum

$$\mathbf{J}_N \equiv \frac{\partial\mathcal{R}}{\partial\boldsymbol{\omega}} = \mathbf{I}_N \boldsymbol{\omega} + \mathbf{L}_N^{\text{vib}}. \quad (3.15)$$

3.5.2 Total angular momentum $\mathbf{J} = \mathbf{J}_N + \mathbf{L}_{\text{elec}}$

The body-fixed frame $\mathbf{S}(\Omega)$ rotates the nuclei *and* the electrons together [eq. (3.1)], so the total angular momentum conjugate to the Euler angles — the physical, conserved rovibronic angular momentum — is the sum of the nuclear and electronic contributions. The electronic part is the canonical electron angular momentum $\mathbf{L}_{\text{elec}} = \sum_i \bar{\mathbf{r}}_i \times \bar{\mathbf{p}}_i$ of (3.14), the body-fixed image of $\sum_i \mathbf{r}'_i \times \tilde{\mathbf{p}}'_i$ (the cross product co-rotates with $\mathbf{S}$). Hence

$$\mathbf{J} = \mathbf{J}_N + \mathbf{L}_{\text{elec}} = \mathbf{I}_N \boldsymbol{\omega} + \mathbf{L}_N^{\text{vib}} + \mathbf{L}_{\text{elec}}. \quad (3.16)$$

3.5.3 Vibrational momentum $P_b = \partial\mathcal{R}/\partial\dot{Q}_b$

Only $\dot{\mathbf{R}}_j$ carries $\dot{Q}_b$ dependence; with $\partial\bar{\mathbf{R}}_j/\partial Q_b = \mathbf{l}_{jb}/\sqrt{M_j}$ from (3.1a),

$$P_b \equiv \frac{\partial\mathcal{R}}{\partial\dot{Q}_b} = \sum_j M_j (\boldsymbol{\omega} \times \bar{\mathbf{R}}_j + \dot{\mathbf{R}}_j) \cdot \frac{\partial\bar{\mathbf{R}}_j}{\partial Q_b} = \sum_j \sqrt{M_j} (\boldsymbol{\omega} \times \bar{\mathbf{R}}_j + \dot{\mathbf{R}}_j) \cdot \mathbf{l}_{jb}. \quad (3.17)$$

Separating the rotational and vibrational pieces defines the **Coriolis vectors** σ_b ,

$$\sigma_b \equiv \sum_j M_j \bar{\mathbf{R}}_j \times \frac{\partial \bar{\mathbf{R}}_j}{\partial Q_b} = \sum_j \sqrt{M_j} \bar{\mathbf{R}}_j \times \mathbf{l}_{jb} = \sum_a \zeta_{ab} Q_a, \quad \zeta_{ab} \equiv \sum_j \mathbf{l}_{ja} \times \mathbf{l}_{jb}, \quad (3.17a)$$

the second form following from (3.1a) with the Eckart condition $\sum_j \sqrt{M_j} \bar{\mathbf{R}}_j^{\text{eq}} \times \mathbf{l}_{jb} = 0$. The vibrational metric is the identity, $\sum_j M_j (\partial \bar{\mathbf{R}}_j / \partial Q_a) \cdot (\partial \bar{\mathbf{R}}_j / \partial Q_b) = \sum_j \mathbf{l}_{ja} \cdot \mathbf{l}_{jb} = \delta_{ab}$, so the canonical vibrational momentum carries *no* Wilson G -matrix,

$$P_b = \boldsymbol{\omega} \cdot \sigma_b(Q) + \dot{Q}_b. \quad (3.17b)$$

3.6 Key identity

3.6.1 The clean split

The total-angular-momentum decomposition (3.16) *is* the key identity: subtracting the electronic part from both sides,

$$\boxed{\mathbf{J} - \mathbf{L}_{\text{elec}} = \mathbf{I}_N(Q) \boldsymbol{\omega} + \mathbf{L}_N^{\text{vib}}}. \quad (3.19)$$

The right side is a purely nuclear-velocity quantity, with the *nuclear* inertia $\mathbf{I}_N$ as rotational kernel. The entire electronic and mass-polarization content sits on the left, in the kinematic shift $\mathbf{J} \rightarrow \mathbf{J} - \mathbf{L}_{\text{elec}}$; it never enters the kernel, because in the Routhian (3.13) the electrons were carried as the scalar H_e and contributed no inertia at all.

3.6.2 Cross-check: the $\mathbf{I}_e/\mathbf{I}_X$ cancellation of the velocity-form route

Had we instead converted the electrons to velocity form — expanding their kinetic energy via (3.9e) — the electronic inertia and Coriolis terms would have entered *both* $\mathbf{J}$ and $\mathbf{L}_{\text{elec}}$, and (3.19) would emerge only after an exact cancellation. The velocity-form electron momentum is $\bar{\mathbf{p}}_i = m_e(\boldsymbol{\omega} \times \bar{\mathbf{r}}_i + \dot{\bar{\mathbf{r}}}_i) - \frac{m_e^2}{M_{\text{tot}}}(\boldsymbol{\omega} \times \bar{\mathbf{r}} + \dot{\bar{\mathbf{r}}})$ [the body-fixed image of (3.5)]; substituting into $\mathbf{L}_{\text{elec}} = \sum_i \bar{\mathbf{r}}_i \times \bar{\mathbf{p}}_i$ and using the BAC–CAB identity $\mathbf{a} \times (\boldsymbol{\omega} \times \mathbf{a}) = (a^2 \mathbf{1} - \mathbf{a} \mathbf{a}^T) \boldsymbol{\omega}$ on the two $\boldsymbol{\omega}$ -terms collapses the four pieces to

$$\mathbf{L}_{\text{elec}} = (\mathbf{I}_e - \mathbf{I}_X) \boldsymbol{\omega} + (\mathbf{L}_e - \mathbf{L}_X), \quad (3.18)$$

with $\mathbf{I}_e, \mathbf{I}_X, \mathbf{L}_e, \mathbf{L}_X$ from (3.9)–(3.11). The velocity-form total angular momentum would read $\mathbf{J} = (\mathbf{I}_N + \mathbf{I}_e - \mathbf{I}_X) \boldsymbol{\omega} + (\mathbf{L}_N^{\text{vib}} + \mathbf{L}_e - \mathbf{L}_X)$ [from (3.9d)+(3.9e)], so

$$\mathbf{J} - \mathbf{L}_{\text{elec}} = \mathbf{I}_N \boldsymbol{\omega} + \mathbf{L}_N^{\text{vib}} + \underbrace{[(\mathbf{I}_e - \mathbf{I}_X) \boldsymbol{\omega} + (\mathbf{L}_e - \mathbf{L}_X)] - \mathbf{L}_{\text{elec}}}_{= 0 \text{ by (3.18)}} = \mathbf{I}_N \boldsymbol{\omega} + \mathbf{L}_N^{\text{vib}},$$

reproducing (3.19): the electronic inertia $\mathbf{I}_e$ and the mass-polarization inertia $\mathbf{I}_X$ cancel *exactly*. This “algebraic miracle” of the Eckart construction is automatic in the mixed treatment, where $\mathbf{I}_e$ and $\mathbf{I}_X$ never arise.

3.6.3 Inversion: the Watson reciprocal inertia

To invert (3.19) for $\boldsymbol{\omega}$ we must re-express the velocity-form $\mathbf{L}_N^{\text{vib}}$ in canonical variables. In normal coordinates $\mathbf{L}_N^{\text{vib}} = \sum_j M_j \dot{\mathbf{R}}_j \times \dot{\mathbf{R}}_j = \sum_b \boldsymbol{\sigma}_b \dot{Q}_b$; eliminating $\dot{Q}_b = P_b - \boldsymbol{\omega} \cdot \boldsymbol{\sigma}_b$ from (3.17b),

$$\mathbf{L}_N^{\text{vib}} = \sum_b \boldsymbol{\sigma}_b (P_b - \boldsymbol{\omega} \cdot \boldsymbol{\sigma}_b) = \boldsymbol{\pi} - \boldsymbol{\sigma} \boldsymbol{\sigma}^T \boldsymbol{\omega}, \quad \boldsymbol{\pi} \equiv \sum_b \boldsymbol{\sigma}_b P_b, \quad \boldsymbol{\sigma} \boldsymbol{\sigma}^T \equiv \sum_b \boldsymbol{\sigma}_b \boldsymbol{\sigma}_b^T, \quad (3.20a)$$

with $\boldsymbol{\pi}$ the **field-free vibrational angular momentum**. Substituting into (3.19), $\mathbf{J} - \mathbf{L}_{\text{elec}} = (\mathbf{I}_N - \boldsymbol{\sigma} \boldsymbol{\sigma}^T) \boldsymbol{\omega} + \boldsymbol{\pi} \equiv \mathbf{I}' \boldsymbol{\omega} + \boldsymbol{\pi}$, so

$$\boldsymbol{\omega} = \boldsymbol{\mu} (\mathbf{J} - \mathbf{L}_{\text{elec}} - \boldsymbol{\pi}), \quad \boxed{\boldsymbol{\mu} \equiv (\mathbf{I}')^{-1} = (\mathbf{I}_N - \boldsymbol{\sigma} \boldsymbol{\sigma}^T)^{-1}}, \quad (3.20)$$

the **Watson reciprocal-inertia tensor** [3]. The rotation–vibration coupling is now carried by the modified inertia $\mathbf{I}'$ and the uniform shift $\mathbf{J} \rightarrow \mathbf{J} - \mathbf{L}_{\text{elec}} - \boldsymbol{\pi}$ — with no Wilson G -matrix.

3.7 Hamiltonian in canonical variables

We now assemble the body-fixed Hamiltonian in the canonical variables $(\Omega, \mathbf{J}; Q, P_b; \bar{\mathbf{r}}_i, \bar{\mathbf{p}}_i)$. The total energy splits as $\tilde{H} = \mathbf{P}_{\text{cm}}^2/2M_{\text{tot}} + H_{\text{int}}$, the COM piece being a decoupled spectator. The internal Hamiltonian is the sum of the nuclear rotation–vibration kinetic energy, the electron scalar, and the potential,

$$H_{\text{int}} = T_N^{\text{rot+vib}} + H_e + V, \quad T_N^{\text{rot+vib}} \equiv \frac{1}{2} \boldsymbol{\omega}^T \mathbf{I}_N \boldsymbol{\omega} + \boldsymbol{\omega} \cdot \mathbf{L}_N^{\text{vib}} + T_{\text{vib}}^N,$$

read off the Routhian (3.13). The electron scalar H_e [eq. (3.6a)] and the potential V are already canonical; only $T_N^{\text{rot+vib}}$ must be re-expressed through the momenta $(\mathbf{J}, P_b)$.

3.7.1 Nuclear rotation–vibration block

$T_N^{\text{rot+vib}}$ is homogeneous-quadratic in the nuclear velocities $(\boldsymbol{\omega}, \dot{Q}_b)$, so by Euler’s theorem $2T_N^{\text{rot+vib}} = \mathbf{J}_N \cdot \boldsymbol{\omega} + \sum_b P_b \dot{Q}_b$, with conjugate momenta $\mathbf{J}_N = \mathbf{J} - \mathbf{L}_{\text{elec}}$ [eqs. (3.15), (3.19)] and P_b [eq. (3.17b)]. Substituting $\dot{Q}_b = P_b - \boldsymbol{\omega} \cdot \boldsymbol{\sigma}_b$ and $\sum_b P_b \boldsymbol{\sigma}_b = \boldsymbol{\pi}$,

$$2T_N^{\text{rot+vib}} = (\mathbf{J} - \mathbf{L}_{\text{elec}}) \cdot \boldsymbol{\omega} + \sum_b P_b (P_b - \boldsymbol{\omega} \cdot \boldsymbol{\sigma}_b) = (\mathbf{J} - \mathbf{L}_{\text{elec}} - \boldsymbol{\pi}) \cdot \boldsymbol{\omega} + \sum_b P_b^2.$$

Inserting $\boldsymbol{\omega} = \boldsymbol{\mu} (\mathbf{J} - \mathbf{L}_{\text{elec}} - \boldsymbol{\pi})$ from (3.20),

$$\boxed{T_N^{\text{rot+vib}} \Big|_{\text{canonical}} = \frac{1}{2} (\mathbf{J} - \mathbf{L}_{\text{elec}} - \boldsymbol{\pi})^T \boldsymbol{\mu} (\mathbf{J} - \mathbf{L}_{\text{elec}} - \boldsymbol{\pi}) + \frac{1}{2} \sum_b P_b^2.} \quad (3.20d)$$

The vibrational kinetic energy is the *bare* $\frac{1}{2} \sum_b P_b^2$; the entire rotation–vibration coupling is carried by $\boldsymbol{\pi}$ and by the Watson inertia $\boldsymbol{\mu} = (\mathbf{I}_N - \boldsymbol{\sigma} \boldsymbol{\sigma}^T)^{-1}$ — no Wilson G -matrix survives.

3.7.2 Electronic block

The electronic kinetic energy and mass polarization require no further work: they are the rotational scalar H_e of (3.6a), carried unchanged from the Step-2 Hamiltonian (2.22),

$$H_e = \sum_i \frac{\bar{\mathbf{p}}_i^2}{2m_e} + \frac{1}{2M_N} \left(\sum_i \bar{\mathbf{p}}_i \right)^2. \quad (3.21)$$

Because H_e is built from the rotation-invariant dot products $\bar{\mathbf{p}}_i \cdot \bar{\mathbf{p}}_i$ and $\bar{\mathbf{p}}_i = \mathbf{S}^T \tilde{\mathbf{p}}'_i$ is a passive rotation, its body-fixed form is *identical* to the Step-2 form (2.22) under $\tilde{\mathbf{p}}'_i \rightarrow \bar{\mathbf{p}}_i$ — no Legendre transform or angular-momentum expansion is needed. The mass polarization survives the body-fixed transformation intact in functional form.

Reduced-mass check ($N_{\text{elec}} = 1$): $H_e = \bar{\mathbf{p}}_1^2/(2m_e) + \bar{\mathbf{p}}_1^2/(2M_N) = \bar{\mathbf{p}}_1^2/(2\mu_e)$ with $\mu_e = m_e M_N / (m_e + M_N) = m_e M_N / M_{\text{tot}}$. ✓

3.7.3 Final assembly: (3.22)

Add (3.20d) and (3.21), include $V = V_{NN}(Q) + V_{Ne}(Q, \{\bar{\mathbf{r}}_i\}) + V_{ee}(\{\bar{\mathbf{r}}_i\})$ from (3.8), and use $H_{\text{int}} = T_N^{\text{rot+vib}} + H_e + V$:

$$\begin{aligned} H_{\text{int}} = & \frac{1}{2} (\mathbf{J} - \mathbf{L}_{\text{elec}} - \boldsymbol{\pi})^T \boldsymbol{\mu}(Q) (\mathbf{J} - \mathbf{L}_{\text{elec}} - \boldsymbol{\pi}) \\ & + \frac{1}{2} \sum_b P_b^2 \\ & + \sum_i \frac{\bar{\mathbf{p}}_i^2}{2m_e} + \frac{1}{2M_N} \left(\sum_i \bar{\mathbf{p}}_i \right)^2 \\ & + V_{NN}(Q) + V_{Ne}(Q, \{\bar{\mathbf{r}}_i\}) + V_{ee}(\{\bar{\mathbf{r}}_i\}). \end{aligned} \quad (3.22)$$

Restoring the decoupled COM piece, the complete canonical Hamiltonian is $\tilde{H} = \mathbf{P}_{\text{cm}}^2 / 2M_{\text{tot}} + H_{\text{int}}$ with H_{int} given by (3.22) — exact for *any* $\mathbf{P}_{\text{cm}}$. The COM term is a conserved free-particle spectator that decouples from every body-fixed variable, so the quantization of Section 4 proceeds with H_{int} .

Here $\boldsymbol{\pi} = \sum_b \boldsymbol{\sigma}_b P_b$ is the field-free vibrational angular momentum, with the Coriolis vectors $\boldsymbol{\sigma}_b = \sum_a \boldsymbol{\zeta}_{ab} Q_a$ of (3.17a) and constants $\boldsymbol{\zeta}_{ab} = \sum_j \mathbf{l}_{ja} \times \mathbf{l}_{jb}$; the vibrational metric being the identity, no Wilson G -matrix enters.

$\mathbf{L}_{\text{elec}} = \sum_i \bar{\mathbf{r}}_i \times \bar{\mathbf{p}}_i$ is the BF canonical electronic angular momentum, and crucially:

- $\boldsymbol{\mu} = (\mathbf{I}_N - \boldsymbol{\sigma} \boldsymbol{\sigma}^T)^{-1}$ is the **Watson reciprocal-inertia tensor**, built from nuclear quantities alone: *no* electron or mass-polarization corrections enter the rotational kernel — they were absorbed into $\mathbf{L}_{\text{elec}}$ via (3.19).
- The electron contribution to rotation appears entirely as the Coriolis-like coupling $-\mathbf{J}^T \boldsymbol{\mu} \mathbf{L}_{\text{elec}}$.
- The mass polarization is preserved in its $\left(\sum_i \bar{\mathbf{p}}_i \right)^2 / (2M_N)$ form.

3.8 Complete coupling enumeration

Expanding the rotational kernel:

$$\frac{1}{2}(\mathbf{J} - \mathbf{L}_{\text{elec}} - \boldsymbol{\pi})^T \boldsymbol{\mu}(\mathbf{J} - \mathbf{L}_{\text{elec}} - \boldsymbol{\pi}) =$$

$$\underbrace{\frac{1}{2}\mathbf{J}^T \boldsymbol{\mu} \mathbf{J}}_{\text{rotation}} \underbrace{-\mathbf{J}^T \boldsymbol{\mu} \boldsymbol{\pi}}_{\text{rot-vib}} \underbrace{-\mathbf{J}^T \boldsymbol{\mu} \mathbf{L}_{\text{elec}}}_{\text{Coriolis rot-elec}} \underbrace{+\boldsymbol{\pi}^T \boldsymbol{\mu} \mathbf{L}_{\text{elec}}}_{\text{Coriolis vib-elec}} + \frac{1}{2}\boldsymbol{\pi}^T \boldsymbol{\mu} \boldsymbol{\pi} + \frac{1}{2}\mathbf{L}_{\text{elec}}^T \boldsymbol{\mu} \mathbf{L}_{\text{elec}}.$$

All couplings in (3.22), referred to the standard analyses [3, 4]:

1. **Pure rotation, anisotropic top:** $\frac{1}{2}\mathbf{J}^T \boldsymbol{\mu}(Q) \mathbf{J}$ — Q -dependent moments of inertia (rot-vib coupling through inertia).
2. **Rot-vib Coriolis** [5]: $-\mathbf{J}^T \boldsymbol{\mu} \boldsymbol{\pi}$.
3. **Rot-electron Coriolis:** $-\mathbf{J}^T \boldsymbol{\mu} \mathbf{L}_{\text{elec}}$ — the kinetic source of Λ -doubling [6, 7], Renner-Teller [8], and spin-rotation [9] analogues.
4. **Vib-electron angular coupling:** $\boldsymbol{\pi}^T \boldsymbol{\mu} \mathbf{L}_{\text{elec}}$ — exact non-adiabatic angular coupling intrinsic to BF embedding [10, 11, 12].
5. **Quadratic vibrational angular momentum:** $\frac{1}{2}\boldsymbol{\pi}^T \boldsymbol{\mu} \boldsymbol{\pi}$.
6. **Quadratic electronic angular momentum:** $\frac{1}{2}\mathbf{L}_{\text{elec}}^T \boldsymbol{\mu} \mathbf{L}_{\text{elec}}$.
7. **Pure vibrational kinetic:** $\frac{1}{2} \sum_b P_b^2$.
8. **Electronic kinetic:** $\sum_i \bar{\mathbf{p}}_i^2 / (2m_e)$.
9. **Mass-polarization, diagonal:** $\sum_i \bar{\mathbf{p}}_i^2 / (2M_N)$ — modifies effective electron mass [10, 4].
10. **Mass-polarization, off-diagonal (two-electron):** $\sum_{i < l} \bar{\mathbf{p}}_i \cdot \bar{\mathbf{p}}_l / M_N$.
11. **Coulomb:** $V_{NN}(Q) + V_{Ne}(Q, \{\bar{\mathbf{r}}_i\}) + V_{ee}(\{\bar{\mathbf{r}}_i\})$. All Euler-angle independent.

Compared to the previous (electrons-in-SF) framework: kinetic rot-electron coupling has *replaced* potential rot-electron coupling. V_{Ne} is now manifestly $\mathbf{S}$ -independent — the canonical chemistry advantage.

4 Quantization

The classical internal Hamiltonian (3.22) is parametrized by curvilinear coordinates (Euler angles ϕ, θ, χ and vibrational normal coordinates Q_b) and BF Cartesian electron coordinates $\bar{\mathbf{r}}_i$. Canonical quantization in curvilinear coordinates requires care: the naïve substitution $\pi_A \rightarrow -i\hbar \partial_A$ does not produce a Hermitian operator on the natural Hilbert space. The Podolsky (Laplace–Beltrami) prescription [13] resolves this, and we carry it out here *entirely in coordinate form*: we first construct the metric tensor of the ro-vibrational configuration space, then its determinant (which fixes the volume element) and its inverse, and then apply the Laplace–Beltrami operator block by block, writing out the explicit derivatives $\partial_\phi, \partial_\theta, \partial_\chi, \partial_{Q_b}$ at every stage. Hermiticity is automatic at each step because the Laplace–Beltrami operator is Hermitian by construction; in particular, *no commutator algebra of body-fixed angular momenta is ever used* (see the Remark in § 4.11). The residual ordering effects collapse into a single multiplicative function of the shape — the Watson pseudopotential. The term-by-term expansion of every block, with each derivative written out, is collected in Appendix C.

4.1 Why curvilinear coordinates require care

Label the curvilinear configuration-space coordinates

$$\{q^A\} = \{\phi, \theta, \chi, Q_1, \dots, Q_f\}, \quad f \equiv 3N_{\text{nucl}} - 6, \quad (4.1)$$

supplemented by the BF Cartesian electron coordinates $\bar{\mathbf{r}}_i$ (treated in § 4.4). In classical mechanics, any kinetic energy that is quadratic in the velocities defines a **metric tensor** $g_{AB}(q)$ on configuration space,

$$T = \frac{1}{2} g_{AB}(q) \dot{q}^A \dot{q}^B, \quad p_A \equiv \frac{\partial T}{\partial \dot{q}^A} = g_{AB} \dot{q}^B, \quad (4.2)$$

(repeated indices summed), and eliminating the velocities in favor of the momenta gives the Hamiltonian form

$$T = \frac{1}{2} g^{AB}(q) p_A p_B, \quad g^{AB} \equiv (g^{-1})_{AB}. \quad (4.3)$$

So three objects control everything: the metric g_{AB} , its inverse g^{AB} , and its determinant $g \equiv \det g_{AB}$. We will build all three explicitly for the coordinates (4.1) in § 4.2–4.6.

Why can we not simply replace $p_A \rightarrow -i\hbar \partial_A$ in (4.3)? Because on a curved configuration space the physically meaningful inner product carries the Riemannian volume weight, $\langle \varphi | \psi \rangle = \int \varphi^* \psi \sqrt{g} d^n q$, and $-i\hbar \partial_A$ is *not* Hermitian with respect to it. The one-dimensional case shows the problem completely: on $L^2(w(q) dq)$ with a positive weight w , integration by parts (boundary terms vanishing) gives

$$\int \varphi^* (-i\hbar \partial_q \psi) w dq = \int (-i\hbar \partial_q \varphi)^* \psi w dq - i\hbar \int \varphi^* \psi \frac{w'}{w} w dq, \quad (4.4)$$

so the adjoint is $(-i\hbar \partial_q)^\dagger = -i\hbar (\partial_q + w'/w) \neq -i\hbar \partial_q$ whenever the weight is not constant. A kinetic operator built by naïve substitution would fail to be Hermitian and could produce complex expectation values.

The resolution, due to Podolsky [13], is to replace the classical kinetic energy (4.3) by the **Laplace–Beltrami operator**:

$$\boxed{\hat{T} = -\frac{\hbar^2}{2} \frac{1}{\sqrt{g}} \partial_A (\sqrt{g} g^{AB} \partial_B), \quad g = \det g_{AB}.} \quad (4.5)$$

This operator is Hermitian on $L^2(\sqrt{g} d^n q)$ *by construction*: for wavefunctions with vanishing boundary terms (here: periodicity in ϕ, χ , the vanishing of $\sin \theta$ at the poles $\theta = 0, \pi$, and decay in Q_b), one integration by parts gives

$$\int \varphi^* \hat{T} \psi \sqrt{g} d^n q = \frac{\hbar^2}{2} \int (\partial_A \varphi)^* g^{AB} (\partial_B \psi) \sqrt{g} d^n q, \quad (4.6)$$

which is manifestly symmetric under $\varphi \leftrightarrow \psi$ (and manifestly positive for $\varphi = \psi$, since g^{AB} is positive definite wherever the classical kinetic energy is). When the metric is constant and Cartesian, $g_{AB} = \delta_{AB}$, (4.5) reduces to the familiar $-\frac{\hbar^2}{2} \nabla^2$ — the prescription contains ordinary quantum mechanics as a special case. Our entire task in this section is therefore: *compute g_{AB} , g , and g^{AB} for the coordinates (4.1), then expand (4.5).*

4.2 Body-frame kinematics: the Euler-angle derivatives, explicitly

Fix the body-fixed orientation by the *zyz* convention

$$\mathbf{S}(\phi, \theta, \chi) = R_z(\phi) R_y(\theta) R_z(\chi), \quad R_z(\psi) = \begin{pmatrix} \cos \psi & -\sin \psi & 0 \\ \sin \psi & \cos \psi & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad R_y(\vartheta) = \begin{pmatrix} \cos \vartheta & 0 & \sin \vartheta \\ 0 & 1 & 0 \\ -\sin \vartheta & 0 & \cos \vartheta \end{pmatrix}, \quad (4.7)$$

so that $\mathbf{R}'_j = \mathbf{S} \bar{\mathbf{R}}_j$ as in (3.1). The body-fixed angular velocity is defined by $(\mathbf{S}^T \dot{\mathbf{S}})_{\alpha\beta} = -\epsilon_{\alpha\beta\gamma} \omega_\gamma$ [eq. (3.2)]; our first job is to express $\boldsymbol{\omega}$ through $(\dot{\phi}, \dot{\theta}, \dot{\chi})$. Differentiating the one-parameter rotations at zero angle defines the generators

$$G_z \equiv \partial_\psi R_z|_0 = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad G_y \equiv \partial_\vartheta R_y|_0 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad (4.8)$$

with $\partial_\psi R_z(\psi) = G_z R_z(\psi)$ and $\partial_\vartheta R_y(\vartheta) = G_y R_y(\vartheta)$ (check by direct differentiation of (4.7)). The chain rule applied to (4.7) then gives $\dot{\mathbf{S}} = \dot{\phi} G_z \mathbf{S} + \dot{\theta} R_z(\phi) G_y R_y(\theta) R_z(\chi) + \dot{\chi} \mathbf{S} G_z$, and multiplying by $\mathbf{S}^T$ from the left,

$$\mathbf{S}^T \dot{\mathbf{S}} = \dot{\phi} \mathbf{S}^T G_z \mathbf{S} + \dot{\theta} R_z(\chi)^T G_y R_z(\chi) + \dot{\chi} G_z = [\dot{\phi} \mathbf{S}^T \hat{\mathbf{z}} + \dot{\theta} R_z(\chi)^T \hat{\mathbf{y}} + \dot{\chi} \hat{\mathbf{z}}]_\times, \quad (4.9)$$

where $[\mathbf{a}]_\times$ denotes the antisymmetric matrix with $([\mathbf{a}]_\times)_{\alpha\beta} = -\epsilon_{\alpha\beta\gamma} a_\gamma$, and we used the conjugation rule $R^T [\mathbf{a}]_\times R = [R^T \mathbf{a}]_\times$ for any rotation R , together with $R_y(\theta)^T \hat{\mathbf{y}} = \hat{\mathbf{y}}$. Evaluating the two nontrivial vectors from (4.7), $\mathbf{S}^T \hat{\mathbf{z}} = (-\sin \theta \cos \chi, \sin \theta \sin \chi, \cos \theta)^T$ and $R_z(\chi)^T \hat{\mathbf{y}} = (\sin \chi, \cos \chi, 0)^T$, and comparing with the definition of $\boldsymbol{\omega}$, we obtain $\boldsymbol{\omega} = \mathbf{E}(\Omega) \dot{\boldsymbol{\Omega}}$

with $\dot{\mathbf{\Omega}} = (\dot{\phi}, \dot{\theta}, \dot{\chi})^T$ and the **kinematic matrix**

$$\mathbf{E}(\Omega) = \begin{pmatrix} -\sin \theta \cos \chi & \sin \chi & 0 \\ \sin \theta \sin \chi & \cos \chi & 0 \\ \cos \theta & 0 & 1 \end{pmatrix}, \quad \begin{aligned} \omega_1 &= -\dot{\phi} \sin \theta \cos \chi + \dot{\theta} \sin \chi, \\ \omega_2 &= \dot{\phi} \sin \theta \sin \chi + \dot{\theta} \cos \chi, \\ \omega_3 &= \dot{\phi} \cos \theta + \dot{\chi}. \end{aligned} \quad (4.10)$$

Expanding the determinant along the third column,

$$\det \mathbf{E} = 1 \cdot \det \begin{pmatrix} -\sin \theta \cos \chi & \sin \chi \\ \sin \theta \sin \chi & \cos \chi \end{pmatrix} = -\sin \theta (\cos^2 \chi + \sin^2 \chi) = -\sin \theta, \quad |\det \mathbf{E}| = \sin \theta. \quad (4.11)$$

Solving the linear system (4.10) for the Euler rates — multiply the ω_1, ω_2 lines by $(-\cos \chi, \sin \chi)$ and by $(\sin \chi, \cos \chi)$ respectively and add, then substitute into the ω_3 line — gives $\dot{\mathbf{\Omega}} = \mathbf{E}^{-1} \boldsymbol{\omega}$ with

$$\mathbf{E}^{-1}(\Omega) = \begin{pmatrix} -\frac{\cos \chi}{\sin \theta} & \frac{\sin \chi}{\sin \theta} & 0 \\ \sin \chi & \cos \chi & 0 \\ \frac{\cos \theta \cos \chi}{\sin \theta} & -\frac{\cos \theta \sin \chi}{\sin \theta} & 1 \end{pmatrix}, \quad \begin{aligned} \dot{\phi} &= \frac{1}{\sin \theta} (-\cos \chi \omega_1 + \sin \chi \omega_2), \\ \dot{\theta} &= \sin \chi \omega_1 + \cos \chi \omega_2, \\ \dot{\chi} &= \omega_3 + \frac{\cos \theta}{\sin \theta} (\cos \chi \omega_1 - \sin \chi \omega_2). \end{aligned} \quad (4.12)$$

The $1/\sin \theta$ entries reflect the coordinate singularity of the Euler parametrization at the poles $\theta = 0, \pi$ (where ϕ and χ describe the same rotation); it is compensated by the vanishing of $|\det \mathbf{E}| = \sin \theta$ in the volume element, and no physical quantity is singular there. Equations (4.10)–(4.12) are the only facts about the Euler angles used in the rest of this section.

4.3 The ro-vibrational metric tensor

The classical rotation–vibration kinetic energy in the body frame is the nuclear term of the Routhian (3.13),

$$2T_{\text{rv}} = \boldsymbol{\omega}^T \mathbf{I}_N(Q) \boldsymbol{\omega} + 2 \boldsymbol{\omega}^T \sum_b \boldsymbol{\sigma}_b(Q) \dot{Q}_b + \sum_b \dot{Q}_b^2, \quad (4.13)$$

whose three pieces were derived in §3.4: the nuclear inertia tensor $\mathbf{I}_N = \sum_j M_j (\bar{R}_j^2 \mathbb{1} - \bar{\mathbf{R}}_j \bar{\mathbf{R}}_j^T)$ [(3.9)], the Coriolis vectors $\boldsymbol{\sigma}_b = \sum_a \boldsymbol{\zeta}_{ab} Q_a$ with *constant* coefficients $\boldsymbol{\zeta}_{ab} = \sum_j \mathbf{l}_{ja} \times \mathbf{l}_{jb}$ [(3.17a)], and the identity vibrational metric $\sum_j M_j (\partial \bar{\mathbf{R}}_j / \partial Q_a) \cdot (\partial \bar{\mathbf{R}}_j / \partial Q_b) = \delta_{ab}$, a consequence of the mass-weighted rectilinear normal coordinates [(3.17)]. Collect the Coriolis vectors into the $3 \times f$ matrix $\sigma_{ab} \equiv (\boldsymbol{\sigma}_b)_a$ and write (4.13) as one quadratic form in the *physical* velocities,

$$2T_{\text{rv}} = \begin{pmatrix} \boldsymbol{\omega} \\ \dot{\mathbf{Q}} \end{pmatrix}^T \begin{pmatrix} \mathbf{I}_N & \boldsymbol{\sigma} \\ \boldsymbol{\sigma}^T & \mathbb{1}_f \end{pmatrix} \begin{pmatrix} \boldsymbol{\omega} \\ \dot{\mathbf{Q}} \end{pmatrix}. \quad (4.14)$$

The metric of (4.2), however, must be expressed in the *coordinate* velocities $\dot{q} = (\dot{\phi}, \dot{\theta}, \dot{\chi}, \dot{Q}_b)$. The two are related linearly by (4.10): $\begin{pmatrix} \boldsymbol{\omega} \\ \dot{\mathbf{Q}} \end{pmatrix} = \mathbf{A} \dot{q}$ with $\mathbf{A} = \text{diag}(\mathbf{E}, \mathbb{1}_f)$. Substituting into

(4.14), the metric g_{AB} defined by $2T_{\text{rv}} = g_{AB}\dot{q}^A\dot{q}^B$ is $g = \mathbf{A}^T(\cdots)\mathbf{A}$:

$$g_{AB} = \begin{pmatrix} \mathbf{E}^T \mathbf{I}_N \mathbf{E} & \mathbf{E}^T \boldsymbol{\sigma} \\ \boldsymbol{\sigma}^T \mathbf{E} & \mathbb{1}_f \end{pmatrix}, \quad \begin{array}{ll} \text{Euler-Euler block: } \mathbf{E}^T \mathbf{I}_N \mathbf{E} & (3 \times 3), \\ \text{Euler-mode block: } \mathbf{E}^T \boldsymbol{\sigma} & (3 \times f), \\ \text{mode-mode block: } \mathbb{1}_f. & \end{array} \quad (4.15)$$

Every entry is now an explicit function of (θ, χ, Q) : the inertia tensor pulled back through the kinematic matrix, the Coriolis coupling pulled back through the same matrix, and a flat vibrational block. This single object encodes all four kinetic structures of the classical Hamiltonian (3.22) — the rotational kernel, the Coriolis coupling, the vibrational kinetic energy, and the internal $\boldsymbol{\pi}^T \boldsymbol{\mu} \boldsymbol{\pi}$ term — as we verify in § 4.7 after inverting it.

4.4 The electron sector: flat, constant, and decoupled from the measure

Before quantizing (4.15) we must account for the electrons, since the Podolsky prescription requires the metric of the *full* configuration space. Include the electron velocities: expanding the body-fixed electronic and mass-polarization kinetic terms via (3.9e), the complete velocity-form kinetic energy is

$$2T = \mathbf{v}^T \mathbf{M} \mathbf{v}, \quad \mathbf{v} = (\boldsymbol{\omega}, \dot{\mathbf{Q}}, \dot{\mathbf{r}}_i), \quad \mathbf{M} = \begin{pmatrix} \mathbf{I} & \boldsymbol{\sigma} & \mathbf{C} \\ \boldsymbol{\sigma}^T & \mathbb{1}_f & 0 \\ \mathbf{C}^T & 0 & \mathbf{D} \end{pmatrix}, \quad (4.16)$$

with $\mathbf{I} = \mathbf{I}_N + \mathbf{I}_e - \mathbf{I}_X$ the full instantaneous inertia, $\mathbf{C}$ the rotation–electron coupling block [from $\boldsymbol{\omega} \cdot (\mathbf{L}_e - \mathbf{L}_X)$ in (3.9e)], and $\mathbf{D}_{(i\alpha)(j\beta)} = (m_e \delta_{ij} - m_e^2/M_{\text{tot}}) \delta_{\alpha\beta}$ the *constant* electron mass matrix. Eliminating the electron block by its Schur complement — i.e. completing the square in $\dot{\mathbf{r}}_i$ — removes all $\boldsymbol{\omega}$ -dependence from the electron sector: the correction is precisely the electronic-plus-mass-polarization inertia, $\mathbf{C} \mathbf{D}^{-1} \mathbf{C}^T = \mathbf{I}_e - \mathbf{I}_X$, so $\mathbf{I} - \mathbf{C} \mathbf{D}^{-1} \mathbf{C}^T = \mathbf{I}_N$ — the same exact cancellation proven classically in the cross-check of § 3.6 [eq. (3.18)]. Hence

$$\det \mathbf{M} = \det \mathbf{D} \det \begin{pmatrix} \mathbf{I}_N & \boldsymbol{\sigma} \\ \boldsymbol{\sigma}^T & \mathbb{1}_f \end{pmatrix}, \quad \det \mathbf{D} = \text{const}, \quad (4.17)$$

and the surviving determinant is exactly that of the ro-vibrational block (4.14). Two consequences. (i) The electron coordinates contribute only a constant factor to the volume element — their measure is the flat $\prod_i d^3 \bar{\mathbf{r}}_i$, and $\hat{\mathbf{p}}_i = -i\hbar \nabla_{\bar{\mathbf{r}}_i}$ is Hermitian on it with no weight corrections. (ii) In the canonical (momentum) form (3.22), the electrons couple to rotation only through the shift $\mathbf{J} \rightarrow \mathbf{J} - \mathbf{L}_{\text{elec}}$, and the operator $\hat{L}_{\text{elec}, \alpha} = -i\hbar \sum_i \epsilon_{\alpha\beta\gamma} \bar{r}_{i\beta} \partial_{\bar{r}_{i\gamma}}$ is Hermitian on the flat electron measure *without any symmetrization*, because its vector field is divergence-free:

$$\sum_{i\gamma} \partial_{\bar{r}_{i\gamma}} (\epsilon_{\alpha\beta\gamma} \bar{r}_{i\beta}) = \sum_i \epsilon_{\alpha\beta\gamma} \delta_{\beta\gamma} = 0. \quad (4.18)$$

Consequently the entire quantization problem lives in the ro-vibrational sector: we quantize the metric (4.15) by the Podolsky prescription, and the electrons enter the result only through the flat kinetic operator of §4.10 and the uniform shift $\hat{J}_\alpha \rightarrow \hat{J}_\alpha - \hat{L}_{\text{elec},\alpha}$, which preserves Hermiticity term by term because $\hat{L}_{\text{elec},\alpha}$ is Hermitian and acts on a different set of coordinates than every ro-vibrational object (the term-by-term verification is Appendix C).

4.5 Determinant and the volume element

We need $g = \det g_{AB}$ for the metric (4.15). Use the block-determinant identity: for any square blocks with invertible lower-right corner,

$$\det \begin{pmatrix} \mathbf{a} & \mathbf{b} \\ \mathbf{b}^T & \mathbb{1} \end{pmatrix} = \det(\mathbf{a} - \mathbf{b} \mathbf{b}^T), \quad (4.19)$$

which follows by block row reduction — subtracting $\mathbf{b} \times$ (bottom row) from the top row produces the block-triangular matrix $\begin{pmatrix} \mathbf{a} - \mathbf{b} \mathbf{b}^T & 0 \\ \mathbf{b}^T & \mathbb{1} \end{pmatrix}$, whose determinant is the product of the diagonal blocks. Applying (4.19) to (4.15) with $\mathbf{a} = \mathbf{E}^T \mathbf{I}_N \mathbf{E}$, $\mathbf{b} = \mathbf{E}^T \boldsymbol{\sigma}$:

$$\det g_{AB} = \det[\mathbf{E}^T (\mathbf{I}_N - \boldsymbol{\sigma} \boldsymbol{\sigma}^T) \mathbf{E}] = (\det \mathbf{E})^2 \det \mathbf{I}' \implies \boxed{\det g_{AB} = \sin^2 \theta \gamma(Q), \quad \gamma \equiv \det \mathbf{I}', \quad \mathbf{I}' \equiv \mathbf{I}_N - \boldsymbol{\sigma} \boldsymbol{\sigma}^T,} \quad (4.20)$$

using $\det(\mathbf{E}^T X \mathbf{E}) = (\det \mathbf{E})^2 \det X$ and (4.11). Remarkably, the determinant is governed by the *Watson-modified* inertia $\mathbf{I}' = \mathbf{I}_N - \boldsymbol{\sigma} \boldsymbol{\sigma}^T$ of (3.20) — not the bare $\mathbf{I}_N$: the Coriolis coupling $\boldsymbol{\sigma}$ reduces the rotational determinant, and it is the same $\mathbf{I}'$ whose inverse $\boldsymbol{\mu} = (\mathbf{I}')^{-1}$ is the rotational kernel of (3.22). The Riemannian volume element is therefore

$$\boxed{\sqrt{g} d^{3+f} q = \underbrace{\sin \theta d\phi d\theta d\chi}_{\text{Haar measure on } SO(3)} \underbrace{\sqrt{\gamma(Q)} \prod_b dQ_b}_{\text{rot.-vib. Jacobian}}, \quad \sqrt{g} = \sin \theta \sqrt{\gamma(Q)},} \quad (4.21)$$

times the flat, constant electron factor $\prod_i d^3 \mathbf{r}_i$ from §4.4. The two curved factors have transparent origins: $\sin \theta$ is the Euler-angle Jacobian $|\det \mathbf{E}|$ of (4.11), and $\sqrt{\gamma}$ is the shape-dependent part carrying the Coriolis-reduced inertia.

4.6 The inverse metric

The Podolsky operator (4.5) needs g^{AB} . We claim

$$\boxed{g^{AB} = \begin{pmatrix} \mathbf{E}^{-1} \boldsymbol{\mu} \mathbf{E}^{-T} & -\mathbf{E}^{-1} \boldsymbol{\mu} \boldsymbol{\sigma} \\ -\boldsymbol{\sigma}^T \boldsymbol{\mu} \mathbf{E}^{-T} & \mathbb{1}_f + \boldsymbol{\sigma}^T \boldsymbol{\mu} \boldsymbol{\sigma} \end{pmatrix}, \quad \boldsymbol{\mu} \equiv (\mathbf{I}')^{-1} = (\mathbf{I}_N - \boldsymbol{\sigma} \boldsymbol{\sigma}^T)^{-1},} \quad (4.22)$$

with $\mathbf{E}^{-T} \equiv (\mathbf{E}^{-1})^T$ from (4.12). Rather than derive it, *verify* it — multiply (4.15) by (4.22) block by block, using only $\boldsymbol{\mu}\mathbf{I}' = \mathbb{1}$:

$$\begin{aligned}
& \text{(top-left)} \quad \mathbf{E}^T \mathbf{I}_N \mathbf{E} \cdot \mathbf{E}^{-1} \boldsymbol{\mu} \mathbf{E}^{-T} - \mathbf{E}^T \boldsymbol{\sigma} \cdot \boldsymbol{\sigma}^T \boldsymbol{\mu} \mathbf{E}^{-T} = \mathbf{E}^T (\mathbf{I}_N - \boldsymbol{\sigma} \boldsymbol{\sigma}^T) \boldsymbol{\mu} \mathbf{E}^{-T} = \mathbf{E}^T \mathbf{E}^{-T} = \mathbb{1}_3, \\
& \text{(top-right)} \quad -\mathbf{E}^T \mathbf{I}_N \mathbf{E} \cdot \mathbf{E}^{-1} \boldsymbol{\mu} \boldsymbol{\sigma} + \mathbf{E}^T \boldsymbol{\sigma} (\mathbb{1}_f + \boldsymbol{\sigma}^T \boldsymbol{\mu} \boldsymbol{\sigma}) = \mathbf{E}^T [-(\mathbf{I}_N - \boldsymbol{\sigma} \boldsymbol{\sigma}^T) \boldsymbol{\mu} \boldsymbol{\sigma} + \boldsymbol{\sigma}] = \mathbf{E}^T (-\boldsymbol{\sigma} + \boldsymbol{\sigma}) = 0, \\
& \text{(bottom-left)} \quad \boldsymbol{\sigma}^T \mathbf{E} \cdot \mathbf{E}^{-1} \boldsymbol{\mu} \mathbf{E}^{-T} - \mathbb{1}_f \cdot \boldsymbol{\sigma}^T \boldsymbol{\mu} \mathbf{E}^{-T} = 0, \\
& \text{(bottom-right)} \quad -\boldsymbol{\sigma}^T \mathbf{E} \cdot \mathbf{E}^{-1} \boldsymbol{\mu} \boldsymbol{\sigma} + \mathbb{1}_f + \boldsymbol{\sigma}^T \boldsymbol{\mu} \boldsymbol{\sigma} = \mathbb{1}_f.
\end{aligned} \tag{4.23}$$

So $g_{AB} g^{BC} = \delta_A^C$, confirming (4.22). Note how the Watson reciprocal inertia $\boldsymbol{\mu}$ — introduced classically in (3.20) by inverting the momentum–velocity relation — reappears here as the geometric inverse of the metric: the two inversions are the same computation.

4.7 The rotational derivative operators (defined, not postulated)

Expanding the Podolsky operator will repeatedly produce the combination $(\mathbf{E}^{-1})_{s\alpha} \partial_{\Omega_s}$ (index s running over ϕ, θ, χ). It is convenient to give it a name:

$$\boxed{
\begin{aligned}
& \hat{J}_1 = -i\hbar \left[-\frac{\cos \chi}{\sin \theta} \partial_\phi + \sin \chi \partial_\theta + \frac{\cos \theta \cos \chi}{\sin \theta} \partial_\chi \right], \\
\hat{J}_\alpha \equiv -i\hbar \sum_s (\mathbf{E}^{-1})_{s\alpha} \partial_{\Omega_s} : & \quad \hat{J}_2 = -i\hbar \left[\frac{\sin \chi}{\sin \theta} \partial_\phi + \cos \chi \partial_\theta - \frac{\cos \theta \sin \chi}{\sin \theta} \partial_\chi \right], \\
& \hat{J}_3 = -i\hbar \partial_\chi,
\end{aligned}
} \tag{4.24}$$

reading the rows of $\mathbf{E}^{-1}$ off (4.12). We stress the logic: (4.24) is a *definition* — shorthand for three explicit combinations of Euler-angle derivatives — and we will never compute a commutator $[\hat{J}_\alpha, \hat{J}_\beta]$ or use any angular-momentum algebra. The definition is natural because classically $J_\alpha = \partial T_{rv} / \partial \omega_\alpha$ is conjugate to $\boldsymbol{\omega}$: from $\boldsymbol{\omega} = \mathbf{E} \dot{\boldsymbol{\Omega}}$ and the chain rule, $p_{\Omega_s} = \partial T / \partial \dot{\Omega}_s = (\partial \omega_\alpha / \partial \dot{\Omega}_s) J_\alpha = (\mathbf{E})_{\alpha s} J_\alpha$, i.e. $p_\Omega = \mathbf{E}^T \mathbf{J}$ and $J_\alpha = (\mathbf{E}^{-1})_{s\alpha} p_{\Omega_s}$ — (4.24) is its coordinate representation. As a consistency check, substituting $p_\Omega = \mathbf{E}^T \mathbf{J}$ and P_b into $\frac{1}{2} p_A g^{AB} p_B$ with (4.22) collapses all $\mathbf{E}$ factors,

$$\frac{1}{2} p_A g^{AB} p_B = \frac{1}{2} \mathbf{J}^T \boldsymbol{\mu} \mathbf{J} - \mathbf{J}^T \boldsymbol{\mu} \boldsymbol{\pi} + \frac{1}{2} \boldsymbol{\pi}^T \boldsymbol{\mu} \boldsymbol{\pi} + \frac{1}{2} \sum_b P_b^2, \quad \boldsymbol{\pi} \equiv \boldsymbol{\sigma} P = \sum_b \boldsymbol{\sigma}_b P_b, \tag{4.25}$$

which is exactly the nuclear kernel of (3.22) [the electronic shift $\mathbf{J} \rightarrow \mathbf{J} - \mathbf{L}_{\text{elec}}$ enters per §4.4]. So the classical correspondence is verified before any quantization.

Two coordinate identities make the operators (4.24) as good as Cartesian momenta. First, a **divergence identity**: the weighted divergence of each column of $\mathbf{E}^{-1}$ vanishes.

Compute all three explicitly from (4.12):

$$\begin{aligned}
\alpha = 1 : \quad \sum_s \partial_{\Omega_s} (\sin \theta (E^{-1})_{s1}) &= \partial_\phi (-\cos \chi) + \partial_\theta (\sin \theta \sin \chi) + \partial_\chi (\cos \theta \cos \chi) \\
&= 0 + \cos \theta \sin \chi - \cos \theta \sin \chi = 0, \\
\alpha = 2 : \quad \sum_s \partial_{\Omega_s} (\sin \theta (E^{-1})_{s2}) &= \partial_\phi (\sin \chi) + \partial_\theta (\sin \theta \cos \chi) + \partial_\chi (-\cos \theta \sin \chi) \\
&= 0 + \cos \theta \cos \chi - \cos \theta \cos \chi = 0, \\
\alpha = 3 : \quad \sum_s \partial_{\Omega_s} (\sin \theta (E^{-1})_{s3}) &= \partial_\chi (\sin \theta) = 0.
\end{aligned} \tag{4.26}$$

Second, this identity makes each $\hat{J}_\alpha$ **Hermitian on the Haar measure** $\sin \theta d\phi d\theta d\chi$: integrating by parts (periodicity in ϕ, χ ; $\sin \theta \rightarrow 0$ at the poles),

$$\int \varphi^* \hat{J}_\alpha \psi \sin \theta d\Omega = \int (\hat{J}_\alpha \varphi)^* \psi \sin \theta d\Omega + i\hbar \int \varphi^* \psi \underbrace{\left[\sum_s \partial_{\Omega_s} (\sin \theta (E^{-1})_{s\alpha}) \right]}_{=0 \text{ by (4.26)}} d\Omega, \tag{4.27}$$

so $\hat{J}_\alpha^\dagger = \hat{J}_\alpha$ with *no* symmetrization needed. In differential-geometric language, (4.26) says the vector fields generating body-frame rotations are divergence-free with respect to the Haar measure; it is the coordinate-level fact that replaces every appeal to angular-momentum algebra in what follows. [The same argument with the flat measure and (4.18) makes $\hat{L}_{\text{elec}, \alpha}$ Hermitian, and the vibrational analogue appears in § 4.9.]

4.8 The Podolsky operator, block by block

Insert the measure (4.21) and the inverse metric (4.22) into the Podolsky operator (4.5) and split the double sum over A, B into the four blocks:

$$\hat{T}_{\text{rv}} = \underbrace{-\frac{\hbar^2}{2\sqrt{g}} \partial_{\Omega_s} (\sqrt{g} g_{st}^{\Omega\Omega} \partial_{\Omega_t})}_{\hat{T}_{\text{rot}}} - \underbrace{\frac{\hbar^2}{2\sqrt{g}} \left[\partial_{\Omega_s} (\sqrt{g} g_{sb}^{\Omega Q} \partial_{Q_b}) + \partial_{Q_b} (\sqrt{g} g_{bs}^{Q\Omega} \partial_{\Omega_s}) \right]}_{\hat{T}_{\text{Cor}}} - \underbrace{\frac{\hbar^2}{2\sqrt{g}} \partial_{Q_b} (\sqrt{g} g_{bc}^{QQ} \partial_{Q_c})}_{\hat{T}_{\text{vib}}}. \tag{4.28}$$

Rotational block. In $\hat{T}_{\text{rot}}$, the factors $\sqrt{g}$, $\mu_{\alpha\beta}(Q)$ are independent of the Euler angles, so they pass through ∂_{Ω_s} and the $\sqrt{g}$ cancels against the prefactor. Writing $g_{st}^{\Omega\Omega} = (E^{-1})_{s\alpha} \mu_{\alpha\beta} (E^{-1})_{t\beta}$ and expanding the outer derivative by the product rule,

$$\begin{aligned}
\hat{T}_{\text{rot}} &= -\frac{\hbar^2}{2} \mu_{\alpha\beta} \left[\underbrace{(E^{-1})_{s\alpha} \partial_{\Omega_s} ((E^{-1})_{t\beta} \partial_{\Omega_t})}_{\text{both derivatives survive}} + \underbrace{\frac{1}{\sin \theta} \left(\sum_s \partial_{\Omega_s} (\sin \theta (E^{-1})_{s\alpha}) \right)}_{=0 \text{ by (4.26)}} (E^{-1})_{t\beta} \partial_{\Omega_t} \right] \\
&= \frac{1}{2} \sum_{\alpha\beta} \mu_{\alpha\beta}(Q) \hat{J}_\alpha \hat{J}_\beta,
\end{aligned} \tag{4.29}$$

by the definition (4.24). The divergence identity (4.26) has done all the work: the rotational block is *exactly* the symmetric kernel $\frac{1}{2} \mu_{\alpha\beta} \hat{J}_\alpha \hat{J}_\beta$, Hermitian as written [each $\hat{J}_\alpha$ is Hermitian by (4.27), $\mu_{\alpha\beta}$ is a real symmetric function of Q commuting with both, and taking the adjoint merely reverses the order, $(\mu_{\alpha\beta} \hat{J}_\alpha \hat{J}_\beta)^\dagger = \mu_{\alpha\beta} \hat{J}_\beta \hat{J}_\alpha$, which upon relabeling the

summed indices $\alpha \leftrightarrow \beta$ and using $\mu_{\beta\alpha} = \mu_{\alpha\beta}$ is the original operator], and it contributes *no* extra potential term. In the traditional treatment this last fact is derived from the anomalous commutator algebra; here it is simply the statement that a rigid rotor has no pseudopotential, verified by the three explicit derivative computations (4.26).

Coriolis block. In $\hat{T}_{\text{Cor}}$, the first group carries ∂_{Ω_s} , which passes through the Q -functions $\mu, \sigma, \sqrt{\gamma}$ and, by the same divergence identity (4.26), collapses onto $\hat{J}_\alpha$ acting from the left. The second group carries ∂_{Q_b} , which passes through $\mathbf{E}^{-1}(\Omega)$ and $\sin\theta$ but *acts* on $\sqrt{\gamma}, \mu$, and σ . The result (each step is one product rule; the full expansion is Appendix C) is

$$\hat{T}_{\text{Cor}} = -\frac{1}{2} \left[\hat{J}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta + \frac{1}{\sqrt{\gamma}} \hat{P}_b \sqrt{\gamma} \sigma_{\alpha b} \mu_{\alpha\beta} \hat{J}_\beta \right], \quad \hat{P}_b \equiv -i\hbar \partial_{Q_b}, \quad \hat{\pi}_\alpha \equiv \sum_b \sigma_{\alpha b}(Q) \hat{P}_b. \quad (4.30)$$

The two groups are adjoints of each other on the measure (4.21) [the $\gamma^{\pm 1/2}$ factors in the second group are precisely what integration by parts over $\sqrt{\gamma} dQ$ produces], so their sum is Hermitian — again automatically, with no ordering imposed by hand.

Vibrational block. With $g_{bc}^{QQ} = \delta_{bc} + (\sigma^T \mu \sigma)_{bc}$ and the $\sin\theta$ cancelling,

$$\hat{T}_{\text{vib}} = -\frac{\hbar^2}{2} \frac{1}{\sqrt{\gamma}} \sum_{b,c} \partial_{Q_b} \left(\sqrt{\gamma} [\delta_{bc} + (\sigma^T \mu \sigma)_{bc}] \partial_{Q_c} \right). \quad (4.31)$$

Expanding the δ_{bc} part by one product rule exhibits the problem announced in (4.4): $-\frac{\hbar^2}{2} \gamma^{-1/2} \partial_{Q_b} (\gamma^{1/2} \partial_{Q_b} \psi) = \frac{1}{2} \hat{P}_b^2 \psi - \frac{\hbar^2}{2} (\frac{1}{2} \partial_{Q_b} \ln \gamma) \partial_{Q_b} \psi$ — the bare $\hat{P}_b$ is not Hermitian on the curved measure $\sqrt{\gamma} dQ$, and the first-order remainder is the compensation. Removing these remainders once and for all is the job of the rescaling below.

4.9 Rescaling to the flat measure and the Watson pseudopotential

To work with the simpler measure $d\mu_W \equiv \sin\theta d\phi d\theta d\chi \prod_b dQ_b \prod_i d^3\mathbf{r}_i$, absorb the shape factor of the volume element into the wavefunction:

$$\psi_W \equiv \gamma^{1/4} \psi \quad \implies \quad |\psi|^2 \sqrt{g} d^{3+f}q = |\psi_W|^2 d\mu_W, \quad (4.32)$$

so $\psi_W \in L^2(d\mu_W)$, and conjugate every operator: $\hat{O}_W = \gamma^{1/4} \hat{O} \gamma^{-1/4}$. Only the Q -sector is affected (γ depends on Q alone); the $\sin\theta$ stays in the measure, on which the $\hat{J}_\alpha$ are already Hermitian by (4.27), and the electron sector is flat. Conjugating the identity part of (4.31), working from the innermost derivative outward (each line is one product rule):

$$\begin{aligned} \partial_{Q_b} (\gamma^{-1/4} \psi_W) &= \gamma^{-1/4} \left[\partial_{Q_b} \psi_W - \frac{1}{4} (\partial_{Q_b} \ln \gamma) \psi_W \right], \\ \gamma^{1/2} \partial_{Q_b} (\gamma^{-1/4} \psi_W) &= \gamma^{1/4} \left[\partial_{Q_b} \psi_W - \frac{1}{4} (\partial_{Q_b} \ln \gamma) \psi_W \right], \\ \partial_{Q_b} \left(\gamma^{1/4} [\dots] \right) &= \gamma^{1/4} \left[\partial_{Q_b}^2 \psi_W - \frac{1}{16} (\partial_{Q_b} \ln \gamma)^2 \psi_W - \frac{1}{4} (\partial_{Q_b}^2 \ln \gamma) \psi_W \right], \end{aligned} \quad (4.33)$$

where in the last line the two cross terms $\pm \frac{1}{4} (\partial_{Q_b} \ln \gamma) \partial_{Q_b} \psi_W$ (one from differentiating $\gamma^{1/4}$, one from the bracket) cancel. Multiplying by $-\frac{\hbar^2}{2} \gamma^{1/4} \cdot \gamma^{-1/2} = -\frac{\hbar^2}{2} \gamma^{-1/4}$ and summing

over b : the identity part becomes the bare $\frac{1}{2} \sum_b \hat{P}_b^2$ plus a pure multiplicative function,

$$U_{\text{vib}} = \frac{\hbar^2}{8} \sum_b \left[\partial_{Q_b}^2 \ln \gamma + \frac{1}{4} (\partial_{Q_b} \ln \gamma)^2 \right]. \quad (4.34)$$

The Coriolis block responds even more simply. Conjugating (4.30): in the first group, $\gamma^{1/4} \hat{P}_b \gamma^{-1/4} = \hat{P}_b + \frac{i\hbar}{4} (\partial_{Q_b} \ln \gamma)$; in the second, $\gamma^{-1/4} \hat{P}_b \gamma^{1/4} = \hat{P}_b - \frac{i\hbar}{4} (\partial_{Q_b} \ln \gamma)$. The two induced terms are equal and opposite [they multiply the same function, since $\hat{J}_\alpha \mu_{\alpha\beta} \sigma_{\beta b} = \sigma_{\beta b} \mu_{\alpha\beta} \hat{J}_\alpha$ upon relabeling the summed Cartesian indices — these are functions of different variables and commute], so they cancel, leaving the manifestly symmetric flat-measure form

$$\hat{T}_{\text{Cor},W} = -\frac{1}{2} \sum_{\alpha\beta} \hat{J}_\alpha (\mu_{\alpha\beta} \hat{\pi}_\beta + \hat{\pi}_\beta \mu_{\alpha\beta}) = -\frac{1}{2} (\hat{J}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta + \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{J}_\beta), \quad (4.35)$$

with *no* extra potential from this block. Finally, conjugating the $\sigma^T \mu \sigma$ dressing of (4.31) and reorganizing it into the symmetric kernel ordering $\frac{1}{2} \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta$ leaves a second multiplicative residue, U_{Cor} (every derivative of this bookkeeping is written out in Appendix C).

The total extra potential $\hat{U} = U_{\text{vib}} + U_{\text{Cor}}$ collapses through two exact identities, both pure Q -calculus. First, Jacobi's formula — for a symmetric positive-definite matrix, $\ln \det \mathbf{I}' = \text{tr} \ln \mathbf{I}'$, so differentiating,

$$\partial_{Q_b} \ln \gamma = \text{tr}(\mu \partial_{Q_b} \mathbf{I}'), \quad \partial_{Q_b} \mu = -\mu (\partial_{Q_b} \mathbf{I}') \mu \quad [\text{from } \partial_{Q_b}(\mu \mathbf{I}') = 0], \quad (4.36)$$

expresses every derivative of γ and μ through $\partial_{Q_b} \mathbf{I}'$. Second, because $\mathbf{I}' = \mathbf{I}_N - \sigma \sigma^T$ with $\sigma_b = \sum_a \zeta_{ab} Q_a$ linear in Q with constant ζ 's,

$$\partial_{Q_c} \mathbf{I}'_{\alpha\beta} = -\sum_b (\zeta_{cb}^\alpha \sigma_{\beta b} + \sigma_{\alpha b} \zeta_{cb}^\beta), \quad \zeta_{cb}^\alpha \equiv (\zeta_{cb})_\alpha = \partial_{Q_c} \sigma_{\alpha b} \quad (\text{constant}), \quad (4.37)$$

so all derivatives reduce to the constant Coriolis matrices $\zeta_{cb} = \sum_j \mathbf{l}_{jc} \times \mathbf{l}_{jb}$. Substituting (4.36)–(4.37) into $U_{\text{vib}} + U_{\text{Cor}}$, the gradient-type terms generated by the two groups cancel identically, and the surviving Q -local remainder is fixed by the sum rules the ζ 's inherit from the orthonormality $\sum_j \mathbf{l}_{ja} \cdot \mathbf{l}_{jb} = \delta_{ab}$ of the Eckart mode vectors; the outcome, first obtained by Watson [3], is the single trace

$$\boxed{\hat{U}(Q) = -\frac{\hbar^2}{8} \sum_{\alpha=1}^3 \mu_{\alpha\alpha}(Q), \quad \mu = (\mathbf{I}_N - \sigma \sigma^T)^{-1}.} \quad (4.38)$$

Two remarks. (a) The collapse is special to *rectilinear* normal coordinates: for general curvilinear internal coordinates the linearity (4.37) fails and additional extra-potential terms survive. (b) The electrons do not modify $\hat{U}$: γ and μ depend on Q alone, and the electronic Jacobian is flat (§ 4.4). The pseudopotential is a c -number on the nuclear shape space, fixed entirely by the modified inertia $\mathbf{I}'$.

4.10 Electronic kinetic operator

In BF Cartesian electron coordinates, $\hat{\mathbf{p}}_i = -i\hbar \nabla_{\mathbf{r}_i}$ is Hermitian on $L^2(\mathbb{R}^{3N_{\text{elec}}}, \prod_i d^3 \mathbf{r}_i)$ (§ 4.4). The electronic + mass-polarization kinetic operator is therefore the straightforward

quantization of (3.21):

$$\hat{T}_e = \sum_i \frac{\hat{\mathbf{p}}_i^2}{2m_e} + \frac{1}{2M_N} \left(\sum_i \hat{\mathbf{p}}_i \right)^2. \quad (4.39)$$

Expanding the second piece:

$$\frac{1}{2M_N} \left(\sum_i \hat{\mathbf{p}}_i \right)^2 = \sum_i \frac{\hat{\mathbf{p}}_i^2}{2M_N} + \frac{1}{M_N} \sum_{i < l} \hat{\mathbf{p}}_i \cdot \hat{\mathbf{p}}_l. \quad (4.40)$$

The diagonal $1/(2M_N)$ piece renormalizes the effective electron mass through $1/m_e + 1/M_N = M_{\text{tot}}/(m_e M_N) = 1/\mu_e$. The off-diagonal two-electron term is the mass-polarization *sensu stricto*, dynamically coupling electrons through nuclear recoil — a purely classical effect (Section 2.4) inherited by the quantum theory.

4.11 Complete quantum molecular Hamiltonian

Collect the flat-measure blocks: the rotational kernel (4.29), the Coriolis coupling (4.35), the vibrational operator [bare $\frac{1}{2} \sum_b \hat{P}_b^2$ plus the symmetric $\frac{1}{2} \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta$], the pseudopotential (4.38), the electronic operator (4.39), the shift $\hat{J}_\alpha \rightarrow \hat{J}_\alpha - \hat{L}_{\text{elec},\alpha}$ of § 4.4, and the rotationally invariant Coulomb potentials of (3.8). The pieces assemble into the sandwich-ordered kernel [the term-by-term identification of the nine products with the blocks above, including every derivative, is Appendix C]:

$$\begin{aligned} \hat{H}_{\text{int}} = & \frac{1}{2} \sum_{\alpha\beta} (\hat{J}_\alpha - \hat{L}_{\text{elec},\alpha} - \hat{\pi}_\alpha) \mu_{\alpha\beta}(Q) (\hat{J}_\beta - \hat{L}_{\text{elec},\beta} - \hat{\pi}_\beta) \\ & + \frac{1}{2} \sum_b \hat{P}_b^2 \\ & + \sum_i \frac{\hat{\mathbf{p}}_i^2}{2m_e} + \frac{1}{2M_N} \left(\sum_i \hat{\mathbf{p}}_i \right)^2 \\ & - \frac{\hbar^2}{8} \sum_\alpha \mu_{\alpha\alpha}(Q) \\ & + V_{NN}(Q) + V_{Ne}(Q, \{\mathbf{r}_i\}) + V_{ee}(\{\mathbf{r}_i\}), \end{aligned} \quad (4.41)$$

acting on $L^2(\mathcal{C}, d\mu_W)$ with $d\mu_W = \sin \theta d\phi d\theta d\chi \prod_b dQ_b \prod_i d^3\mathbf{r}_i$. Every operator in (4.41) is an explicit differential operator in the coordinates — the $\hat{J}_\alpha$ are the Euler-angle combinations (4.24), $\hat{\pi}_\alpha = \sum_b \sigma_{\alpha b} \hat{P}_b$, $\hat{P}_b = -i\hbar \partial_{Q_b}$, $\hat{\mathbf{p}}_i = -i\hbar \nabla_{\mathbf{r}_i}$ — and every one is Hermitian on $d\mu_W$ by the integration-by-parts arguments given above.

This is the **exact non-relativistic molecular Hamiltonian** after separation of the total COM motion, in BF + Eckart frame, with electrons referred to the nuclear COM. The only approximations made along the way were: (i) non-relativistic dynamics; (ii) Eckart's choice of the BF orientation (a convention, not an approximation); (iii) parametrization of the nuclear shape by $3N_{\text{nuc}} - 6$ internal coordinates Q_b (a parametrization, also exact).

Remark (the anomalous commutator, not needed). The traditional route to (4.41) [14, 7, 4] defines the body-fixed angular momentum by projecting the space-fixed operator through the direction cosines, derives its *anomalous* commutation relations $[\hat{J}_\alpha, \hat{J}_\beta] = -i\hbar \epsilon_{\alpha\beta\gamma} \hat{J}_\gamma$, and uses that algebra to control the operator ordering of the rotational ker-

nel. Nothing in this section used it: the $\hat{J}_\alpha$ entered only as the shorthand (4.24) for explicit Euler-angle derivatives, Hermiticity came from integration by parts with the divergence identity (4.26), and the pseudopotential came from the Q -derivatives of the volume element. The anomalous algebra is a correct and sometimes useful property of the operators (4.24), but it is a *consequence* of the coordinate representation, not an input to the quantization.

4.12 Born–Huang expansion

4.12.1 Clamped-nuclei electronic Hamiltonian

At each nuclear configuration Q , define the **BF clamped-nuclei electronic Hamiltonian** as the sum of all Q -parametric electron-only pieces of (4.41):

$$\hat{H}_{\text{el}}(Q) \equiv \sum_i \frac{\hat{\mathbf{p}}_i^2}{2m_e} + \frac{1}{2M_N} \left(\sum_i \hat{\mathbf{p}}_i \right)^2 + V_{Ne}(Q, \{\bar{\mathbf{r}}_i\}) + V_{ee}(\{\bar{\mathbf{r}}_i\}). \quad (4.42)$$

Remark. The mass-polarization piece $(\sum_i \hat{\mathbf{p}}_i)^2/(2M_N)$ is included here as a small correction with $1/M_N$. Some treatments (“clamped-nuclei BO”) drop it; we keep it explicit because it is parametric in Q only through the absent dependence (i.e., M_N is just a number), which means it acts trivially on the parametric family of electronic Hilbert spaces.

Solve the parametric eigenvalue problem:

$$\hat{H}_{\text{el}}(Q) |\phi_n(Q)\rangle = E_n^{\text{el}}(Q) |\phi_n(Q)\rangle, \quad (4.43)$$

in the electronic Hilbert space $\mathcal{H}_e = L^2(\mathbb{R}^{3N_{\text{elec}}}, d^{3N_{\text{elec}}} \bar{\mathbf{r}})$. The eigenstates $\{|\phi_n(Q)\rangle\}$ depend *parametrically* on Q — Q is treated as a fixed classical parameter, not a quantum degree of freedom — and form a complete orthonormal basis of $\mathcal{H}_e$ at each Q :

$$\sum_n |\phi_n(Q)\rangle \langle \phi_n(Q)| = \hat{\mathbb{1}}_e, \quad \langle \phi_m(Q) | \phi_n(Q) \rangle_e = \delta_{mn}. \quad (4.44)$$

4.12.2 The exact Born–Huang expansion

The *exact* molecular wavefunction $|\Psi\rangle \in L^2(\mathcal{C}, d\mu_W)$ admits a resolution against the parametric electronic basis. Insert the resolution of identity (4.44) at each Q :

$$|\Psi\rangle = \sum_n |\chi_n\rangle \otimes |\phi_n(Q)\rangle, \quad \chi_n(\Omega, Q) \equiv \langle \phi_n(Q) | \Psi \rangle_e, \quad (4.45)$$

where $\Omega = (\phi, \theta, \chi)$, the electronic inner product is taken at fixed Q , and χ_n are the **nuclear coefficient functions**. The expansion (4.45) is **exact**, not an ansatz: it follows from the completeness (4.44) at each Q , with the only subtlety being that the basis $|\phi_n(Q)\rangle$ varies parametrically. There is *no* simple tensor factorization $\mathcal{H}_{\text{molecule}} = \mathcal{H}_N \otimes \mathcal{H}_e$ because of this parametric dependence; the Born–Huang expansion [11] is the rigorous replacement.

4.12.3 Coupled-channel equations

Substitute (4.45) into $\hat{H}_{\text{int}}|\Psi\rangle = E|\Psi\rangle$, project from the left by $\langle\phi_m(Q)|$, and use (4.43):

$$[\hat{T}_N + E_n^{\text{el}}(Q) + V_{NN}(Q) + \hat{U}(Q)]\chi_n(\Omega, Q) + \sum_m \hat{\Lambda}_{nm} \chi_m(\Omega, Q) = E \chi_n(\Omega, Q), \quad (4.46)$$

where $\hat{T}_N$ denotes the nuclear (rotational + vibrational) kinetic operator from the first two lines of (4.41). The **non-adiabatic coupling operators** $\hat{\Lambda}_{nm}$ arise from the action of $\hat{T}_N$ (and the rot–electronic couplings through $\hat{\mathbf{L}}_{\text{elec}}$) on the Q -parametric dependence of $|\phi_m(Q)\rangle$:

$$\hat{\Lambda}_{nm} \sim -\hbar^2 \langle\phi_n|\nabla_Q\phi_m\rangle_e \cdot \nabla_Q - \frac{\hbar^2}{2} \langle\phi_n|\nabla_Q^2\phi_m\rangle_e + \dots, \quad (4.47)$$

with analogous terms from the rotation–electron Coriolis $\hat{J}_\alpha \hat{L}_{\text{elec},\alpha}$ kernel.

4.12.4 Born–Oppenheimer approximation as truncation

The **Born–Oppenheimer (BO) approximation** is the diagonal truncation of (4.46): set $\hat{\Lambda}_{nm} \rightarrow 0$ for $m \neq n$ and (optionally) also $\hat{\Lambda}_{nn} \rightarrow 0$. The resulting effective nuclear problem is

$$[\hat{T}_N + E_n^{\text{el}}(Q) + V_{NN}(Q) + \hat{U}(Q)]\chi_n^{\text{BO}}(\Omega, Q) = E^{\text{BO}} \chi_n^{\text{BO}}(\Omega, Q), \quad (4.48)$$

where the **Born–Oppenheimer potential energy surface** for electronic state n is

$$V_n^{\text{BO}}(Q) \equiv E_n^{\text{el}}(Q) + V_{NN}(Q) + \hat{U}(Q). \quad (4.49)$$

The exactness of (4.45) and the well-definedness of (4.46) are mathematical facts. BO is the physical assumption that off-diagonal $\hat{\Lambda}_{nm}$ are small — justified when electronic gaps are large compared to nuclear kinetic energies, breaking down at conical intersections and avoided crossings, where the parametric variation of $|\phi_n(Q)\rangle$ becomes singular.

4.12.5 Summary

1. (4.41) is the exact non-relativistic quantum molecular Hamiltonian in BF + Eckart + nuclear-COM coordinates, obtained entirely from coordinate derivatives: metric (4.15) $\rightarrow$ measure (4.21) $\rightarrow$ inverse (4.22) $\rightarrow$ Laplace–Beltrami blocks (4.29)–(4.31) $\rightarrow$ rescaling and pseudopotential (4.38).
2. The Born–Huang expansion (4.45) is exact, following from completeness of the parametric electronic basis (4.44).
3. The infinite coupled system (4.46) is equivalent to the Schrödinger equation $\hat{H}_{\text{int}}|\Psi\rangle = E|\Psi\rangle$.
4. The Born–Oppenheimer truncation reduces it to (4.48), giving the familiar single-PES picture (4.49).

Appendix

Appendix A: Mass-Polarization Term in Classical Molecular Hamiltonian

We start from the laboratory-frame kinetic energy

$$T = \frac{1}{2} \sum_j M_j \dot{\mathbf{R}}_j^2 + \frac{1}{2} \sum_i m_e \dot{\mathbf{r}}_i^2, \quad (1)$$

where M_j is the mass of the j -th nucleus, $\mathbf{R}_j$ is its Cartesian coordinate, m_e is the electron mass, and $\mathbf{r}_i$ is the coordinate of the i -th electron.

Nuclear Center-of-Mass Coordinate

Define the nuclear total mass

$$M_N = \sum_j M_j. \quad (2)$$

The nuclear center-of-mass coordinate is

$$\mathbf{R}_{\text{cm}} = \mathbf{R}_N^{\text{CoM}} = \frac{1}{M_N} \sum_j M_j \mathbf{R}_j. \quad (3)$$

We introduce coordinates shifted by the nuclear center of mass:

$$\mathbf{R}'_j = \mathbf{R}_j - \mathbf{R}_{\text{cm}}, \quad (4)$$

$$\mathbf{r}'_i = \mathbf{r}_i - \mathbf{R}_{\text{cm}}. \quad (5)$$

Therefore,

$$\mathbf{R}_j = \mathbf{R}_{\text{cm}} + \mathbf{R}'_j, \quad \mathbf{r}_i = \mathbf{R}_{\text{cm}} + \mathbf{r}'_i. \quad (6)$$

The shifted nuclear coordinates satisfy the constraint

$$\sum_j M_j \mathbf{R}'_j = 0, \quad (7)$$

and therefore

$$\sum_j M_j \dot{\mathbf{R}}'_j = 0. \quad (8)$$

This is important: the $\mathbf{R}'_j$ are not all independent.

Kinetic Energy in Velocity Form

Substitute

$$\dot{\mathbf{R}}_j = \dot{\mathbf{R}}_{\text{cm}} + \dot{\mathbf{R}}'_j, \quad \dot{\mathbf{r}}_i = \dot{\mathbf{R}}_{\text{cm}} + \dot{\mathbf{r}}'_i \quad (9)$$

into the kinetic energy. For the nuclear part,

$$T_N = \frac{1}{2} \sum_j M_j \left(\dot{\mathbf{R}}_{\text{cm}} + \dot{\mathbf{R}}'_j \right)^2 \quad (10)$$

$$= \frac{1}{2} M_N \dot{\mathbf{R}}_{\text{cm}}^2 + \dot{\mathbf{R}}_{\text{cm}} \cdot \sum_j M_j \dot{\mathbf{R}}'_j + \frac{1}{2} \sum_j M_j \dot{\mathbf{R}}_j'^2. \quad (11)$$

Using

$$\sum_j M_j \dot{\mathbf{R}}'_j = 0, \quad (12)$$

we get

$$T_N = \frac{1}{2} M_N \dot{\mathbf{R}}_{\text{cm}}^2 + \frac{1}{2} \sum_j M_j \dot{\mathbf{R}}_j'^2. \quad (13)$$

For the electronic part,

$$T_e = \frac{1}{2} \sum_i m_e \left(\dot{\mathbf{R}}_{\text{cm}} + \dot{\mathbf{r}}'_i \right)^2 \quad (14)$$

$$= \frac{1}{2} N_{\text{elec}} m_e \dot{\mathbf{R}}_{\text{cm}}^2 + m_e \dot{\mathbf{R}}_{\text{cm}} \cdot \sum_i \dot{\mathbf{r}}'_i + \frac{1}{2} \sum_i m_e \dot{\mathbf{r}}_i'^2, \quad (15)$$

where N_{elec} is the number of electrons. Hence the total kinetic energy is

$$T = \frac{1}{2} (M_N + N_{\text{elec}} m_e) \dot{\mathbf{R}}_{\text{cm}}^2 + m_e \dot{\mathbf{R}}_{\text{cm}} \cdot \sum_i \dot{\mathbf{r}}'_i + \frac{1}{2} \sum_j M_j \dot{\mathbf{R}}_j'^2 + \frac{1}{2} \sum_i m_e \dot{\mathbf{r}}_i'^2. \quad (16)$$

Canonical Momenta

The momentum conjugate to the nuclear center-of-mass coordinate $\mathbf{R}_{\text{cm}}$ is

$$\mathbf{P}_{\text{cm}} = \frac{\partial \mathcal{L}}{\partial \dot{\mathbf{R}}_{\text{cm}}} \quad (17)$$

$$= (M_N + N_{\text{elec}} m_e) \dot{\mathbf{R}}_{\text{cm}} + m_e \sum_i \dot{\mathbf{r}}'_i. \quad (18)$$

Since

$$\mathbf{p}'_i = \frac{\partial \mathcal{L}}{\partial \dot{\mathbf{r}}'_i} = m_e \left(\dot{\mathbf{R}}_{\text{cm}} + \dot{\mathbf{r}}'_i \right), \quad (19)$$

we can write

$$\sum_i \mathbf{p}'_i = N_{\text{elec}} m_e \dot{\mathbf{R}}_{\text{cm}} + m_e \sum_i \dot{\mathbf{r}}'_i. \quad (20)$$

Therefore,

$$\boxed{\mathbf{P}_{\text{cm}} = M_N \dot{\mathbf{R}}_{\text{cm}} + \sum_i \mathbf{p}'_i} \quad (21)$$

or equivalently,

$$\boxed{M_N \dot{\mathbf{R}}_{\text{cm}} = \mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}'_i} \quad (22)$$

and hence

$$\dot{\mathbf{R}}_{\text{cm}} = \frac{1}{M_N} \left(\mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}'_i \right). \quad (23)$$

The electron momentum conjugate to $\mathbf{r}'_i$ is

$$\mathbf{p}'_i = m_e \left(\dot{\mathbf{R}}_{\text{cm}} + \dot{\mathbf{r}}'_i \right). \quad (24)$$

Thus $\mathbf{p}'_i$ is the lab-frame electron momentum, not simply $m_e \dot{\mathbf{r}}'_i$. Equivalently,

$$m_e \dot{\mathbf{r}}'_i = \mathbf{p}'_i - m_e \dot{\mathbf{R}}_{\text{cm}}. \quad (25)$$

The nuclear shifted momenta are

$$\mathbf{P}'_j = M_j \dot{\mathbf{R}}'_j. \quad (26)$$

Because the shifted nuclear coordinates satisfy

$$\sum_j M_j \mathbf{R}'_j = 0, \quad (27)$$

their conjugate momenta satisfy

$$\sum_j \mathbf{P}'_j = 0. \quad (28)$$

In terms of the laboratory-frame nuclear momenta

$$\mathbf{P}_j = M_j \dot{\mathbf{R}}_j, \quad (29)$$

one has

$$\mathbf{P}'_j = \mathbf{P}_j - \frac{M_j}{M_N} \mathbf{P}_N, \quad (30)$$

where

$$\mathbf{P}_N = \sum_j \mathbf{P}_j = M_N \dot{\mathbf{R}}_{\text{cm}}. \quad (31)$$

Using

$$\mathbf{P}_N = \mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}'_i, \quad (32)$$

we may also write

$$\mathbf{P}'_j = \mathbf{P}_j - \frac{M_j}{M_N} \left(\mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}'_i \right). \quad (33)$$

Kinetic Energy in Momentum Form

The original kinetic energy can be written as

$$T = T_{\text{nuc,int}} + T_{\text{nuc,CoM}} + T_e. \quad (34)$$

The nuclear internal kinetic energy is

$$T_{\text{nuc,int}} = \sum_j \frac{\mathbf{P}_j'^2}{2M_j}. \quad (35)$$

The electron kinetic energy is

$$T_e = \sum_i \frac{\mathbf{p}_i'^2}{2m_e}. \quad (36)$$

The nuclear center-of-mass kinetic energy is

$$T_{\text{nuc,CoM}} = \frac{1}{2} M_N \dot{\mathbf{R}}_{\text{cm}}^2. \quad (37)$$

Using

$$M_N \dot{\mathbf{R}}_{\text{cm}} = \mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}_i', \quad (38)$$

we get

$$T_{\text{nuc,CoM}} = \frac{1}{2M_N} \left(\mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}_i' \right)^2. \quad (39)$$

Therefore the kinetic energy in terms of the new momenta is

$$T = \sum_j \frac{\mathbf{P}_j'^2}{2M_j} + \sum_i \frac{\mathbf{p}_i'^2}{2m_e} + \frac{1}{2M_N} \left(\mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}_i' \right)^2. \quad (40)$$

The accompanying constraints are

$$\sum_j M_j \mathbf{R}_j' = 0, \quad \sum_j \mathbf{P}_j' = 0. \quad (41)$$

Expanded Form

Expanding the center-of-mass recoil term gives

$$T = \sum_j \frac{\mathbf{P}_j'^2}{2M_j} + \sum_i \frac{\mathbf{p}_i'^2}{2m_e} + \frac{\mathbf{P}_{\text{cm}}^2}{2M_N} - \frac{\mathbf{P}_{\text{cm}}}{M_N} \cdot \sum_i \mathbf{p}_i' + \frac{1}{2M_N} \left(\sum_i \mathbf{p}_i' \right)^2. \quad (42)$$

Thus,

$$T = \sum_j \frac{\mathbf{P}_j'^2}{2M_j} + \sum_i \frac{\mathbf{p}_i'^2}{2m_e} + \frac{\mathbf{P}_{\text{cm}}^2}{2M_N} - \frac{\mathbf{P}_{\text{cm}}}{M_N} \cdot \sum_i \mathbf{p}_i' + \frac{1}{2M_N} \left(\sum_i \mathbf{p}_i' \right)^2. \quad (43)$$

Hamiltonian

If the potential energy is translationally invariant, then

$$V = V(\{\mathbf{R}_j'\}, \{\mathbf{r}_i'\}). \quad (44)$$

The Hamiltonian is therefore

$$H = \sum_j \frac{\mathbf{P}_j'^2}{2M_j} + \sum_i \frac{\mathbf{p}_i'^2}{2m_e} + \frac{1}{2M_N} \left(\mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}_i' \right)^2 + V(\{\mathbf{R}_j'\}, \{\mathbf{r}_i'\}). \quad (45)$$

Molecular Rest Frame

If the whole molecule is taken to be in its rest frame, then

$$\mathbf{P}_{\text{cm}} = 0. \quad (46)$$

The kinetic energy becomes

$$T_{\text{rest}} = \sum_j \frac{\mathbf{P}_j'^2}{2M_j} + \sum_i \frac{\mathbf{p}_i'^2}{2m_e} + \frac{1}{2M_N} \left(\sum_i \mathbf{p}_i' \right)^2. \quad (47)$$

The last term,

$$\frac{1}{2M_N} \left(\sum_i \mathbf{p}_i' \right)^2, \quad (48)$$

is the nuclear center-of-mass recoil term, also called the mass-polarization term.

Important Interpretation

The coordinate transformation used here shifts both nuclei and electrons by the nuclear center of mass, not by the total molecular center of mass. Therefore,

$$\mathbf{p}_i' = m_e \left(\dot{\mathbf{R}}_{\text{cm}} + \dot{\mathbf{r}}_i' \right) \quad (49)$$

is the canonical momentum conjugate to $\mathbf{r}_i'$. It is equal to the electron lab-frame mechanical momentum. However,

$$m_e \dot{\mathbf{r}}_i' \quad (50)$$

is only the electron momentum relative to the moving nuclear center of mass. It is not the canonical momentum conjugate to $\mathbf{r}_i'$ unless the nuclear center-of-mass motion has been fixed or eliminated. The final kinetic energy in nuclear-CoM coordinates is therefore

$$T = \sum_j \frac{\mathbf{P}_j'^2}{2M_j} + \sum_i \frac{\mathbf{p}_i'^2}{2m_e} + \frac{1}{2M_N} \left(\mathbf{P}_{\text{cm}} - \sum_i \mathbf{p}_i' \right)^2, \quad (51)$$

with

$$\sum_j M_j \mathbf{R}_j' = 0, \quad \sum_j \mathbf{P}_j' = 0. \quad (52)$$

Appendix C: Term-by-term expansion of the Podolsky operator, with every derivative

Section 4 quantized the classical Hamiltonian (3.22) by applying the Podolsky operator (4.5) to the ro-vibrational metric (4.15) and rescaling to the flat measure. This appendix supplies what the main text compressed: the expansion of the quantized kernel of (4.41),

$$\hat{T}_{\text{kernel}} = \frac{1}{2} \sum_{\alpha\beta} \hat{A}_\alpha \mu_{\alpha\beta}(Q) \hat{A}_\beta, \quad \hat{A}_\alpha \equiv \hat{J}_\alpha - \hat{L}_\alpha - \hat{\pi}_\alpha, \quad \hat{L}_\alpha \equiv \hat{L}_{\text{elec},\alpha}, \quad (\text{C.1})$$

into its nine products, with *every* coordinate derivative written out and the Hermiticity of each block verified from first principles. All arguments use only: the adjoint of a product reverses the order, $(\hat{X}\hat{Y})^\dagger = \hat{Y}^\dagger\hat{X}^\dagger$; integration by parts on the measure $d\mu_W$; and the product rule. No commutator algebra of body-fixed angular momenta appears anywhere (cf. the Remark in §4.11). Repeated Cartesian indices $\alpha, \beta \in \{1, 2, 3\}$ and mode indices $b, c \in \{1, \dots, f\}$ are summed.

C.1 Toolkit

The three momentum-like operators live in three separate coordinate sectors:

$$\hat{J}_\alpha = -i\hbar (\mathbf{E}^{-1})_{s\alpha} \partial_{\Omega_s} \text{ [eq. (4.24), Euler angles]}, \quad \hat{P}_b = -i\hbar \partial_{Q_b} \text{ [modes]}, \quad \hat{L}_\alpha = -i\hbar \epsilon_{\alpha\beta\gamma} \sum_i \bar{r}_{i\beta} \partial_{\bar{r}_{i\gamma}} \text{ [electrons]}. \quad (\text{C.2})$$

Because they differentiate disjoint sets of variables, *operators from different sectors commute*, and each sector's functions are constants for the other sectors' derivatives. Each is Hermitian on its factor of $d\mu_W$: $\hat{J}_\alpha$ on $\sin\theta d\Omega$ by the divergence identity (4.26) [via (4.27)], $\hat{L}_\alpha$ on the flat electron measure by (4.18), and $\hat{P}_b$ on the flat $\prod_b dQ_b$ trivially. The only non-commutativity in the problem is the product rule within the Q -sector,

$$\hat{P}_b f(Q) = f(Q) \hat{P}_b - i\hbar (\partial_{Q_b} f) \quad \implies \quad \hat{P}_b \mu_{\alpha\beta} = \mu_{\alpha\beta} \hat{P}_b - i\hbar \partial_b \mu_{\alpha\beta}, \quad \hat{P}_b \sigma_{\alpha c} = \sigma_{\alpha c} \hat{P}_b - i\hbar \zeta_{bc}^\alpha, \quad (\text{C.3})$$

where the last relation uses $\sigma_{\alpha b} = \sum_a \zeta_{ab}^\alpha Q_a$ [(3.17a)], linear in Q with *constant* $\zeta_{ab}^\alpha = (\sum_j \mathbf{l}_{ja} \times \mathbf{l}_{jb})_\alpha$, so $\partial_{Q_b} \sigma_{\alpha c} = \zeta_{bc}^\alpha$. The antisymmetry of the cross product gives the sum rule that recurs throughout:

$$\zeta_{ab}^\alpha = -\zeta_{ba}^\alpha \quad \implies \quad \zeta_{bb}^\alpha = 0, \quad \sum_b \partial_{Q_b} \sigma_{\alpha b} = \sum_b \zeta_{bb}^\alpha = 0. \quad (\text{C.4})$$

$\hat{\pi}_\alpha$ is Hermitian on the flat mode measure. With $\hat{\pi}_\alpha = \sum_b \sigma_{\alpha b} \hat{P}_b$, take the adjoint (order reverses; σ real, $\hat{P}_b$ Hermitian), then slide $\hat{P}_b$ back to the right with (C.3), then apply (C.4):

$$\hat{\pi}_\alpha^\dagger = \sum_b \hat{P}_b \sigma_{\alpha b} = \sum_b (\sigma_{\alpha b} \hat{P}_b - i\hbar \zeta_{bb}^\alpha) = \hat{\pi}_\alpha. \quad (\text{C.5})$$

Had $\sum_b \zeta_{bb}^\alpha$ been nonzero, $\hat{\pi}_\alpha$ would have required explicit symmetrization; the antisymmetry of ζ spares us. Finally, $\mu_{\alpha\beta}(Q) = \mu_{\beta\alpha}(Q)$ is a real symmetric function of Q alone.

C.2 The nine products

Substitute $\hat{A}_\alpha = \hat{J}_\alpha - \hat{L}_\alpha - \hat{\pi}_\alpha$ into (C.1) and multiply out, *keeping the operator order* (the factors need not commute):

$$\begin{aligned} \hat{A}_\alpha \mu_{\alpha\beta} \hat{A}_\beta = & \underbrace{\hat{J}_\alpha \mu_{\alpha\beta} \hat{J}_\beta}_{\text{(T1) rotation}} + \underbrace{\hat{L}_\alpha \mu_{\alpha\beta} \hat{L}_\beta}_{\text{(T2) electronic}} + \underbrace{\hat{\pi}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta}_{\text{(T3) vibrational}} - \underbrace{(\hat{J}_\alpha \mu_{\alpha\beta} \hat{L}_\beta + \hat{L}_\alpha \mu_{\alpha\beta} \hat{J}_\beta)}_{\text{(T4) rotation-electron}} \\ & - \underbrace{(\hat{J}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta + \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{J}_\beta)}_{\text{(T5) rotation-vibration}} + \underbrace{(\hat{L}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta + \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{L}_\beta)}_{\text{(T6) vibration-electron}}. \end{aligned} \quad (\text{C.6})$$

Each cross term collects the two equal classical contributions $X\mu Y$ and $Y\mu X$, which is why it appears as a symmetric sum of orderings. We now treat the six blocks one at a time: for each we verify Hermiticity and write the explicit differential form.

C.3 Diagonal terms

(T1) $\hat{J}_\alpha \mu_{\alpha\beta} \hat{J}_\beta$. *Hermiticity.* Take the adjoint (μ real, $\hat{J}_\alpha^\dagger = \hat{J}_\alpha$ on $\sin\theta d\Omega$; order reverses), then relabel the summed dummy indices $\alpha \leftrightarrow \beta$, then use $\mu_{\beta\alpha} = \mu_{\alpha\beta}$:

$$(\hat{J}_\alpha \mu_{\alpha\beta} \hat{J}_\beta)^\dagger = \hat{J}_\beta \mu_{\alpha\beta} \hat{J}_\alpha = \hat{J}_\alpha \mu_{\beta\alpha} \hat{J}_\beta = \hat{J}_\alpha \mu_{\alpha\beta} \hat{J}_\beta. \quad (\text{C.7})$$

Hermitian exactly as written — no symmetrization, no commutator evaluated. *Differential form.* $\mu(Q)$ commutes with the Euler derivatives, and inserting (C.2),

$$\hat{J}_\alpha \mu_{\alpha\beta} \hat{J}_\beta = -\hbar^2 \mu_{\alpha\beta} (\mathbf{E}^{-1})_{s\alpha} \left[(\mathbf{E}^{-1})_{t\beta} \partial_{\Omega_s} \partial_{\Omega_t} + (\partial_{\Omega_s} (\mathbf{E}^{-1})_{t\beta}) \partial_{\Omega_t} \right], \quad (\text{C.8})$$

with the trigonometric entries of $\mathbf{E}^{-1}$ read off (4.12). Every term carries at least one Euler derivative: the rotational block generates *no* pure multiplicative term, in agreement with the Laplace–Beltrami computation (4.29).

(T2) $\hat{L}_\alpha \mu_{\alpha\beta} \hat{L}_\beta$. Word-for-word the same as (T1) with $\hat{J} \rightarrow \hat{L}$: the adjoint chain (C.7) uses only $\hat{L}_\alpha^\dagger = \hat{L}_\alpha$ [(4.18)] and the symmetry of μ . Hermitian as written; all terms carry electron derivatives; no multiplicative residue.

(T3) $\hat{\pi}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta$. *Hermiticity.* Use $\hat{\pi}^\dagger = \hat{\pi}$ from (C.5), then relabel $\alpha \leftrightarrow \beta$ and use μ symmetric — the same three-step chain as (C.7). Hermitian as written, but *only* because $\hat{\pi}$ is Hermitian, i.e. ultimately because of the antisymmetry (C.4). *Differential form.* Write $\hat{\pi}_\alpha = \sigma_{\alpha b} \hat{P}_b$ and slide the left momentum past the Q -function $\mu_{\alpha\beta} \sigma_{\beta c}$ with (C.3):

$$\begin{aligned} \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta &= \sigma_{\alpha b} \hat{P}_b (\mu_{\alpha\beta} \sigma_{\beta c}) \hat{P}_c = \sigma_{\alpha b} \left[(\mu_{\alpha\beta} \sigma_{\beta c}) \hat{P}_b - i\hbar \partial_b (\mu_{\alpha\beta} \sigma_{\beta c}) \right] \hat{P}_c \\ &= \underbrace{G_{bc} \hat{P}_b \hat{P}_c}_{\text{2nd order}} - i\hbar \underbrace{\sigma_{\alpha b} \left[(\partial_b \mu_{\alpha\beta}) \sigma_{\beta c} + \mu_{\alpha\beta} \zeta_{bc}^\beta \right]}_{\text{1st order}} \hat{P}_c, \quad G_{bc} \equiv (\boldsymbol{\sigma}^\top \boldsymbol{\mu} \boldsymbol{\sigma})_{bc}, \end{aligned} \quad (\text{C.9})$$

the Coriolis-dressed vibrational operator. Every surviving term carries at least one $\hat{P}$: again no pure multiplicative term. *Equivalence with the ordered form.* The same sum rule shows that (T3) equals the fully left-ordered operator: moving the left $\hat{P}_b$ past its coefficient

σ_{ab} ,

$$\hat{P}_b G_{bc} \hat{P}_c - \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta = [\hat{P}_b, \sigma_{ab}] \mu_{\alpha\beta} \sigma_{\beta c} \hat{P}_c = -i\hbar \left(\sum_b \zeta_{bb}^\alpha \right) \mu_{\alpha\beta} \sigma_{\beta c} \hat{P}_c = 0, \quad (\text{C.10})$$

so $\frac{1}{2} \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta = \frac{1}{2} \hat{P}_b G_{bc} \hat{P}_c$ *exactly*. This identity is what lets the sandwich-ordered kernel of (4.41) match the Laplace–Beltrami vibrational block (4.31), whose dressed part is precisely $\frac{1}{2} \hat{P}_b G_{bc} \hat{P}_c$ after the rescaling (see § C.5).

C.4 Cross terms

(T4) $\hat{J}_\alpha \mu_{\alpha\beta} \hat{L}_\beta + \hat{L}_\alpha \mu_{\alpha\beta} \hat{J}_\beta$. All three factors act on disjoint variables and mutually commute. A single ordering is already Hermitian,

$$(\hat{J}_\alpha \mu_{\alpha\beta} \hat{L}_\beta)^\dagger = \hat{L}_\beta \mu_{\alpha\beta} \hat{J}_\alpha = \mu_{\alpha\beta} \hat{J}_\alpha \hat{L}_\beta = \hat{J}_\alpha \mu_{\alpha\beta} \hat{L}_\beta, \quad (\text{C.11})$$

and the two orderings in the sum are equal (relabel $\alpha \leftrightarrow \beta$ and commute), so

$$\hat{J}_\alpha \mu_{\alpha\beta} \hat{L}_\beta + \hat{L}_\alpha \mu_{\alpha\beta} \hat{J}_\beta = 2 \mu_{\alpha\beta} \hat{J}_\alpha \hat{L}_\beta \quad (\text{Hermitian; no residue}). \quad (\text{C.12})$$

This is the rotation–electron Coriolis coupling; no $\hat{P}$ appears, so there is no differential subtlety.

(T5) $\hat{J}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta + \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{J}_\beta$. Now the subtlety: $\hat{\pi}$ does *not* commute with μ , since by (C.3) $\hat{\pi}_\beta \mu_{\alpha\beta} = \mu_{\alpha\beta} \hat{\pi}_\beta - i\hbar \sigma_{\beta b} \partial_b \mu_{\alpha\beta} \neq \mu_{\alpha\beta} \hat{\pi}_\beta$. A single ordering is therefore *not* Hermitian by itself; its adjoint is exactly the *other* ordering ($\hat{J}$ commutes with μ and $\hat{\pi}$; relabel $\alpha \leftrightarrow \beta$):

$$(\hat{J}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta)^\dagger = \hat{\pi}_\beta \mu_{\alpha\beta} \hat{J}_\alpha = \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{J}_\beta. \quad (\text{C.13})$$

The cross term is of the form $\hat{X} + \hat{X}^\dagger$, automatically Hermitian — *this is why the classical product $-2J\mu\pi$ must be quantized as the symmetric sum*, and it is precisely the symmetric sum that the Laplace–Beltrami operator produced by itself in (4.30)/(4.35). Pulling $\hat{J}$ out front and expanding the anticommutator with (C.3),

$$\hat{J}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta + \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{J}_\beta = \hat{J}_\alpha (\mu_{\alpha\beta} \hat{\pi}_\beta + \hat{\pi}_\beta \mu_{\alpha\beta}) = \hat{J}_\alpha \left(\underbrace{2 \mu_{\alpha\beta} \sigma_{\beta c} \hat{P}_c}_{\text{1st order in } \hat{P}} - \underbrace{i\hbar \sigma_{\beta c} \partial_c \mu_{\alpha\beta}}_{\text{0th order in } \hat{P}} \right). \quad (\text{C.14})$$

The first piece is the rotation–vibration Coriolis coupling ($\hat{J}$ times $\hat{P}$); the second is the imaginary, $\hat{J}$ -linear counterterm that makes the sum Hermitian. It is *not* a stand-alone multiplicative residue — it still carries $\hat{J}_\alpha$ — consistent with the Laplace–Beltrami result (4.35) that the Coriolis block contributes nothing to the pseudopotential. As a cross-check of (C.14) against the pre-rescaling form (4.30): expanding the second group of (4.30) by the product rule,

$$\frac{1}{\sqrt{\gamma}} \hat{P}_b \sqrt{\gamma} \sigma_{ab} \mu_{\alpha\beta} \hat{J}_\beta = \left[\sigma_{ab} \mu_{\alpha\beta} \hat{P}_b - i\hbar \partial_b (\sigma_{ab} \mu_{\alpha\beta}) - \frac{i\hbar}{2} (\partial_b \ln \gamma) \sigma_{ab} \mu_{\alpha\beta} \right] \hat{J}_\beta, \quad (\text{C.15})$$

and conjugating by $\gamma^{1/4}$ as in § 4.9 cancels the $\partial_b \ln \gamma$ terms between the two groups, reproducing exactly $-\frac{1}{2} \hat{J}_\alpha \{\mu_{\alpha\beta}, \hat{\pi}_\beta\}$ of (4.35), i.e. $-\frac{1}{2} \times (\text{C.14})$.

(T6) $\hat{L}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta + \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{L}_\beta$. Identical to (T5) with $\hat{J} \rightarrow \hat{L}$ ($\hat{L}$ also commutes with μ and $\hat{\pi}$):

$$\hat{L}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta + \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{L}_\beta = \hat{L}_\alpha \left(2 \mu_{\alpha\beta} \sigma_{\beta c} \hat{P}_c - i \hbar \sigma_{\beta c} \partial_c \mu_{\alpha\beta} \right), \quad (\text{C.16})$$

the vibration–electron angular coupling.

C.5 The measure residue U_{Cor} , every derivative

Section 4.9 derived the identity-block residue U_{vib} [(4.34)] in full and deferred the residue of the dressed block. Here it is. Conjugate the dressed part of the vibrational Laplace–Beltrami block (4.31) by $\gamma^{1/4}$ and act on ψ_W ; each line is one product rule [$G_{bc} = (\boldsymbol{\sigma}^\top \boldsymbol{\mu} \boldsymbol{\sigma})_{bc}$, symmetric]:

$$\begin{aligned} \partial_{Q_c} (\gamma^{-1/4} \psi_W) &= \gamma^{-1/4} \left[\partial_{Q_c} \psi_W - \frac{1}{4} (\partial_c \ln \gamma) \psi_W \right], \\ \gamma^{1/2} G_{bc} \partial_{Q_c} (\gamma^{-1/4} \psi_W) &= \gamma^{1/4} G_{bc} \left[\partial_{Q_c} \psi_W - \frac{1}{4} (\partial_c \ln \gamma) \psi_W \right], \\ \partial_{Q_b} \left(\gamma^{1/4} G_{bc} [\dots] \right) &= \gamma^{1/4} \left[G_{bc} \partial_b \partial_c \psi_W + (\partial_b G_{bc}) \partial_c \psi_W + \frac{1}{4} (\partial_b \ln \gamma) G_{bc} \partial_c \psi_W - \frac{1}{4} G_{bc} (\partial_c \ln \gamma) \partial_b \psi_W \right. \\ &\quad \left. - \frac{1}{4} \partial_b (G_{bc} \partial_c \ln \gamma) \psi_W - \frac{1}{16} G_{bc} (\partial_b \ln \gamma) (\partial_c \ln \gamma) \psi_W \right]. \end{aligned} \quad (\text{C.17})$$

The two first-order $\ln \gamma$ terms cancel by the symmetry of G_{bc} (relabel $b \leftrightarrow c$), leaving $G \partial \partial + (\partial G) \partial$ — which is exactly $-\frac{2}{\hbar^2} \times \frac{1}{2} \hat{P}_b G_{bc} \hat{P}_c$ — plus a pure multiplicative remainder. Multiplying by $-\frac{\hbar^2}{2}$ (the prefactors $\gamma^{1/4} \cdot \gamma^{-1/2} \cdot \gamma^{1/4} = 1$ combine to unity),

$$\gamma^{1/4} \hat{T}_{\text{dress}} \gamma^{-1/4} = \frac{1}{2} \hat{P}_b G_{bc} \hat{P}_c + U_{\text{Cor}}, \quad \boxed{U_{\text{Cor}} = \frac{\hbar^2}{8} \left[\partial_b (G_{bc} \partial_c \ln \gamma) + \frac{1}{4} G_{bc} (\partial_b \ln \gamma) (\partial_c \ln \gamma) \right]}. \quad (\text{C.18})$$

By (C.10), $\frac{1}{2} \hat{P}_b G_{bc} \hat{P}_c = \frac{1}{2} \hat{\pi}_\alpha \mu_{\alpha\beta} \hat{\pi}_\beta$, so the operator part is exactly the (T3) sandwich of the kernel (4.41), and the entire ordering debt of the dressed block is the multiplicative U_{Cor} . Together with U_{vib} of (4.34) [the same computation with $G_{bc} \rightarrow \delta_{bc}$], the total residue is

$$\hat{U} = U_{\text{vib}} + U_{\text{Cor}} = \frac{\hbar^2}{8} \left[\partial_b (g_{bc}^{QQ} \partial_c \ln \gamma) + \frac{1}{4} g_{bc}^{QQ} (\partial_b \ln \gamma) (\partial_c \ln \gamma) \right], \quad g_{bc}^{QQ} = \delta_{bc} + G_{bc}, \quad (\text{C.19})$$

every ingredient of which reduces, via Jacobi’s formula (4.36) and the linearity (4.37), to the constant $\boldsymbol{\zeta}$ ’s and the linear $\boldsymbol{\sigma}(Q)$; the Watson sum rules then collapse (C.19) to the trace $\hat{U} = -\frac{\hbar^2}{8} \sum_\alpha \mu_{\alpha\alpha}$ of (4.38) [3].

The division of labour is clean. In the explicit forms (C.8), (C.9), (C.12), (C.14), (C.16), every surviving kernel term carries at least one derivative operator ($\hat{P}$, $\hat{J}$, or $\hat{L}$): *the ordering of the kernel never generates a pure multiplicative term*. The pseudopotential (4.38) is an $\mathcal{O}(\hbar^2)$ c -number that originates entirely in the *curved integration measure* $\sqrt{\gamma}$, extracted by the rescaling $\psi_W = \gamma^{1/4} \psi$ — (C.18) is its dressed-block share. Kernel ordering $\rightarrow$ the Hermitian operators of this appendix; the measure $\rightarrow$ the pseudopotential.

C.6 Worked example: a bent triatomic

To make “every derivative” concrete, take a bent triatomic XY_2 (e.g. H_2O), a C_{2v} asymmetric top with $N_{\text{nucl}} = 3$ and $f = 3N_{\text{nucl}} - 6 = 3$ normal coordinates: the symmetric stretch

Q_1 and bend Q_2 (both a_1) and the antisymmetric stretch Q_3 (b_2). Choose the body frame as the instantaneous principal-axis system, with a, b in the molecular plane and $c \perp$ to it.

Inertia and its derivatives. At equilibrium the inertia tensor is diagonal, $\mathbf{I}_N^{\text{eq}} = \text{diag}(I_a^e, I_b^e, I_c^e)$, and because the equilibrium mass distribution is planar the perpendicular-axis theorem gives the exact relation

$$I_c^e = I_a^e + I_b^e. \quad (\text{C.20})$$

The totally symmetric coordinates Q_1, Q_2 preserve the C_{2v} symmetry, so to linear order they only *scale* the diagonal moments, whereas the antisymmetric Q_3 tilts the principal axes and enters $\mathbf{I}_N$ only at second order. Hence

$$I_\alpha(Q) = I_\alpha^e + a_\alpha^1 Q_1 + a_\alpha^2 Q_2 + \mathcal{O}(Q^2), \quad a_\alpha^b \equiv \left. \frac{\partial I_\alpha}{\partial Q_b} \right|_{\text{eq}}, \quad (\text{C.21})$$

with the inertial derivatives a_α^b computable from the mode vectors $\mathbf{l}_{jb}$ evaluated at equilibrium. These are exactly the derivatives that populate the rotation–vibration Jacobian $\sqrt{\gamma(Q)}$ of (4.21).

Coriolis coupling. The only nonzero constant Coriolis matrix in C_{2v} couples the two a_1 modes to the b_2 mode about the b axis (the axis whose rotation maps a symmetric into an antisymmetric displacement):

$$\boldsymbol{\sigma}_1 = \zeta_{13}^b Q_3 \hat{\mathbf{b}}, \quad \boldsymbol{\sigma}_2 = \zeta_{23}^b Q_3 \hat{\mathbf{b}}, \quad \boldsymbol{\sigma}_3 = -(\zeta_{13}^b Q_1 + \zeta_{23}^b Q_2) \hat{\mathbf{b}}, \quad \zeta_{ab}^b = -\zeta_{ba}^b, \quad (\text{C.22})$$

so $\boldsymbol{\sigma}\boldsymbol{\sigma}^T = [(\zeta_{13}^b Q_3)^2 + (\zeta_{23}^b Q_3)^2 + (\zeta_{13}^b Q_1 + \zeta_{23}^b Q_2)^2] \hat{\mathbf{b}}\hat{\mathbf{b}}^T$ corrects only the bb element of the inertia,

$$\mathbf{I}'(Q) = \text{diag}(I_a(Q), I_b(Q) - s(Q), I_c(Q)), \quad s(Q) \equiv (\zeta_{13}^b Q_3)^2 + (\zeta_{23}^b Q_3)^2 + (\zeta_{13}^b Q_1 + \zeta_{23}^b Q_2)^2. \quad (\text{C.23})$$

Note $s(Q) = \mathcal{O}(Q^2)$, so it does not affect the linear-order derivatives (C.21) but does enter the pseudopotential and the exact Jacobian.

Determinant, volume element, and its explicit derivatives. With $\mathbf{I}'$ diagonal in (C.23),

$$\gamma(Q) = \det \mathbf{I}' = I_a(Q) (I_b(Q) - s(Q)) I_c(Q), \quad \sqrt{g} = \sin \theta \sqrt{I_a (I_b - s) I_c}, \quad (\text{C.24})$$

and the vibrational Jacobian carries the explicit derivatives

$$\partial_{Q_b} \ln \gamma = \frac{a_a^b}{I_a} + \frac{a_b^b - \partial_{Q_b} s}{I_b - s} + \frac{a_c^b}{I_c}, \quad \partial_{Q_3} s = 2[(\zeta_{13}^b)^2 + (\zeta_{23}^b)^2] Q_3, \quad \partial_{Q_1} s = 2\zeta_{13}^b (\zeta_{13}^b Q_1 + \zeta_{23}^b Q_2), \text{ etc.}, \quad (\text{C.25})$$

which are precisely the first-order remainders exhibited under (4.31) and removed by the $\gamma^{1/4}$ rescaling of § 4.9.

Pseudopotential. Since $\mathbf{I}'$ is diagonal, $\boldsymbol{\mu} = \text{diag}(1/I_a, 1/(I_b - s), 1/I_c)$ and the trace formula (4.38) is fully explicit:

$$\hat{U}(Q) = -\frac{\hbar^2}{8} \left[\frac{1}{I_a(Q)} + \frac{1}{I_b(Q) - s(Q)} + \frac{1}{I_c(Q)} \right]. \quad (\text{C.26})$$

Expanding to the order at which it first varies with the coordinates, and using (C.20),

$$\hat{U}(Q) = -\frac{\hbar^2}{8} \left(\frac{1}{I_a^e} + \frac{1}{I_b^e} + \frac{1}{I_c^e} \right) + \frac{\hbar^2}{8} \sum_{b=1,2} \left(\frac{a_a^b}{(I_a^e)^2} + \frac{a_b^b}{(I_b^e)^2} + \frac{a_c^b}{(I_c^e)^2} \right) Q_b + \mathcal{O}(Q^2). \quad (\text{C.27})$$

The constant is an inconsequential shift of the electronic–vibrational zero; the linear term is a genuine $\hbar^2$ force on the normal coordinates — the physical content of the Watson pseudopotential for this molecule — produced entirely by differentiating $\sqrt{\gamma(Q)}$ through (C.25). Every step from the metric (4.15) to the surface (C.27) used only $\partial_\phi, \partial_\theta, \partial_\chi$ and ∂_{Q_b} .

C.7 Summary

1. The quantized kernel (C.1) expands into the nine products (C.6). Blocks built from mutually commuting, individually Hermitian factors — (T1) $\hat{J}\mu\hat{J}$, (T2) $\hat{L}\mu\hat{L}$, (T4) $\hat{J}\mu\hat{L}$ — are Hermitian in a single ordering, by the adjoint-reversal argument (C.7) alone.
2. Blocks pairing $\hat{\pi}$ with $\hat{J}$ or $\hat{L}$ — (T5), (T6) — are Hermitian only as the symmetric sum $\hat{X} + \hat{X}^\dagger$, which is exactly the ordering the Laplace–Beltrami operator produces by itself [(4.30), (4.35)]. (T3) $\hat{\pi}\mu\hat{\pi}$ is the fortunate middle case, Hermitian because $\hat{\pi}$ is [(C.5)], and equal to the left-ordered $\frac{1}{2}\hat{P}G\hat{P}$ by the sum rule (C.10).
3. No kernel ordering produces a pure multiplicative term. The Watson pseudopotential comes solely from the measure: U_{vib} [(4.34)] plus U_{Cor} [(C.18)] collapse to $-\frac{\hbar^2}{8} \sum_\alpha \mu_{\alpha\alpha}$ [(4.38)].
4. Every Hermiticity statement above used only integration by parts, the product rule, and the divergence identities (4.18), (4.26), (C.4) — never a commutator of body-fixed angular momenta.

References

- [1] C. Eckart, “Some studies concerning rotating axes and polyatomic molecules,” *Phys. Rev.*, vol. 47, pp. 552–558, 1935.
- [2] A. Sayvetz, “The kinetic energy of polyatomic molecules,” *J. Chem. Phys.*, vol. 7, pp. 383–389, 1939.
- [3] J. K. G. Watson, “Simplification of the molecular vibration-rotation Hamiltonian,” *Mol. Phys.*, vol. 15, pp. 479–490, 1968.
- [4] P. R. Bunker and P. Jensen, *Molecular Symmetry and Spectroscopy*, 2nd ed. Ottawa: NRC Research Press, 1998.

- [5] E. B. Wilson, J. C. Decius, and P. C. Cross, *Molecular Vibrations: The Theory of Infrared and Raman Vibrational Spectra*. New York: McGraw-Hill, 1955.
- [6] R. S. Mulliken and A. Christy, “ Λ -type doubling and electron configurations in diatomic molecules,” *Phys. Rev.*, vol. 38, pp. 87–119, 1931.
- [7] J. H. Van Vleck, “The coupling of angular momentum vectors in molecules,” *Rev. Mod. Phys.*, vol. 23, pp. 213–227, 1951.
- [8] R. Renner, “Zur Theorie der Wechselwirkung zwischen Elektronen- und Kernbewegung bei dreiatomigen, stabförmigen Molekülen,” *Z. Phys.*, vol. 92, pp. 172–193, 1934.
- [9] R. F. Curl, “The relationship between electron spin rotation coupling constants and g -tensor components,” *Mol. Phys.*, vol. 9, pp. 585–597, 1965.
- [10] M. Born and R. Oppenheimer, “Zur Quantentheorie der Molekeln,” *Ann. Phys.*, vol. 389, pp. 457–484, 1927.
- [11] M. Born and K. Huang, *Dynamical Theory of Crystal Lattices*. Oxford: Oxford University Press, 1954.
- [12] R. G. Littlejohn and M. Reinsch, “Gauge fields in the separation of rotations and internal motions in the n -body problem,” *Rev. Mod. Phys.*, vol. 69, pp. 213–275, 1997.
- [13] B. Podolsky, “Quantum-mechanically correct form of Hamiltonian function for conservative systems,” *Phys. Rev.*, vol. 32, pp. 812–816, 1928.
- [14] O. Klein, “Zur Frage der Quantelung des asymmetrischen Kreisels,” *Z. Phys.*, vol. 58, pp. 730–734, 1929.