

# Quantum Physics II Spring 2019

## Midterm Exam

Instructor: Pai-hsien Jennifer Hsu

Tuesday April 16, 2019

You must show your work. No credits will be given if you don't show how you get your answers.

You may use the following formula:

- One can determine how a state  $|\psi\rangle$  evolves with time

$$\psi(x', t') = \int_{-\infty}^{\infty} dx_0 \langle x', t' | x_0, t_0 \rangle \psi(x_0, t_0)$$

if one knows the amplitude  $\langle x', t' | x_0, t_0 \rangle$ :

$$\langle x', t' | x_0, t_0 \rangle = \int_{x_0}^{x'} D[x(t)] \exp\left(i \frac{1}{\hbar} S[x(t)]\right),$$

where  $D[x(t)]$  means summing over all possible path  $x(t)$  between  $(x_0, t_0)$  and  $(x', t')$ .  $S[x(t)]$  is the action for a path  $x(t)$ :

$$S[x(t)] = \int_{t_0}^{t'} dt L(x, \dot{x}),$$

and  $L(x, \dot{x})$  being the Lagrangian,

$$L(x, \dot{x}) = \frac{1}{2} m \dot{x}^2 - V(x).$$

The Planck constant  $\hbar = 1.055 \times 10^{-34}$  Joule  $\cdot$  sec.

- The Schrödinger equation:

$$i\hbar \frac{d|\psi(t)\rangle}{dt} = \hat{H} |\psi(t)\rangle$$

where  $\hat{H}$  is the Hamiltonian  $\frac{\hat{p}^2}{2\mu} + V(\hat{r})$  and  $\mu$  is the mass of the particle.

For energy eigenstates, this reduces to the time-independent Schrödinger equation (in spherical coordinates)

$$-\frac{\hbar^2}{2\mu}\left(\frac{\partial^2}{\partial r^2} + \frac{2}{r}\frac{\partial}{\partial r}\right)\psi(r) + \langle r | \frac{\hat{\mathbf{L}}^2}{2\mu r^2} | \psi \rangle + V(r)\psi(r) = E\psi(r)$$

$\hat{\mathbf{L}}$  is the orbital angular momentum operator,  $\hat{\mathbf{L}} = \hat{\mathbf{r}} \times \hat{\mathbf{p}}$ , which satisfies the usual commutation relations for angular momenta, e.g.  $[\hat{L}_x, \hat{L}_y] = i\hbar\hat{L}_z$ . In the spherical coordinate,  $\hat{L}_z \rightarrow -i\hbar\frac{\partial}{\partial\phi}$ .

- For spherical symmetric potential  $V(r)$ , there exist simultaneous eigenstates of  $\hat{H}$ ,  $\hat{\mathbf{L}}^2$ , and  $\hat{L}_z$ :

$$\hat{H} |E, l, m\rangle = E |E, l, m\rangle, \hat{\mathbf{L}}^2 |E, l, m\rangle = l(l+1)\hbar^2 |E, l, m\rangle, \hat{L}_z |E, l, m\rangle = m\hbar |E, l, m\rangle.$$

- In terms of wavefunctions,  $|E, l, m\rangle$  is of the form  $R(r)Y_{l,m}(\theta, \phi)$ , where the spherical harmonics  $Y_{l,m}(\theta, \phi) = \langle \theta, \phi | l, m \rangle$ , and

$$\int d\Omega |Y_{l,m}(\theta, \phi)|^2 = 1,$$

with  $d\Omega$  being the solid angle.

- Given one  $|l, m\rangle$ , one can obtain other eigenkets with the same  $l$  by the ladder operators  $\hat{L}_{\pm} = \hat{L}_x \pm i\hat{L}_y$ :

$$\hat{L}_{\pm} |l, m\rangle = \sqrt{l(l+1) - m(m \pm 1)}\hbar |l, m \pm 1\rangle.$$

- The parity operator  $\hat{\pi}$  acts on a ket  $|\psi\rangle$  and flips the sign of the position expectation value:

$$|\psi'\rangle = \hat{\pi} |\psi\rangle, \langle x \rangle_{\psi'} = -\langle x \rangle_{\psi}$$

From this definition one can show that  $\hat{\pi}\hat{x}\hat{\pi} = -\hat{x}$ ,  $\hat{\pi}|x\rangle = |-x\rangle$ , and  $\hat{\pi} = \hat{\pi}^{-1} = \hat{\pi}^\dagger$ . The eigenvalues of  $\hat{\pi}$  are  $\pm 1$ . The wavefunctions of its eigenkets have the property  $\psi(-x) = \pm\psi(x)$ , and thus they are called parity-even or parity-odd states.

- The energy eigenvalues of a hydrogen-like atom (with Coulomb potential  $-\frac{Ze^2}{r}$ ) can be written as  $E_n = -\frac{\mu c^2 Z^2 \alpha^2}{2n^2}$ , where  $\mu$  is the reduced mass (can be take as the mass of an electron  $m_e$  for most cases), and  $\alpha$  is the fine structure constant  $\alpha = e^2/\hbar c \approx 1/137$ .

- For a 1-D simple harmonic oscillator (SHO),  $\hat{H} = \frac{\hat{p}_x^2}{2m} + \frac{1}{2}m\omega^2\hat{x}^2$ :

The raising and lowering operators are

$$\hat{a}^\dagger = \sqrt{\frac{m\omega}{2\hbar}}(\hat{x} - \frac{i}{m\omega}\hat{p}_x), \quad \hat{a} = \sqrt{\frac{m\omega}{2\hbar}}(\hat{x} + \frac{i}{m\omega}\hat{p}_x)$$

and  $[\hat{a}, \hat{a}^\dagger] = 1$ . The operators get their names from the facts that

$$\hat{a}^\dagger |n\rangle = \sqrt{n+1} |n+1\rangle, \quad \hat{a} |n\rangle = \sqrt{n} |n-1\rangle.$$

One can rewrite  $\hat{x}$  and  $\hat{p}_x$  as

$$\hat{x} = \sqrt{\frac{\hbar}{2m\omega}}(\hat{a} + \hat{a}^\dagger), \quad \hat{p}_x = -i\sqrt{\frac{m\omega\hbar}{2}}(\hat{a} - \hat{a}^\dagger)$$

and the Hamiltonian as

$$\hat{H} = \hbar\omega(\hat{a}^\dagger\hat{a} + \frac{1}{2}) = \hbar\omega(\hat{N} + \frac{1}{2}),$$

where the number operator  $\hat{N} \equiv \hat{a}^\dagger\hat{a}$ .

- The energy eigenvalues of an SHO are  $E_n = (n + \frac{1}{2})\hbar\omega$ ,  $n = 0, 1, 2, \dots$
- Expectation value of an operator  $\hat{A}$  for a state  $|\psi\rangle$  is  $\langle A \rangle = \langle \psi | \hat{A} | \psi \rangle$
- Uncertainty of an operator  $\hat{A}$  is defined as  $\Delta A = \sqrt{\langle A^2 \rangle - \langle A \rangle^2}$
- Heisenberg Equation:

$$\frac{d}{dt}\langle A \rangle = \langle \frac{\partial A}{\partial t} \rangle + \frac{1}{i\hbar}\langle [\hat{A}, \hat{H}] \rangle$$

where  $\hat{H}$  is the Hamiltonian.

1. Quantum v.s. Classical descriptions:

- (a) Use the Heisenberg equation prove that (in 1-D)

$$\frac{d}{dt}\langle xp \rangle = 2\langle T \rangle - \langle x \frac{dV}{dx} \rangle,$$

where  $\hat{T} = \hat{H} - \hat{V}$  is the kinetic energy. (2 points)

[Hint:  $[\hat{A}\hat{B}, \hat{C}] = \hat{A}[\hat{B}, \hat{C}] + [\hat{A}, \hat{C}]\hat{B}$ , and  $[\hat{p}_x, f(x)] = -i\hbar \frac{df}{dx}$ .]

- (b) Explain why the result reduces to

$$2\langle T \rangle = \langle x \frac{dV}{dx} \rangle$$

for energy eigenstates. This is the Virial Theorem you've seen in Classical Mechanics. (2 points)

- (c) Express the Virial Theorem in 3-D, and show that  $\langle T \rangle = -\frac{1}{2}\langle V \rangle$  for a hydrogen atom (Coulomb potential). (2 points)

- (d) Argue that the motion of a ground-state electron in a hydrogen atom has to be described *quantum-mechanically*. (2 points)

[Hint: Use the results from part (c) to estimate the size of  $v/c$  ( $v$  is the typical speed) for the electron, and take the typical size of a hydrogen atom as the traveling distance, and estimate the size of its classical action.  $m_e \approx 9 \times 10^{-31}$  kg. An order-of-magnitude estimation is all you need.]

2. Electron in a Coulomb potential: Let  $\psi_{nlm}(\mathbf{r})$  denote the normalized energy eigenfunctions of a Coulomb potential, with principle quantum number  $n$  and angular momentum quantum numbers  $l$  and  $m$ . Consider an electron in the state

$$\Psi(\mathbf{r}) = c[\psi_{100}(\mathbf{r}) + 4i\psi_{210}(\mathbf{r}) - 2\sqrt{2}\psi_{21-1}(\mathbf{r})]$$

at time  $t = 0$ .

- (a) Find the normalization constant  $c$ . (1 point)
- (b) Let  $E_1$  be the energy eigenvalue of the state  $\psi_{100}(\mathbf{r})$ , express the expectation value of energy  $\langle E \rangle$  at  $t = 0$  in terms of  $E_1$ . (1 point)
- (c) Evaluate the expectation values of the angular momentum  $\langle L_z \rangle$  and  $\langle L_x \rangle$  at time  $t = 0$ . (2 points)

[Hint: Make use of the ladder operators  $\hat{L}_{\pm}$ ]

- (d) Now we turn on an external magnetic field along the z-axis, and the resulting Hamiltonian is

$$\hat{H} = \frac{\hat{\mathbf{p}}^2}{2\mu} - \frac{Ze^2}{r} + \omega_0 \hat{L}_z.$$

Evaluate  $\langle L_z \rangle$  and  $\langle L_x \rangle$  as functions of time. (2 points).

[Note: You can save some efforts by arguing which is/are constant(s) of motion.]

### 3. Parity of SHO eigenstates:

- (a) Consider a 1-D SHO, explain why its energy eigenstates have definite parity (i.e. each energy eigenstate  $|n\rangle$  is an eigenstate of the parity operator  $\hat{\pi}$ ). (1 point)
- (b) Show that  $\hat{\pi} \hat{a}^\dagger \hat{\pi} = -\hat{a}^\dagger$ , where  $\hat{a}^\dagger$  is the raising operator of the 1-D SHO. (1 point)
- (c) Show that the wavefunction of the ground state  $|0\rangle$  for a 1-D SHO is Gaussian (you don't need to derive the normalization constant). (2 points)  
[Hint: Use the property  $\hat{a}|0\rangle = 0$  ( $\hat{a}$  is the lowering operator). Note that  $\hat{p}_x \rightarrow -i\hbar \frac{d}{dx}$  in 1-D.]
- (d) Given the fact  $\langle x|0\rangle$  is Gaussian and the results from part (b), show that  $\hat{\pi}|n\rangle = (-1)^n |n\rangle$ . (2 points)

### 4. Schrödinger equation in the presence of EM field: We've shown in class that the canonical momentum (to be identified with $-i\hbar\nabla$ in the coordinate space) when there exists a magnetic vector potential $\mathbf{A}$ is

$$\mathbf{p}_c = m\mathbf{v} + \frac{q}{c}\mathbf{A}$$

And the Hamiltonian is

$$H = \frac{(\mathbf{p}_c - q\mathbf{A}/c)^2}{2m} + q\phi$$

where  $\phi$  is the electric potential. Different  $\mathbf{A}$  and  $\phi$  can result in the same  $\mathbf{E}$  and  $\mathbf{B}$  fields, under the following gauge transformation:

$$\mathbf{A}' = \mathbf{A} - \nabla f(\mathbf{r}, t), \quad \phi' = \phi + \frac{1}{c} \frac{\partial f}{\partial t}.$$

with some real function  $f(\mathbf{r}, t)$ .

- (a) One can show that if  $\psi(\mathbf{r}, t)$  is a solution of the Schrödinger equation with  $\mathbf{A}$  and  $\phi$ , then

$$\psi'(\mathbf{r}, t) = \exp\left(-\frac{iq}{\hbar c} f(\mathbf{r}, t)\right) \psi(\mathbf{r}, t)$$

is a solution of the the Schrödinger equation with  $\mathbf{A}'$  and  $\phi'$ . Is it worrisome that  $\psi(\mathbf{r}, t)$  is *not* gauge-invariant? Explain why. (1 point)

- (b) Define the “velocity” operator as

$$\hat{v}_i = \frac{1}{m}(\hat{p}_{c,i} - \frac{q}{c} A_i).$$

Please rewrite the Hamiltonian in terms of  $\hat{v}_i$ . (1 point)

- (c) Explicitly show whether the expectation values  $\langle \hat{p}_{c,i} \rangle$  and  $\langle \hat{v}_i \rangle$  are gauge-invariant or not (i.e. a few lines of derivations are needed). (2 points)

[Note: Partial credits will be given if you only argue by physics.]

- (d) Suppose we are in an EM field with (certain gauge in which)  $\phi = 0$  and

$$\mathbf{A} = \frac{1}{2}(-B_0 y, B_0 x, 0),$$

where  $B_0$  is a constant. What are the corresponding electric and magnetic fields? (2 points)

$$\lambda = \frac{h}{p}$$

- (e) Show that the commutator

$$[\hat{v}_x, \hat{v}_y] = \frac{i\hbar\omega_L}{m},$$

where  $\omega_L = \frac{qB_0}{mc}$ . (2 points)

- (f) Introduce the annihilation and creation operators

$$\gamma, \gamma^\dagger, \bar{\psi}, \psi$$

$$\hat{a} = \sqrt{\frac{m}{2\hbar\omega_L}}(\hat{v}_x + i\hat{v}_y), \quad \hat{a}^\dagger = \sqrt{\frac{m}{2\hbar\omega_L}}(\hat{v}_x - i\hat{v}_y).$$

Rewrite the Hamiltonian in terms of  $\hat{a}$  and  $\hat{a}^\dagger$ . What are the energy eigenvalues of this system? (2 points)

[Hint: You need to show that  $[\hat{a}, \hat{a}^\dagger] = 1$ .]

The energy spectrum is called the Landau Levels, which is essential in understanding the Quantum Hall Effect.