

# Thermal and Statistical Physics Fall 2019

## Final Exam

Instructor: Pai-hsien Jennifer Hsu

Monday January 6th, 2020

You must show your work. No credits will be given if you don't show how you get your answers.

You may use the following formula:

- Let  $p_j$  be the probability of a system being at the  $j$ -th state, the entropy  $\sigma$  of the system is defined as  $\sigma = -\sum_j p_j \ln p_j$ .
- The internal energy  $U$  is defined as  $U = \sum_j p_j E_j$ , where  $E_j$  is the energy of the  $j$ -th state.
- For a system at thermal equilibrium with temperature  $\tau$  and diffusive equilibrium with chemical potential  $\mu$ , the Gibbs sum is defined as

$$\mathfrak{Z} = \sum_N \sum_s e^{\beta \mu N} e^{-\beta \epsilon_s},$$

where  $\beta = \frac{1}{\tau}$ .  $N$  and  $\epsilon_s$  are all possible number of particles and energy levels of the system. The probability of the system having  $N_i$  particles at energy  $\epsilon_j$  is

$$p_j^{(N_i)} = \frac{e^{\beta(N_i \mu - \epsilon_j)}}{\mathfrak{Z}}.$$

- The first law the thermal dynamics, taking into account the change in particle numbers, can be written as

$$dU = \tau d\sigma - P dV + \mu dN,$$

where  $P$  and  $V$  are the pressure and volume of the system.

- The heat capacity at fixed volume is  $C_V = \left( \frac{\tau \partial \sigma}{\partial \tau} \right)_{V, N}$ .
- The Gibbs free energy  $G = U - \tau \sigma + PV$  tends to be minimized during a process at constant  $\tau$  and  $P$ . The chemical potential can be identified

as the Gibbs free energy per particle, i.e.  $G = N\mu(\tau, P)$ .

- The expected number of particles at a given energy  $\varepsilon$  is

$$f(\varepsilon) = \frac{1}{e^{\beta(\varepsilon-\mu)} \pm 1},$$

where  $\beta = \frac{1}{\tau}$ ,  $+$  for fermions and  $-$  for bosons.

- The density of states  $\mathcal{D}(\varepsilon)d\varepsilon$  is the number of (one-particle) orbitals within  $\varepsilon$  and  $\varepsilon + d\varepsilon$ . For non-relativistic particles in 3D,

$$\mathcal{D}(\varepsilon) = \frac{V}{4\pi^2} \left( \frac{2m}{\hbar^2} \right)^{3/2} \sqrt{\varepsilon}.$$

(ignoring spin degrees of freedom.) The expected number of particles and internal energy are thus given by

$$N = \int_0^\infty d\varepsilon \mathcal{D}(\varepsilon) f(\varepsilon), \quad U = \int_0^\infty d\varepsilon \mathcal{D}(\varepsilon) f(\varepsilon) \varepsilon.$$

- The quantum concentration is  $n_Q = \left( \frac{m\tau}{2\pi\hbar^2} \right)^{3/2}$ .
- Special functions for integrals:

$$\int dx \frac{x^{\nu-1}}{Z^{-1}e^x - 1} = \Gamma(\nu) Li_\nu(Z),$$

where  $\Gamma(\nu)$  is the Gamma function and  $Li_\nu$  is the polylogarithm. For example, the number of particles *in the excited states* in Bose-Einstein condensation (BEC) is  $N_e = \frac{2}{\sqrt{\pi}} n_Q V \Gamma\left(\frac{3}{2}\right) Li_{3/2}(\lambda)$ , where  $\lambda = e^{\beta\mu}$ . At BEC  $\lambda \rightarrow 1$  and  $Li_{3/2}(1) \approx 2.612$ ,  $\Gamma\left(\frac{3}{2}\right) = \frac{\sqrt{\pi}}{2}$ .

- The sum of a geometric series

$$\sum_0^\infty ar^k = a + ar + ar^2 + \dots = \frac{a}{1-r}$$

for  $|r| < 1$ .

- $\ln(AB) = \ln A + \ln B$ ,  $\ln x^n = n \ln x$ ,  $\frac{d}{dx} \ln x = \frac{1}{x}$ .

1. The Gibbs Sum is defined as  $\mathfrak{Z} = \sum_N \sum_s e^{\beta\mu N} e^{-\beta\epsilon_s}$ , where  $\beta = \frac{1}{T}$ .

(a) Show that

$$\langle N \rangle = \frac{1}{\beta} \frac{\partial}{\partial \mu} \ln \mathfrak{Z},$$

and

$$\mu \langle N \rangle - U = \frac{\partial}{\partial \beta} \ln \mathfrak{Z},$$

where  $\langle N \rangle$  is the expected number of particles and  $U$  is the internal energy. (4 points)

- (b) From the definition of entropy  $\sigma = - \sum_{N,s} p_{N,s} \ln p_{N,s}$ , plug in the expression for probability (at certain energy level  $\epsilon_s$  with some number of particles  $N$ ), and **use the meaning of Gibbs free energy**, to show that

$$\beta PV = \ln \mathfrak{Z}$$

with  $P$  and  $V$  being the pressure and the volume of the system. (4 points)

2. In HW (and in class) we've shown that the Gibbs sum for a system with many energy levels  $\epsilon_s$  can be written as the product of the Gibbs sum for *one energy level*, i.e.  $\mathfrak{Z} = \mathfrak{Z}_0 \mathfrak{Z}_1 \mathfrak{Z}_2 \cdots = \prod_s \mathfrak{Z}_s$ .

- (a) For a single state with energy  $\epsilon_r$ , write down the Gibbs sum *for this state*  $\mathfrak{Z}_r$  for a system of identical bosons and a system of identical fermions. (Ignore the spin degrees of freedom.) (4 points)
- (b) Use the formula of  $\langle N \rangle$  from Prob. 1(a) to show that

$$\langle N \rangle = \sum_s \langle N(\epsilon_s) \rangle = \sum_s f(\epsilon_s),$$

where  $f(\epsilon_s)$  is the expected number of particles at energy  $\epsilon_s$  for *fermions or bosons*. (You need to explain both cases.) (2 points)

- (c) Consider a 3D system, use  $\beta PV = \ln \mathfrak{Z}$  from Prob. 1(b) and replace  $\sum_s$  by integrating over  $\int d\epsilon$ , to show that

$$PV = \frac{2}{3}U,$$

as you'd expect for non-relativistic free particles. (4 points)

[Hint: In 3D the density of states  $\mathcal{D}(\epsilon) \propto \sqrt{\epsilon}$ . To do the integral, use  $\sqrt{\epsilon}g(\epsilon) = \frac{d}{d\epsilon} \left( \frac{2}{3}\epsilon^{3/2}g(\epsilon) \right) - \frac{2}{3}\epsilon^{3/2}\frac{dg(\epsilon)}{d\epsilon}$  where  $g(\epsilon)$  is some function of  $\epsilon$ , and integration by parts.]

3. Consider a (3D) system of identical bosons:

- (a) Write down an expression of its internal energy  $U$  using a **dimensionless** polylogarithm function. In other words, you don't need to evaluate the integral. (2 points)

[Hint: Change the variable to  $x = \beta\epsilon$ .]

- (b) If the system is cooled down below the Einstein temperature (in BEC phase), show that the heat capacity  $C_V$  is proportional to  $\tau^{3/2}$ . (3 points)

[Hint: The polylogarithm  $Li_{5/2}(\lambda)$  is just a number at  $\lambda = 1$ , and  $\Gamma(\frac{5}{2})$  is also some constant. How do you evaluate  $C_V$  from  $U$ ?]

4. Consider a 3D system of **relativistic** ( $E \approx pc$ ) identical fermions with spin-1/2:

- (a) Show that for particles numbers  $N$  and volume  $V$ , the fermi energy  $\epsilon_F$  is

$$\epsilon_F = \hbar c (3\pi^2 \frac{N}{V})^{\frac{1}{3}},$$

and the internal energy  $U$  of the ground state is

$$U = \frac{3}{4} N \epsilon_F.$$

(4 points)

[Hint: Use the relation of  $p = \hbar k$ , with  $k = \frac{n\pi}{L}$ ,  $n$  is integer, and  $L = V^{1/3}$ .]

- (b) Show that  $PV = \frac{1}{3}U$  for this system, in contrast to the case of non-relativistic particles. (3 points)

[Hint: Use the relation of  $U$  and  $P$  from the first law of thermodynamics.]