

Quantum Physics II Spring 2019

Final Exam

Instructor: Pai-hsien Jennifer Hsu

Tuesday June 18, 2019

You must show your work. No credits will be given if you don't show how you get your answers.

You may use the following formula:

- The Schrödinger equation:

$$i\hbar \frac{d|\psi(t)\rangle}{dt} = \hat{H} |\psi(t)\rangle$$

where \hat{H} is the Hamiltonian $\frac{\hat{p}^2}{2\mu} + V(\hat{r})$ and μ is the mass of the particle.

If there is an electromagnetic (EM) field, the Hamiltonian is

$$\hat{H} = \frac{(\mathbf{p}_c - q\mathbf{A}/c)^2}{2m} + q\phi$$

where A is the vector potential and ϕ is the electric potential.

- For spherical symmetric potential $V(r)$, there exist simultaneous eigenstates of \hat{H} , \hat{L}^2 , and \hat{L}_z :

$$\hat{H} |E, l, m\rangle = E |E, l, m\rangle, \hat{L}^2 |E, l, m\rangle = l(l+1)\hbar^2 |E, l, m\rangle, \hat{L}_z |E, l, m\rangle = m\hbar |E, l, m\rangle.$$

- Time-independent Perturbation: Consider $\hat{H} = \hat{H}_0 + \hat{H}_1$, where \hat{H}_1 is a small perturbation. The first order energy correction to the n-th energy eigenvalue of \hat{H}_0 is

$$E_n^1 = \langle \psi_n^{(0)} | \hat{H}_1 | \psi_n^{(0)} \rangle$$

where $\psi_n^{(0)}$ is the n-th eigenstate of \hat{H}_0 .

In case $\psi_n^{(0)}$ is degenerated, diagonalize \hat{H}_1 in the subspace of $\psi_n^{(0)}$ to find the "good basis" to use this formula.

- Time-dependent Perturbation: Consider $\hat{H} = \hat{H}_0 + \hat{H}_1(t)$, where $\hat{H}_1(t)$

is a small perturbation. The state $|\psi(t)\rangle$ can be expanded in terms of the unperturbed eigenstates $|E_n^{(0)}\rangle$:

$$|\psi(t)\rangle = \sum_{n=0} c_n(t) e^{-i \frac{E_n^{(0)} t}{\hbar}} |E_n^{(0)}\rangle.$$

Initially ($t = 0$) the particle is at state $|i\rangle$ ($c_i(0) = 1$), the transition amplitude for the particle to be in state $|f\rangle$ at time t is

$$c_f(t) = -\frac{i}{\hbar} \int_0^t dt' \langle f | \hat{H}_1(t') | i \rangle e^{i \frac{(E_f^{(0)} - E_i^{(0)})t'}{\hbar}}.$$

The probability of the particle being at $|f\rangle$ at time t is $|c_f(t)|^2$.

- Fermi's Golden Rule: Suppose there exists a set of final state around $E_f^{(0)}$, the transition probability is

$$\text{Prob.}(t) = \frac{2\pi}{\hbar} \rho(\omega_0) |\langle f | \hat{H}_1 | i \rangle|^2 t,$$

where $\omega_0 = (E_f^{(0)} - E_i^{(0)})/\hbar$ and ρ is the density of states.

- Interaction of a hydrogen atom and EM field (photons): To the first order in \mathbf{A} ,

$$\hat{H}_1 = \frac{e}{m_e} \hat{\mathbf{A}} \cdot \hat{\mathbf{p}}$$

in the Coulomb gauge ($\nabla \cdot \mathbf{A} = 0$).

In the big-box approximation, \mathbf{A} can be expanded as

$$\hat{\mathbf{A}} = \sum_{\mathbf{k}, s} A_0(\omega) (\hat{a}_{\mathbf{k}, s} \epsilon(\mathbf{k}, s) e^{i(\mathbf{k} \cdot \mathbf{r} - \omega t)} + \hat{a}_{\mathbf{k}, s}^\dagger \epsilon^*(\mathbf{k}, s) e^{-i(\mathbf{k} \cdot \mathbf{r} - \omega t)})$$

where s is the polarization, $A_0(\omega)$ is an overall factor that depends on $\omega = c|\mathbf{k}|$, ϵ 's are the polarization vectors, and \hat{a}, \hat{a}^\dagger are the usual creation and annihilation operators of an SHO.

- 3-D Scattering: The wave function $\psi(\vec{r})$ of a particle scattering off a potential $V(\vec{r})$ has the asymptotic behavior

$$\psi(\vec{r}) = A e^{ikz} + A f(\theta, \phi) \frac{e^{ikr}}{r}, \quad r \rightarrow \infty,$$

where the incident wave is in the z -direction.

The differential cross section

$$\frac{d\sigma}{d\Omega} = \frac{\text{Number of particles scattered in to } d\Omega/\text{time}}{\text{Number of incident particles}/(\text{time} \times \text{area})},$$

and $\frac{d\sigma}{d\Omega} = |f(\theta, \phi)|^2$.

- Born Approximation: Useful for high-energy scattering.

$$f(\theta, \phi) = -\frac{m}{2\pi\hbar^2} \int d^3\mathbf{r}' V(\mathbf{r}') e^{i\mathbf{q}\cdot\mathbf{r}'},$$

where m is the mass of the scattering particle, and $\hbar\mathbf{q} = \hbar\mathbf{k}_i - \hbar\mathbf{k}_f$ is the momentum transferred from the initial state $e^{i\mathbf{k}_i\cdot\mathbf{r}}$ to the final state $e^{i\mathbf{k}_f\cdot\mathbf{r}}$.

- Partial wave expansion: Useful for low-energy scattering. For spherically-symmetric $V(r)$,

$$f(\theta) = \sum_{l=0}^{\infty} (2l+1) \frac{e^{i\delta_l}}{k} \sin \delta_l P_l(\cos \theta),$$

where δ_l is the phase shift of the l -th partial wave, and $P_l(\cos \theta)$ is the Legendre polynomial, which relates to the spherical harmonics $Y_{l,m}(\theta, \phi)$ as

$$Y_{l,0}(\theta, \phi) = \sqrt{\frac{2l+1}{4\pi}} P_l(\cos \theta).$$

- For a 1-D simple harmonic oscillator (SHO), $\hat{H} = \frac{\hat{p}_x^2}{2m} + \frac{1}{2}m\omega^2\hat{x}^2$.

The raising and lowering operators are

$$\hat{a}^\dagger = \sqrt{\frac{m\omega}{2\hbar}} \left(\hat{x} - \frac{i}{m\omega} \hat{p}_x \right), \quad \hat{a} = \sqrt{\frac{m\omega}{2\hbar}} \left(\hat{x} + \frac{i}{m\omega} \hat{p}_x \right)$$

and $[\hat{a}, \hat{a}^\dagger] = 1$. The operators get their names from the facts that

$$\hat{a}^\dagger |n\rangle = \sqrt{n+1} |n+1\rangle, \quad \hat{a} |n\rangle = \sqrt{n} |n-1\rangle.$$

One can rewrite \hat{x} and \hat{p}_x as

$$\hat{x} = \sqrt{\frac{\hbar}{2m\omega}} (\hat{a} + \hat{a}^\dagger), \quad \hat{p}_x = -i\sqrt{\frac{m\omega\hbar}{2}} (\hat{a} - \hat{a}^\dagger)$$

and the Hamiltonian as

$$\hat{H} = \hbar\omega \left(\hat{a}^\dagger \hat{a} + \frac{1}{2} \right) = \hbar\omega \left(\hat{N} + \frac{1}{2} \right),$$

where the number operator $\hat{N} \equiv \hat{a}^\dagger \hat{a}$.

- The energy eigenvalues of an SHO are $E_n = (n + \frac{1}{2})\hbar\omega$, $n = 0, 1, 2, \dots$
- Expectation value of an operator \hat{A} for a state $|\psi\rangle$ is $\langle A \rangle = \langle \psi | \hat{A} | \psi \rangle$

$$\omega + \omega^2$$

$$\omega^2 = \frac{k}{m}$$

$$\omega^2 m = k$$

1. Consider a 1-D simple harmonic oscillator (SHO) with an unperturbed Hamiltonian

$$\hat{H}_0 = \frac{\hat{p}_x^2}{2m} + \frac{1}{2}k\hat{x}^2, \quad \omega = \sqrt{\frac{k}{m}}$$

and let $\omega = \sqrt{k/m}$ be the unperturbed frequency of the oscillator.

- (a) Suppose the "spring constant" k is changed a little bit by δk (i.e. $k \rightarrow k + \delta k$). Treat δk as a time-independent perturbation, calculate the first-order energy shift for every unperturbed state $|n\rangle$. (3 points)

- (b) What is the exact energy eigenvalues of the system (after the change of k)? (1 point)

- (c) Expand your result from part (b) to the first order of the small quantity $\delta k/k = \epsilon$. How does that compare to your result from part (a)? (3 points)

[Hint: $(1+x)^n \approx 1+nx$ for small x .]

2. Consider a particle with charge q and mass m_0 . It moves on the surface of a sphere of radius r . The (unperturbed) Hamiltonian is thus

$$\hat{H}_0 = \frac{\hat{L}^2}{2m_0 r^2}.$$

- (a) Write down the ground state(s) and first excited state(s) and their corresponding energies. (2 points)

- (b) Now we turn on an external magnetic field $\mathbf{B} = B_0 \hat{z}$. The resulting Hamiltonian from \mathbf{B} is then $\hat{H}_1 = -\vec{\mu} \cdot \mathbf{B}$, where $\vec{\mu} = \frac{q}{2m_0 c} \mathbf{L}$ is the magnetic moment of the particle.


Treat \hat{H}_1 as a perturbation, evaluate the energy shift of the first excited state(s). Remember to explain how you treat the degeneracies, if there's any. (4 points)

3. Consider a 1-D SHO with charge q , mass m , and frequency ω . It is initially ($t=0$) at the ground state $|0\rangle$. At $t=0$, an external electric field $E = E_0 e^{-\lambda t}$ is turned on. To the lowest order in E_0 , what is the probability that the SHO is in an excited state $|n\rangle$ as $t \rightarrow \infty$? Please give explicit expressions for all $n \geq 1$. (5 points)

4. Scattering:

- (a) Show that for a spherically-symmetric potential $V(\mathbf{r}) = V(r)$, the (first order) Born approximation gives

$$f(\theta, \phi) = -\frac{2m}{\hbar^2 q} \int_0^\infty r' V(r') \sin(qr') dr',$$



where $q = |\mathbf{q}| = |\mathbf{k}_i - \mathbf{k}_f|$. Why is $f(\theta, \phi)$ be independent of ϕ ? (4 points)

[Hint: Choose \mathbf{q} to be the z' -axis in the \mathbf{r}' -integral. Remember that

$$\int d^3\mathbf{r}' = \int_0^\infty r'^2 dr' \int_0^\pi \sin \theta' d\theta' \int_0^{2\pi} d\phi'.$$

]

- (b) Let $V(r) = g\delta(r - a)$. How does the differential cross section $d\sigma/d\Omega$ depend on θ ? (4 points)

[Hint: Use $|\mathbf{k}_i| = |\mathbf{k}_f| = k$.]

- (c) Explain that, *without detailed calculation*, for a spherically-symmetric potential, $d\sigma/d\Omega$ is isotropic (no angular dependence) if we only consider the s-wave ($l = 0$) scattering. (2 points)

[Hint: The zeroth Legendre polynomial $P_0(\cos \theta) = 1$.]

- (d) Take the low-energy limit of your answer in part (b). Is it isotropic? (2 points)

[Note: You have to define what you mean by "low energy".]