

# Quantum Physics I Fall 2018

## Final Exam

Instructor: Pai-hsien Jennifer Hsu

Tuesday January 8, 2019

You must show your work. No credits will be given if you don't show how you get your answers.

You may use the following formula:

- One of the 2-D representations of  $\hat{J}$  is  $\hat{S} = \frac{\hbar}{2}\hat{\sigma}$ , where  $\hat{\sigma}$  are the Pauli matrices

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

- The time-evolution operator  $\hat{U}(t)$ ,  $|\psi(t)\rangle = \hat{U}(t)|\psi(0)\rangle$ , can be written as

$$\hat{U}(t) = \exp\left(-\frac{i}{\hbar}\hat{H}t\right)$$

where  $\hat{H}$  is the time-independent Hamiltonian.  $\hat{H}$  satisfies the Schrödinger equation:

$$i\hbar \frac{d|\psi(t)\rangle}{dt} = \hat{H}|\psi(t)\rangle$$

- The wave function is the projection of the ket vector on the position eigenvector  $\psi(x, t) \equiv \langle x|\psi\rangle$ . The position eigenbasis are normalized as  $\langle x|x'\rangle = \delta(x - x')$ , where the  $\delta$  function satisfies  $\int_{-\infty}^{\infty} f(x)\delta(x - x_0) = f(x_0)$ .

- The Schrödinger equation in terms of wave functions:

$$i\hbar \frac{\partial}{\partial t}\psi(x, t) = \left[-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + V(x)\right]\psi(x, t)$$

For energy eigenstates, this reduces to the time-independent Schrödinger equation

$$\left[-\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x)\right]\psi(x) = E\psi(x)$$

- Expectation value of an operator  $\hat{A}$  for a state  $|\psi\rangle$  is  $\langle A \rangle = \langle \psi|\hat{A}|\psi\rangle$

- Uncertainty of an operator  $\hat{A}$  is defined as  $\Delta A = \sqrt{\langle A^2 \rangle - \langle A \rangle^2}$

- Heisenberg Equation:

$$\frac{d}{dt} \langle A \rangle = \left\langle \frac{\partial A}{\partial t} \right\rangle + \frac{1}{i\hbar} \langle [\hat{A}, \hat{H}] \rangle$$

where  $\hat{H}$  is the Hamiltonian.

- For a spin-1/2 particle, the eigenstates of  $\hat{S}_x$  can be expressed in terms of eigenstates of  $\hat{S}_z$  as

$$|+x\rangle = \frac{1}{\sqrt{2}} |+z\rangle + \frac{1}{\sqrt{2}} |-z\rangle, \quad |-x\rangle = \frac{1}{\sqrt{2}} |+z\rangle - \frac{1}{\sqrt{2}} |-z\rangle$$

- For a system formed by two spin-1/2 particles, there are two choices of eigenbases,  $|m_1, m_2\rangle$  ( $S_z$  of particle 1 and particle 2) and  $|s, m\rangle$  (total spin and total  $S_z$ ), related by ( $\uparrow = +z, \downarrow = -z$ )

Triplet states:

$$|1, 1\rangle = |\uparrow\uparrow\rangle, |1, 0\rangle = \frac{1}{\sqrt{2}} |\uparrow\downarrow\rangle + \frac{1}{\sqrt{2}} |\downarrow\uparrow\rangle, |1, -1\rangle = |\downarrow\downarrow\rangle$$

and the singlet state:

$$|0, 0\rangle = \frac{1}{\sqrt{2}} |\uparrow\downarrow\rangle - \frac{1}{\sqrt{2}} |\downarrow\uparrow\rangle.$$

- Identical particles: The quantum state  $|\psi(a, b)\rangle$  of two particles  $a$  and  $b$  is symmetric  $|\psi(a, b)\rangle = |\psi(b, a)\rangle$  for bosons, and anti-symmetric  $|\psi(a, b)\rangle = -|\psi(b, a)\rangle$  for fermions.

Bosons have integer-spin (spin-0, 1, 2, ...) while fermions have half-integer spin (spin-1/2, 3/2, ...).

- The density operator  $\hat{\rho}$  is defined as

$$\hat{\rho} = \sum_k p_k |\psi^{(k)}\rangle \langle \psi^{(k)}|,$$

where  $p_k$  is the probability to find the system in the state  $|\psi^{(k)}\rangle$ .

- The translational operator  $\hat{T}(a)$ ,  $\hat{T}(a) |x\rangle = |x + a\rangle$ , can be written as

$$\hat{T}(a) = \exp\left(-\frac{i}{\hbar} \hat{p}_x a\right)$$

where  $\hat{p}_x$  is the momentum operator,  $[\hat{x}, \hat{p}_x] = i\hbar$ . In the position space, the momentum operator can be identified as  $\hat{p}_x \rightarrow -i\hbar \frac{\partial}{\partial x}$ .

$$\langle \psi_0 | \hat{p}_x | \psi_0 \rangle$$

- The 1-D probability current is defined as

$$j_x = \frac{\hbar}{2mi}(\psi^* \frac{\partial \psi}{\partial x} - \psi \frac{\partial \psi^*}{\partial x}) = \frac{\hbar}{m} \text{Im}(\psi^* \frac{\partial \psi}{\partial x})$$

- For a simple harmonic oscillator (SHO),  $\hat{H} = \frac{\hat{p}_x^2}{2m} + \frac{1}{2}m\omega^2 \hat{x}^2$ .

The raising and lowering operators are

$$\hat{a}^\dagger = \sqrt{\frac{m\omega}{2\hbar}}(\hat{x} - \frac{i}{m\omega}\hat{p}_x), \quad \hat{a} = \sqrt{\frac{m\omega}{2\hbar}}(\hat{x} + \frac{i}{m\omega}\hat{p}_x)$$

and  $[\hat{a}, \hat{a}^\dagger] = 1$ . The operators get their names from the facts that

$$\hat{a}^\dagger |n\rangle = \sqrt{n+1} |n+1\rangle, \quad \hat{a} |n\rangle = \sqrt{n} |n-1\rangle.$$

One can rewrite  $\hat{x}$  and  $\hat{p}_x$  as

$$\hat{x} = \sqrt{\frac{\hbar}{2m\omega}}(\hat{a} + \hat{a}^\dagger), \quad \hat{p}_x = -i\sqrt{\frac{m\omega\hbar}{2}}(\hat{a} - \hat{a}^\dagger)$$

and the Hamiltonian as

$$\hat{H} = \hbar\omega(\hat{a}^\dagger \hat{a} + \frac{1}{2}) = \hbar\omega(\hat{N} + \frac{1}{2}),$$

where the number operator  $\hat{N} \equiv \hat{a}^\dagger \hat{a}$ .

- The energy eigenvalues of a SHO are  $E_n = (n + \frac{1}{2})\hbar\omega$ ,  $n = 0, 1, 2, \dots$
- The coherent states of a SHO are states which satisfy  $\hat{a} |\alpha\rangle = \alpha |\alpha\rangle$ , where  $\alpha$  can be a complex number.  $|\alpha\rangle$  can be expanded in the energy eigenbasis as

$$|\alpha\rangle = e^{-|\alpha|^2/2} \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle$$

$$\frac{\sqrt{3}}{2} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + \frac{i}{2} \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} \quad A \quad \frac{A}{2} \quad \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & +\hbar\omega & 0 & 0 \\ 0 & 0 & -\hbar\omega & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

1. Consider a system with two spin-1/2 particles in the state

$$|\psi\rangle = \frac{\sqrt{3}}{2} |\uparrow\uparrow\rangle + \frac{i}{2} |\downarrow\downarrow\rangle \quad \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \quad \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

where  $|\uparrow\rangle$  and  $|\downarrow\rangle$  refer to spin along the z-axis.

- Let  $p$  be the probability of finding the system in  $|\uparrow\uparrow\rangle$ , and  $q$  be the probability of  $|\downarrow\downarrow\rangle$ . Can we describe the system as a random "mixture of  $p|\uparrow\uparrow\rangle$  and  $q|\downarrow\downarrow\rangle$ "? Explain in terms of density matrices (i.e. Write down the density matrices for both cases and compare them). (2 points)
- What is the probability of finding particle 1 in the  $|+z\rangle$  state? (1 point) If we find particle 2 in the  $|-z\rangle$  state, what is the probability of getting  $|+z\rangle$  if we measure particle 1? (1 point)
- [Bonus] Is this an entangled state? Briefly explain why. (1 point)
- The system (in the given state  $|\psi\rangle$ ) is subject to a potential  $V(x_1, x_2) = \frac{1}{2}m\omega^2(x_1^2 + x_2^2)$ , where  $x_1$  and  $x_2$  are the positions of particles 1 and 2. What is the ground state energy if they are identical particles? (2 points)
- What is the probability of finding the system with total  $S_x = 0$ ? (2 points)  
[Hint: Rewrite  $|\psi\rangle$  in  $\{|\uparrow\uparrow\rangle, |\uparrow\downarrow\rangle, |\downarrow\uparrow\rangle, |\downarrow\downarrow\rangle\}$  along the x-axis.]
- Suppose we measure total  $S_x$  and find  $S_x = 0$ . At time  $t = 0$  we remove the potential  $V(x_1, x_2)$ , and turn on an external magnetic field. The resulting Hamiltonian is

$$\hat{H} = \omega_0(\hat{S}_{1x} + \hat{S}_{2x}) \quad \begin{pmatrix} \sqrt{3}-i & 0 \\ 0 & \sqrt{3}+i \end{pmatrix} \quad \begin{pmatrix} 3 & -1 \\ -1 & 3 \end{pmatrix}$$

Will the system oscillate between spin-0 and spin-1? Explain why. (2 points)

2. Consider a normalized wave function  $\psi_0(x)$ . Let  $\langle x \rangle = x_0$  and  $\langle p_x \rangle = p_0$  for  $\psi_0(x)$ .

- Let  $\psi_1(x) = e^{iqx/\hbar}\psi_0(x)$ . Express  $\langle x \rangle$  and  $\langle p_x \rangle$  for  $\psi_1(x)$  in terms of  $x_0$ ,  $p_0$ , and  $q$ . (2 points)
- Based on your results, what is the physical meaning of adding an overall factor  $e^{iqx/\hbar}$  to a wave function? (1 point)
- Let  $m$  be the mass of the particle, and write  $\psi_1(x)$  as  $A(x)e^{i\theta(x)}$ , where  $A(x)$  and  $\theta(x)$  are real functions corresponding to its amplitude and phase. Show that the probability current is proportional to the gradient of the phase  $(\partial_x \theta(x))$ . (2 points)

3. A particle with mass  $m$  is trapped in a  $\delta$ -function potential well located at  $x = 0$ . We write the potential as  $V(x) = -\frac{\hbar^2}{mb}\delta(x)$  for some  $b > 0$ .

- (a) The parameter  $b$  carries the unit of length. Use dimensional analysis (analyze the units of available parameters), show that the "natural" energy scale  $E_0$  from the parameters of the system ( $\hbar$ ,  $m$ , and  $b$ ) is

$$E_0 = \frac{\hbar^2}{mb^2}$$

(2 points)

- (b) Assume the wave function of the particle is of the form

$$\psi(x) = ce^{-|x|/\lambda}$$

$$\frac{\hbar^2}{2m} \frac{(-1/\lambda)^2}{\hbar^2} E = 0$$

Use the boundary conditions of the wave functions at (around)  $x = 0$  to determine  $\lambda$  and the bound-state energy  $E_b$  using  $\hbar$ ,  $m$ , and  $b$ . How does  $E_b$  compare with the energy scale  $E_0$  from part (a)? (3 points)

[Hint: Are the wave functions and their derivatives continuous? If not, how do they change?]

- (c) [Bonus] Based on your previous results, briefly explain what the parameter  $b$  represents. For example, can you estimate the size of  $\Delta x$  of the bound state *without calculations*? (2 points)

$$\Delta x \sim \frac{\hbar}{\sqrt{2m|E_b|}} \sim \frac{\hbar}{\sqrt{2m \frac{\hbar^2}{mb^2}}} \sim b$$

4. Consider a coherent state that satisfies  $\hat{a}|\alpha\rangle = \alpha|\alpha\rangle$ , where  $\alpha = |\alpha|e^{-i\delta}$  for some real phase  $\delta$  at time  $t = 0$ .

$$C_0 e^{i\delta} + C_1 e^{-i\delta} + C_2 e^{i\delta} + C_3 e^{-i\delta}$$

- (a) Prove that at time  $t$ , the state  $|\alpha(t)\rangle$  is still a coherent state, with  $\alpha(t) = \alpha e^{-i\omega t}$ . (2 points)
- (b) Find  $\langle x \rangle$  and  $\langle p_x \rangle$  as a function of time. Express your answers in terms of  $|\alpha|$  and  $\delta$ . (2 points)
- (c) Show that for this coherent state,  $\Delta N = \sqrt{\langle N \rangle}$ , where  $\hat{N}$  is the number operator  $\hat{a}^\dagger \hat{a}$  (4 points).
- (d) Express the expectation value of energy  $\langle E \rangle$  in terms of  $\langle x \rangle$  and  $\langle p_x \rangle$ . If we treat  $\langle x \rangle$  and  $\langle p_x \rangle$  as classical  $x(t)$  and  $p_x(t)$ , how does your answer compare to the energy of a classical SHO? (2 points)

$$\langle \alpha | \hat{a}^2 - 2\hat{a}\hat{a}^\dagger + \hat{a}^{\dagger 2} | \alpha \rangle$$

$$\alpha^2 - 2\alpha\alpha^* + \alpha^{*2}$$

$$[a, a^\dagger] = 1$$

$$\hat{a}\hat{a}^\dagger - \hat{a}^\dagger\hat{a} = 1$$