

Geometric Shapes: Quadric Surfaces and Sweeping

Module 3
Lecture 5

CZ2003

1

Quadric Surfaces

- $ax^2+by^2+cz^2+dxy+eyz+fxz+gx+hy+kz+m=0$
- Ellipsoids (including spheres), Elliptic Paraboloid, Hyperbolic Paraboloid, Hyperboloid of One Sheet, Hyperboloid of Two Sheets, Cone, Elliptic Cylinder (including circular cylinders), Hyperbolic Cylinder, Parabolic Cylinder
- Only highlighted surfaces are examinable

CZ2003

3

We have learnt that


- Plane surfaces can be defined by explicit, implicit and parametric functions
- In implicit (linear) equation $Ax+By+Cz+D=0$, $N=[A \ B \ C]$ while D defines displacement from the origin
- To get the plane equation, derive N first (e.g., by cross product of two vectors), then substitute any x,y,z on the plane to derive D .
- Implicit equation in intercepts can be easily written from $x/a+y/b+z/c=1$
- Parametric definition of plane is based on linear equations of two parameters
- Bilinear interpolation is an extension of linear interpolation for interpolating functions of two variables (e.g., u, v). It performs linear interpolation first in one direction, and then again in the other direction. Although each step is linear, the interpolation as a whole is not linear but rather quadratic in the sample location
- Bilinear surface defines a 4-sided polygon, including non-planar surfaces.
- Bilinear surface can be used for writing parametric functions of a triangle as well: two of the vertices are simply merged together which will also simplify the defining formulas.

CZ2003

2

Sphere

- Implicit $r^2 - x^2 - y^2 - z^2 = 0$
- Explicit $z = \pm \sqrt{r^2 - x^2 - y^2}$
- Parametric



$$x = r \cos \varphi \sin \theta$$

$$y = r \sin \varphi$$

$$z = r \cos \varphi \cos \theta$$

$$-\frac{\pi}{2} \leq \varphi \leq \frac{\pi}{2}$$

$$-\pi \leq \theta \leq \pi$$

Two parameters !


CZ2003

4

Ellipsoid

- Implicit $1 - (x/a)^2 - (y/b)^2 - (z/c)^2 = 0$
- Explicit $z = \pm c \sqrt{1 - (x/a)^2 - (y/b)^2}$
- Parametric

By scaling of a sphere with radius 1
by a, b, c



$$x = a * \cos \varphi \sin \theta$$

$$y = b * \sin \varphi$$

$$z = c * \cos \varphi \cos \theta$$

$$-\frac{\pi}{2} \leq \varphi \leq \frac{\pi}{2}$$

$$-\pi \leq \theta \leq \pi$$

Two parameters !


CZ2003

5

Experimenting with Quadrics

$$2^2 - x^2 - y^2 - z^2 = 0$$

$$2^2 - (x-1)^2 - y^2 - z^2 = 0$$




sphere

CZ2003

6

Experimenting with Quadrics

$$1 - (x/1)^2 - (y/0.5)^2 - (z/0.75)^2 = 0$$



ellipsoid


CZ2003

7

Experimenting with Quadrics

$$1^2 - x^4 - y^4 - z^4 = 0$$

$$1^2 - x^{16} - y^{16} - z^{16} = 0$$




Super-ellipsoid

CZ2003

8


Experimenting with Quadrics

$$(z/a)^2 - (x/b)^2 - (y/c)^2 = 0$$



Cone

$$1 - (x/b)^2 - (y/c)^2 = 0$$




Elliptic Cylinder

CZ2003


9

Experimenting with Parametric Sphere

$$\begin{aligned} X &= 0.7 \cdot \cos(u) \cdot \cos(v) \\ Y &= 0.7 \cdot \cos(u) \cdot \sin(v) \\ Z &= 0.7 \cdot \sin(u) \end{aligned}$$




$$u = [0, 2\pi] \quad v = [0, \pi]$$



$$u = [0, \pi] \quad v = [0, \pi]$$

$$\begin{aligned} X &= 0.7 \cdot \cos(u) \cdot \cos(v) + 1 \\ Y &= 0.7 \cdot \cos(u) \cdot \sin(v) \\ Z &= 0.7 \cdot \sin(u) \end{aligned}$$



$$u = [0, 2\pi] \quad v = [0, \pi]$$

CZ2003

10

We have learnt that

- Plane and quadric surfaces can be defined by explicit, implicit and parametric functions
- Parametric definition of plane is based on linear equations of two parameters
- Bilinear interpolation is an extension of linear interpolation for interpolating functions of two variables (e.g., u , v). It performs linear interpolation first in one direction, and then again in the other direction. Although each step is linear, the interpolation as a whole is not linear but rather quadratic in the sample location
- Curves and surfaces were considered as objects sampled by a moving point:
1 DOF for curves and 2 DOFs for surfaces
- Surfaces can also be thought as objects sampled by moving curves:
planes and bilinear surfaces – by moving straight lines
spheres and ellipsoids – by rotating circles and ellipses, etc

CZ2003

11


Sweeping

- Shapes are created by curve moving along some path
- Two particular cases of sweeping--*translational* and *rotational sweeping*--can be easily defined parametrically

CZ2003

12


Translational and Rotational Sweeping



CZ2003

13


Parametric Representation of Translational Sweeping



CZ2003

14

Cylinder by Translational Sweeping a Circle




Translational sweeping

CZ2003

15


Translational Sweeping. Summary



CZ2003

16


Parametric Representation of Rotational Sweeping. Making a Sphere



CZ2003

17


Parametric Representation of Rotational Sweeping. Making a Sphere



CZ2003

18


Parametric Representation of Rotational Sweeping. Making a Cone



CZ2003

19


Parametric Representation of Rotational Sweeping. Making a Cone



CZ2003

20

Rotational Sweeping. Summary



CZ2003

21

Surfaces. Summary

- Polygonal representation - polygon meshes
- Analytic representations
 Define how a point can move along the surface with 2 degrees of freedom (forward, backward, left, right).
 - Explicit representation.
 Seldom used because of axes dependency. Can be derived from implicit functions if needed.
 - Implicit representation
 - Parametric representation.
 Two parameters must be used to define a surface. Often, the parameters can be considered as a kind of latitude and longitude coordinates for locating a point on the surface.

CZ2003

22