

Brain abnormalities in attention-deficit hyperactivity disorder: a review

Katya Rubia, Analucía A. Alegría, Helen Brinson

Aim. To review the magnetic resonance imaging findings in child and adult attention-deficit hyperactivity disorder (ADHD).

Development. Studies have shown that ADHD is characterised by multiple functional and structural neural network abnormalities including most prominently fronto-striatal, but also fronto-parieto-temporal, fronto-cerebellar and even fronto-limbic networks. Evidence from longitudinal structural imaging studies has shown that ADHD is characterised by a delay in structural brain maturation. This is reinforced by indirect evidence from cross-sectional imaging studies for more immature brain function as well as structural and functional connectivity patterns, which, however, needs corroboration by longitudinal studies. Dysfunction of the ventrolateral prefrontal cortex seems to be more pronounced in ADHD relative to other pediatric disorders and there is some evidence for differential abnormalities in the basal ganglia. A meta-analysis of stimulant effects on brain function shows that the most consistent mechanism of action of acute psychostimulant medication is the increased activation of the inferior prefrontal cortex and the basal ganglia. First attempts to use neuroimaging data to make individual diagnostic classifications of ADHD children based on pattern recognition techniques are promising but need replication across centres and scanners.

Conclusions. The last two decades of neuroimaging have shaped out biomarkers of ADHD. Future studies will need to focus on using this information for clinical translation such as using neuroimaging for individual diagnostic and prognostic classification or by using neuroimaging as a neurotherapy to reverse those brain function abnormalities that have been established over the last two decades of neuroimaging.

Key words. Atomoxetine. Attention-deficit hyperactivity disorder (ADHD). Diffusion tensor imaging (DTI). Functional magnetic resonance imaging (fMRI). Methylphenidate. Psychostimulants.

Introduction

Attention-deficit hyperactivity disorder (ADHD) is characterised by symptoms of age-inappropriate inattention, hyperactivity and impulsivity [1]. ADHD is one of the most common childhood psychiatric disorders, affecting 3-8% of school-aged children, with 65% of cases persisting into adulthood [2]. ADHD patients have deficits in higher-level cognitive functions necessary for goal-directed behaviors, so-called 'executive functions' (EF), that are known to be mediated by late developing fronto-striato-parietal and fronto-cerebellar networks [3]. The most consistent deficits are in motor response inhibition, sustained attention, and working memory [4,5] as well as timing processing, in particular time estimation [6,7] and temporal foresight, as measured in temporal discounting and gambling tasks [7].

Objectives

ADHD is the most imaged child psychiatric disorder with over hundreds of published structural and

functional imaging studies. This review discusses the most consistent deficit findings in brain structure, function and structural and functional connectivity. Furthermore, the review will cover current findings on disorder-specificity of these brain deficits in ADHD relative to other child psychiatric disorders. The effects of stimulant and non-stimulant medications on the brain structure and function of ADHD patients will also be reviewed and discussed. Last, we will discuss clinical applications of neuroimaging. This includes the use of multivariate pattern recognition analyses for imaging-based individual diagnostic classification of ADHD and neurotherapy methods that aim to upregulate dysfunctional brain regions in ADHD through neurofeedback or brain stimulation.

Structural MRI studies

The vast majority of structural MRI (sMRI) studies have used region of interest (ROI) analyses, focusing on apriori hypothesised regions, typically with relatively liberally chosen thresholds. The most con-

Department of Child & Adolescent Psychiatry. Institute of Psychiatry. King's College. London, United Kingdom.

Corresponding author:

Prof. Katya Rubia. Department of Child Psychiatry/SGDP, P046. Institute of Psychiatry. 16 De Crespigny Park. London, SE5 8AF, United Kingdom.

Fax:

+44 207 8480866.

E-mail:

katya.rubia@kcl.ac.uk

Financial disclosure:

K.R. has received speaker's honoraria from Lilly and Shire.

Acknowledgements:

K.R. has received support from the National Institute of Health Research (NIHR) Biomedical Research Centre (BRC) for Mental Health at South London and Maudsley NHS Foundation Trust and Institute of Psychiatry, Kings College London. H.B. received post-doctoral support from an Action Medical Research grant to K.R. A.A.A. was supported by an Institute of Psychiatry PhD excellence award.

Accepted:

10.01.14.

How to cite this paper:

Rubia K, Alegría AA, Brinson H. Brain abnormalities in attention-deficit hyperactivity disorder: a review. Rev Neurol 2014; 58 (Supl 1): S3-S18.

Versión española disponible en www.neurologia.com

© 2014 Revista de Neurología

sistent findings of a meta-analysis of ROI studies were reductions in total and right cerebral volumes, including several prefrontal regions, in the posterior or inferior cerebellar vermis, the splenium, the corpus callosum, and the right caudate [8], which were also observed in later larger numbered sMRI studies [9-12]. A few recent ROI sMRI studies also found grey matter (GM) or cortical thickness abnormalities in subcortical limbic regions such as insula [13], amygdala [14] and thalamus [15]. These ROI-based meta-analytic findings confirm the notion that ADHD patients have deficits in fronto-striatal and fronto-cerebellar networks that mediate the late developing EF that are impaired in the disorder [16]. ROI studies, however, are biased towards apriori hypothesised regions. Whole brain imaging studies are more suitable to reveal the most consistent brain abnormalities without unnecessarily restricting the search volume. Three meta-analyses have been published on the remarkably few whole brain voxel-based morphometry (VBM) sMRI studies of ADHD. All studies showed that the right basal ganglia were consistently reduced in GM in ADHD relative to healthy controls [17-19]. In addition, the deficits were most pronounced in pediatric studies with adult studies not showing any deficits [18,19]. Given the extensive connections between the basal ganglia and frontal regions [16], a primary deficit in the basal ganglia, however, still implies fronto-striatal circuit abnormalities in ADHD [16]. These meta-analytic findings tie in well with our neurochemistry abnormality findings in ADHD based on a meta-analysis of positron emission tomography (PET) studies where we showed consistently reduced dopamine transporter levels in the basal ganglia in medication-naïve ADHD patients [20]. The only enhanced GM volume in ADHD patients relative to controls was in the precuneus, which is part of the default mode network (DMN) [18]. The DMN consists of intercorrelated co-activation of medial frontal lobe, anterior and posterior cingulate (ACC/PCC) and inferior temporal and parietal areas during rest, that are parametrically attenuated during cognitive load, and thought to represent 'mental clutter'. ADHD patients have more attentional lapses and attenuated deactivation of the DMN during attention tasks [21-23]. The enlarged volume size in this region could therefore be a plastic consequence of enhanced DMN activation (i.e. diminished deactivation).

In conclusion, the most consistent abnormality in ADHD patients based on whole-brain cross-sectional imaging studies is the reduced GM in the basal ganglia, with additional evidence for abnor-

mal GM and cortical thickness abnormalities in frontal, temporal and cerebellar regions based on ROI studies.

It has been argued that ADHD children suffer from delayed brain maturation, due to their relative immaturity in behavioural features that diminish naturally with age such as impulsiveness and inattention and in cognitive functions that are mediated by late developing fronto-striatal and fronto-cerebellar systems [3]. Seminal longitudinal imaging studies from the National Institute of Mental Health (NIMH) provided direct evidence for this hypothesis by showing that 232 ADHD patients relative to 232 healthy controls had a delay in the peak of cortical thickness and surface area by 2-5 years, with the most prominent delay in frontal, superior temporal and parietal regions [11,24].


Diffusion tensor imaging (DTI) studies show that ADHD children and adults have deficits not only in isolated brain regions but in the white matter (WM) connectivity between these regions, most prominently between fronto-striatal, fronto-parieto-temporal, and fronto-cerebellar connections [25-30], as also shown in a recent meta-analysis of 9 DTI studies using whole brain analyses [31]. Furthermore these WM tract abnormalities, reflecting poor myelination or reduced axonal branching, have been related to clinical behavioural and cognitive abnormalities [30,32]. Consistent with the maturational delay hypothesis of ADHD, one study found slower development of WM in the caudate nucleus over adolescence in ADHD, which reached normal levels by adulthood [33] while 2 other studies that tested for a range of WM tracts, found almost all WM tracts to be attenuated in ADHD, and, given that most of these WM tracts develop well into mid-adulthood [34], hypothesised a possible global delay in WM tract development [35,36]. This will, however, have to be corroborated in longitudinal DTI studies.

Functional MRI studies

Functional magnetic resonance imaging (fMRI) studies have provided consistent evidence for fronto-striatal, fronto-parietal and fronto-cerebellar deficits in ADHD during tasks of cognitive control with some emerging evidence for fronto-limbic abnormalities in the context of reward processing.

A meta-analysis of 55 whole-brain fMRI studies across EF, memory, reward and timing tasks in 16 adult and 39 pediatric studies, including a total of 741 ADHD and 801 control subjects, showed sig-

Figure 1. Four meta-analyses of fMRI studies of ADHD patients for different cognitive domains. The meta-analyses show underactivation in ADHD patients in several dissociated fronto-striatal and fronto-cerebellar networks during the respective cognitive domains. a) During motor response inhibition, ADHD patients show underactivation relative to healthy controls in the right ventral inhibition network, in right IFC, SMA, the basal ganglia and thalamus. They had enhanced activation in posterior cingulate gyrus [38]. b) During interference inhibition, ADHD patients had underactivation in right IFC, ACC, the basal ganglia and thalamus and enhanced activation in ventral anterior cingulate [38]. c) During attention tasks, ADHD patients showed reduced activation relative to healthy controls in the right dorsal attention network, comprising right DLPFC, the posterior part of the basal ganglia and thalamus, inferior parietal lobe and precuneus. ADHD patients have enhanced activation relative to controls in cerebellar and occipital regions [38]. d) During timing tasks, ADHD children had reduced activation in a predominantly left hemispheric timing network, comprising left IFC, left inferior parietal lobe and right cerebellum. ADHD patients had enhanced activation in a default mode region, the posterior cingulate cortex [22]. The enhanced activation in anterior and posterior cingulate could also reflect decreased activation of the default mode network in ADHD versus healthy controls.



nificant hypoactivation in ADHD relative to controls in bilateral ventral attention networks –inferior frontal cortex (IFC), basal ganglia– and predominantly right hemispheric fronto-temporo-parietal networks, including DLPFC/IFC, basal ganglia, thalamus, ACC and SMA. In addition, hyperactivation was observed predominantly in default mode regions as well as in visual and somatomotor regions [37]. Three other meta-analyses focused on specific cognitive domains. Thus, a meta-analysis of 21 whole-brain fMRI studies of cognitive and motor inhibition, including 7 adult and 14 pediatric studies, showed that 287 ADHD patients relative to 320 healthy controls had consistently reduced activation in key regions of inhibition, in right inferior

prefrontal cortex (IFC), supplementary motor area (SMA), ACC, left striatum and right thalamus [38]. When separated by motor response and interference inhibition tasks, the SMA was more prominently underactivated for motor response inhibition and the ACC for interference inhibition (Figs. 1a, 1b). A meta-analysis on attention tasks included 13 mostly pediatric whole-brain fMRI studies and found underactivation in 171 ADHD patients relative to 178 healthy controls in the right hemispheric dorsal attention network, comprising the right DLPFC, right inferior parietal cortex and caudal parts of the basal ganglia and thalamus. In addition, ADHD patients had increased activation relative to controls in right cerebellum and left cuneus, pre-

sumably compensating for the underactivation of the frontal part of the dorsal DLPFC-parieto-cerebellar attention network [38] (Fig. 1c). A meta-analysis of timing functions in ADHD, including 11 fMRI studies showed consistently reduced activation in 150 ADHD patients relative to 145 healthy controls in left IFC, left inferior parietal lobe and the right lateral cerebellum [22], all key regions of timing functions [39]. The timing meta-analysis in addition also showed increased activation in ADHD patients in default mode regions, the PCC and precuneus [22] (Fig. 1d).

Interestingly, the brain regions that are underactivated in ADHD increase progressively in their activation with age [3], which could potentially reflect a developmental delay in brain function in ADHD patients. Problems deactivating the DMN have been associated with more attention lapses, both in normal development and in ADHD [23], and could also reflect a developmental delay [3]. Both the problematic deactivation of the DMN concomitant with poor activation of task-relevant and age-correlated brain activation may hence reflect a developmental delay of brain function and both are likely underlying the poor performance in ADHD patients in attention-demanding higher-level cognitive tasks.

In summary, the meta-analyses of whole-brain fMRI studies in ADHD show that ADHD patients have cognitive-domain dissociated deficits in multiple dorsal and ventral fronto-striato-parietal and fronto-cerebellar networks, including IFC-ACC/SMA-striato-thalamic networks for inhibitory control functions, right DLPFC-parieto-striato-cerebellar networks for attention functions and left IFC-parieto-cerebellar networks for timing functions [22,37,38] (Fig. 1). In addition, poor task-related activation appears to be concomitant with poor deactivation of the DMN, both of which are likely to underlie compromised performance in ADHD.

Abnormalities, however, have also been observed during reward processing. A meta-analysis of 8 ROI fMRI studies of reward anticipation in 340 ADHD patients and healthy controls, most of which used the same monetary reward anticipation task (MID), showed consistent underactivation of the ventral striatum in ADHD adults and children relative to healthy controls with a medium effect size [40]. An important caveat, however, is that deficits in ventral striatum have only been observed in ROI studies. Future large-scale fMRI studies will have to confirm the presence of abnormalities in the ventral striatum using whole brain image analyses.


In addition to deficits in the function of specific frontal, striatal, temporo-parietal and cerebellar re-

gions, functional connectivity studies have demonstrated that ADHD patients also have abnormalities in the functional inter-regional connectivity between these regions both during rest and during cognitive tasks [41]. Thus, during the resting state, ADHD children have reduced functional connectivity in the DMN, mostly between ACC and PCC [42-45], as well as in fronto-striato-thalamic, fronto-temporal and sensorimotor circuitries [46-49]. An interesting double dissociation was observed between attenuated OFC-ventral striatum-limbic functional connectivity and emotion dysregulation and DLPFC-striato-cingulo-parietal functional connectivity abnormalities and poor EF [50]. Reduced functional connectivity has also been observed in children and adults with ADHD within fronto-striato-cerebellar networks during cognitive tasks such as sustained attention [51], inhibition, time estimation [52-54], working memory [55,56] and response preparation [57]. Furthermore, there is evidence for stronger coherence of the DLPFC with DMN and reduced anti-correlation between both in relation to poor attention performance, suggesting a more diffuse connectivity between functional networks in ADHD, where both DMN intrudes during attention-demanding contexts and DLPFC signalling is insufficiently suppressed in relation to DMN activity [21,58]. The findings from these functional connectivity studies suggest both abnormal task-based cortico-striatal-thalamic and abnormal DMN functional 'networks' that are poorly anti-correlated and together contribute to the cognitive and behavioural symptoms of ADHD. Given that DMN as well as task-based functional connectivity mature progressively with increasing age and are associated with progressive cognitive maturation [3,59], the more immature functional connectivity in these networks may reflect delayed functional brain maturation. This hypothesis was supported by a multivariate pattern recognition study that showed that the resting state functional connectivity abnormalities that classified ADHD adults relative to controls was similar to the pattern observed in younger typically developing subjects [60]. Future longitudinal fMRI studies will need to corroborate this hypothesis of a delay in the maturation of brain function and associated networks.

Effects of stimulant and non-stimulant medications on the ADHD brain

Stimulant medication (e.g. methylphenidate or dexamphetamines) are first line treatment for ADHD

Figure 2. Most consistently increased brain activation within ADHD adolescents after a single clinical dose of psychostimulant medication relative to placebo or off-medication. The most consistent activation increase was in right inferior prefrontal cortex ($p < 0.005$) followed by increased activation in ventral ACC and the putamen ($p < 0.05$). There was a significant decrease of activation with placebo in dorsal ACC and supplementary motor area relative to stimulant medication, which could also potentially reflect increased deactivation of these regions for medication relative to placebo [65].



as they reduce the severity of ADHD core symptoms in up to 70% of patients [61]. Several fMRI studies have sought to further our understanding on the acute and chronic effects of stimulant medications on the function of the ADHD brain.

Several well-designed (e.g. randomised placebo-controlled, case-control crossover) whole-brain and ROI fMRI studies have examined the acute effects of Methylphenidate on brain function during a series of cognitive tasks in medication-naïve ADHD patients. An acute dose of methylphenidate in medication-naïve or in chronically medicated ADHD youth has been shown to increase and normalise the underactivation in right and/or left IFC as well as ACC during motor and interference inhibition [62–66], timing [6,65,67], error processing [64] and sustained attention [51], but had no effect during working memory [68,69]. Effects were also observed in the cerebellum during time discrimination [6], interference and motor inhibition [64–66] and attention tasks [51]; and in the striatum during reward [51] and response inhibition tasks [64–66,70].

Our meta-analysis of all 14 published whole-brain fMRI studies that tested the effects of stimulants on brain function in ADHD showed that the most consistent acute effect is the increased activation of right IFC/insula, with additional effects in the putamen at a more lenient threshold [65] (Fig. 2). A few studies investigated the acute effects of stimulants on functional connectivity and found that an acute dose of methylphenidate normalized functional connectivity deficits in fronto-striatal, fronto-parietal and fronto-cerebellar networks during vigilant attention [51], in fronto-parietal networks during working memory [55], in ventral ACC and lateral PFC connectivity during a cognitive

Stroop task [54] and between the amygdala and lateral PFC [71] during an emotional Stroop task.

Relatively little, however, is known about chronic effects of psychostimulants. A 12 months trial of stimulant administration in children with ADHD normalized the enhanced (potentially compensatory) activation in insula and putamen during a reorienting attention process. Also, there was a trend for ACC dysfunction to be more pronounced in 5 unmedicated relative to 9 chronically medicated patients, suggesting long-term amelioration [72]. OROS methylphenidate over 3 months normalized reduced ventral striatum and thalamus activation during the processing of low but not high reward outcomes [73]. In adult ADHD, a 6-week trial of stimulant medication compared to placebo increased activation in the dorsal ACC, DLPFC, pre-motor and parietal cortices, caudate, thalamus and cerebellum during interference inhibition [74]. Our meta-regression analyses across fMRI studies of attention and inhibition found that long-term stimulant administration (between 6 months to 3 years) was associated with normalization of right caudate activity during attention [38] and of right DLPFC activity during timing tasks [22].

In conclusion, the meta-analytic findings suggest that acute and longer-term stimulant medication treatment is most consistently associated with an upregulation of two key dysfunctional areas in ADHD, the right IFC and the basal ganglia.

Unfortunately there are no prospective longitudinal imaging studies of long-term stimulant effects on brain structure. However, retrospective comparisons of medicated and non-medicated ADHD patients suggest that medicated patients with ADHD have more normal size, volumes and/or morphology than unmedicated patients in ADHD-relevant

brain regions including the right ACC [75]; the anterior thalamic pulvinar [15]; the posterior inferior vermis of the cerebellum [76]; the left lateral cerebellar surface [77]; the basal ganglia [78]; and the corpus callosum [79]. Two studies of retrospective comparisons within longitudinal data found more normal WM [80], and a less rapid cortical thinning development in left IFG, premotor and parietal regions in long-term medicated relative to non-medicated children [81]. However, there have also been negative findings in relatively small sampled studies [9,82]. Two meta-regression analyses of VBM studies tested for long-term medication effects. Both studies found that long-term stimulant medication was associated with more normal basal ganglia volumes in ADHD [18,19].

In conclusion, overall sMRI and fMRI studies suggest that stimulant medication may potentially be neuro-protective on brain structure and function. Studies that have tested for neurochemical effects, however, have been less promising. Our meta-analysis of PET studies in mostly adult ADHD patients, showed that long-term stimulant medication was associated with an abnormally increased level of striatal dopamine transporters, which were reduced relative to healthy controls in medication-naïve patients, suggesting potential brain adaptation to stimulants [20]. This was also observed in 10 adults with ADHD in a within-subject study design after a one year follow-up of chronic stimulant medication treatment [83]. These findings of plastic long-term upregulation of DAT with chronic stimulant medication could explain relatively poor long-term efficacy of stimulant medication [84]. However, prospective longitudinal imaging studies within a randomized placebo-controlled design are crucial to confirm these findings of plastic effects based on cross-sectional comparisons on brain structure, function and neurochemistry.

Very few studies have tested brain effects of the only other licensed medication for ADHD, atomoxetine. In a placebo-controlled randomized study, we found shared effects of atomoxetine and methylphenidate of upregulation of right IFC activation during time discrimination and of bilateral IFC activation during inhibition, which was furthermore normalised with both drugs in their underactivation in ADHD patients under placebo relative to healthy controls [62,67]. Both drugs also elicited compensatory fronto-striato-thalamic overactivation in ADHD children during working memory and both drugs deactivated DMN activation [68]. Drug-specific effects, however, were also observed with atomoxetine upregulating and normalised

right DLPFC underactivation during working memory [68] and Methylphenidate increasing compensatory activation in left IFC and the basal ganglia during response execution and one of the WM conditions [68]. Furthermore, methylphenidate had drug-specific effects on the activation of the dopaminergically innervated SMA during motor response execution and time discrimination [62, 67]. A few recent fMRI studies tested chronic effects of atomoxetine. The only pilot study in adult ADHD showed that atomoxetine treatment of 6 weeks increased activation in ROIs of DLPFC, parietal cortex and cerebellum but not in dorsal ACC [85]. A comparative fMRI study using a parallel group design in 36 ADHD patients found that a shared association between clinical improvement after 6-8 weeks of both drugs and reductions in bilateral primary motor cortex activation. There was, however, also a drug-dissociated association in right IFG, left ACC/SMA and bilateral PCC cortex which were all enhanced in activation in relation to clinical response to atomoxetine but decreased in activation in relation to clinical response to methylphenidate [86]. Last, a preliminary analysis found that chronic methylphenidate response in 7 ADHD adolescents was associated with acute stimulant reduction effects in parietal regional homogeneity during rest [87]. Future larger powered imaging studies using multivariate pattern recognition methodology are needed, however, to provide better estimates of whether baseline imaging deficits can predict medication response.

In conclusion, both psychostimulants and atomoxetine appear to enhance the activation of right IFC, presumably via their shared mechanism of action on catecholamines in frontal brain regions, but they seem to also have drug-specific effects in other frontal and subcortical regions. Further studies, however, are needed to disentangle the shared and specific drug effects of atomoxetine relative to stimulant medication on ADHD brain function.

Disorder-specificity of ADHD brain abnormalities compared to other childhood disorders

For neuroimaging to be clinically relevant it is crucial to establish 'disorder-specific' biomarkers that can help with differential diagnosis and differential treatment decisions. The childhood pathology that is most commonly comorbid with ADHD is CD/ODD. However, hardly any imaging studies have controlled for CD/ODD and hence the ADHD im-

aging literature is confounded by this comorbidity. Two sMRI studies compared the two disorders. One found no disorder-specific differences between comorbid children with ADHD with and without CD, but shared volume reductions in posterior and inferior cerebellar vermis [88]. The other study found reduced GM in several frontal and parieto-temporal regions in CD relative to ADHD and control children, while ADHD patients had only reduced regional GM deficit in a left dorsolateral/precentral prefrontal lobe region relative to controls. Also there was a shared GM deficit between both disorders in left dorsolateral/precentral gyri. However, all analyses were conducted at uncorrected thresholds and the ADHD findings were not consistent with the prior literature [89]. A series of fMRI studies from our lab compared well differentiated medication-naïve and IQ-matched groups of children with non-comorbid CD and non-comorbid ADHD during 5 disorder-relevant executive function tasks of response and interference inhibition, sustained attention, saliency detection and cognitive switching. Despite no performance differences, in 4 of the 5 tasks, patients with ADHD had disorder-specific reduced activation compared to both healthy controls and CD patients in the IFC/DLPFC [90-93]. CD children, on the other hand, showed underactivation in paralimbic regions during all tasks, including temporal lobe during performance monitoring [93], the ventromedial prefrontal cortex during reward [90] and limbic areas during sustained attention [90], in line with consistent evidence for structural and function deficits in CD in the paralimbic system, comprising ventromedial OFC and interconnected limbic structures that mediate motivation and affect control [4]. Two studies compared ADHD children with and without CD and psychopathy traits, and found that the comorbid group only had reduced amygdala activation and reduced connectivity between amygdala and vmPFC in relation to fear [94], but enhanced activation in vmPFC during punished reversal errors [95]. Therefore it appears that there are disorder-specific and process-related dissociations in prefrontal lobe deficits, with ADHD children having consistent problems with the recruitment of lateral IFC/DLPFC systems in the context of 'cool' executive inhibitory and attention control, whereas CD children have problems with the recruitment of 'hot' ventromedial OFC-limbic systems that mediate motivation [4].

Disorder-specific reduction in IFC activation in children with ADHD was also observed when compared to children with obsessive-compulsive disorder (OCD) during tasks of motor response inhibition and switching [96,97].

OCD patients, in turn, had shared abnormalities with ADHD patients in the OFC and DLPFC [96,97]. During a saliency task inverse activation patterns were observed with PCC and basal ganglia being underactivated in ADHD relative to OCD, who had increased activation in these regions relative to controls and ADHD boys [97], in line with evidence for diminished saliency processing in ADHD, but enhanced saliency processing in OCD. While no structural studies have directly compared between OCD and ADHD, two meta-analyses of VBM studies in each disorder showed differential abnormalities in the basal ganglia, which were reduced in GM in ADHD [18], but enhanced in OCD [98].

Two sMRI and 2 fMRI studies have compared adolescents with ADHD and autism spectrum disorder (ASD). The first sMRI study found shared abnormalities in limbic and parietal regions which, however, did not survive correction for multiple testing [99]. A study using multivariate pattern recognition analysis found relatively high specificity of about 80% of classifying ADHD patients relative to both controls and ASD patients based on brain patterns of later developing lateral fronto-striato-cerebellar networks for the classification of healthy controls and of earlier developing ventromedial fronto-limbic regions for the classification of ADHD patients [100]. A task-based fMRI study showed shared dysfunction in DLPFC-striato-thalamic regions and shared problems with the deactivation of the DMN in both disorders during a parametric sustained attention task. ASD patients, however, had presumably compensatory increased cerebellar activation and less pronounced DLPFC underfunction relative to ADHD, which may underlie their spared task performance deficits which were only observed in ADHD [21]. A resting state fMRI study reported shared network centrality abnormalities in precuneus for both disorders, but ADHD-specific increases in degree centrality in right striatum/pallidum relative to controls and ASD, while ASD-specific increases in degree centrality were observed in predominantly left temporo-limbic areas, which are typical areas for socio-emotion processing that have consistently been found to be abnormal in ASD [101].

Several imaging studies have compared ADHD with bipolar disorder (BD). Two pediatric sMRI studies found contrasting associations of decreased volumes of caudate and putamen with ADHD and increased volumes of caudate, putamen, globus pallidus and ventral striatum as well as additional lim-

bic abnormalities with BD [102,103]. The comorbid group was either non-impaired [102] or more like the pure BD group [103]. Two adult sMRI found that pure ADHD was associated with reduced GM volumes and cortical thickness in prefrontal, cingulate, parieto-temporal and cerebellar regions, whilst pure BD was associated with smaller left orbital prefrontal and larger right thalamic and parieto-temporal regions [104,105]. In both studies the comorbid disorder showing combinatorial deficits [104,105]. A third study showed that BD alone was associated with thinning in middle, ventral PFC and ACC, while an interaction effect between both disorders was observed in the thinning of left OFC and subgenual ACC [106]. Last, a DTI study testing for differences in 8 fiber tracts reported reduction in 7 WM tracts in ADHD, while BD had shared abnormalities with ADHD in only one of the frontal WM tracts, i.e., the anterior corona radiata and the splenium, connecting posterior attention regions [35]. The findings suggest a global dysmaturation of WM tracts in ADHD, with more specific abnormalities in frontal and parietal WM tracts in BD. Overall, the structural imaging studies show more prominent OFC-limbic abnormalities for BD and more prominent lateral fronto-striato-parieto-temporal deficits in ADHD and differential basal ganglia volumes sizes (enlarged in BD and reduced in ADHD). Similar findings have been observed in fMRI studies. During inhibition tasks, ADHD children showed decreased activation relative to BD in typical inhibition regions of IFG/VLPFC, DLPFC and temporo-parietal regions, whereas BD children had reduced occipital and postcentral activation relative to ADHD [107]. During an emotional valence Stroop task, ADHD patients had reduced VLPFC activation relative to healthy controls while BD relative to healthy controls and ADHD showed increased activation of the VLPFC and ACC [107]. During an affective 2-back working memory task, BD patients had increased activation in bilateral caudate relative to ADHD for happy faces, while for angry faces, the BD group showed increased activation relative to the ADHD group in emotion-regulation areas of left medial OFC and left subgenual ACC, whereas the ADHD group showed increased activation in 'cool' EF working memory regions of DLPFC and SMA [108]. Overall, both the structural and functional comparisons suggest that the emotional impulsiveness of non-comorbid BD is more consistently associated with OFC-limbic abnormalities, while the cognitive impulsivity of ADHD appears to be associated with 'cool' abnormalities in cognitive VLPFC/IFC-striato-parietal circuits [109].

In conclusion, relatively few and small-sampled studies have compared between ADHD and other disorders. There is evidence that ADHD patients have more pronounced dysfunctions in lateral IFC/DLPFC relative to CD, OCD [4], BD [109] and possibly ASD [21]. In addition, there is evidence that the basal ganglia structure and function may be different between ADHD and both BD and OCD [18, 97,98,102,103,108]. Future larger sampled comparisons in groups of non-comorbid and comorbid patients, however, will be necessary to establish disorder-specific structural and functional biomarkers.

Clinical application of neuroimaging in ADHD

Despite consistent evidence for brain structure and function deficits in ADHD, currently ADHD is diagnosed solely on the basis of subjective clinical and rating measures, which are often unreliable. Attempts to find objective neuroimaging biomarkers for ADHD, however, have been limited by the use of traditional univariate group statistical analyses, where subjects in both groups typically overlap and effect sizes are small. In contrast, multivariate pattern analyses for imaging data can make predictions (e.g. of class membership) for individual subjects as opposed to group-level inferences and have been successfully applied to other disorders [110].

To date, few imaging studies have used multivariate pattern recognition analyses techniques to classify ADHD patients. A recent competition to apply multivariate methods on a multicenter resting state functional and anatomical imaging dataset of 285 children and adolescents with ADHD and 491 healthy controls (ADHD-200 Consortium; http://fcon_1000.projects.nitrc.org/indi/adhd200/) was met by a range of classification approaches [111-114]. Accuracies derived by internal cross-validation ranged from 55-78% [114].

A few recent studies used probabilistic classification models such as Gaussian Process Classifiers (GPCs), which provide estimates of predictive uncertainty and can accommodate unbalanced diagnostic settings or variations in disease prevalence, which are crucial for clinical applications [115,116]. Three studies from our lab used GPC in structural and functional imaging data in adolescents with ADHD and showed a relatively high overall classification accuracy of between 75% and 80% with relatively small numbers of between 20 and 33 patients [100,117,118]. Interestingly, across all studies the brain structure and function patterns that classified controls were in later developing lateral

fronto-striato-parieto-cerebellar networks, while patterns that classified ADHD were in earlier developing ventromedial prefrontal and limbic regions [100,117,118]. Importantly, one of the studies showed that the classification accuracy was disorder-specific compared to ASD adolescents [100], which is crucial for the potential future use of neuroimaging as an aid for 'differential' diagnosis. Another recent study used a semi-supervised clustering algorithm based on the spatial patterns of variation across the morphological surfaces of numerous cortical and subcortical brain regions to disorder-specifically diagnose, with relatively high accuracy of almost 90%, youth with ADHD relative to Tourette syndrome [119]. These studies are a promising first step towards the translational use of neuroimaging as a potential differential diagnostic aid. Future studies will have to test whether classification algorithms are stable across patient population, countries and scanners as well as their ability to classify ADHD subgroups and to determine disorder-specific classification. While imaging-based classification algorithms are unlikely to replace clinical assessment and diagnosis, they may be a useful objective, automated, and reliable complementary diagnostic tool that could reduce variability in clinical practice and, ultimately, help to improve diagnostic accuracy. They also have the potential to help with prognosis of disease progression or treatment choice.

Discussion

The last two decades of neuroimaging have significantly broadened our understanding of the underlying neurobiology of ADHD. They have shown that ADHD is most prominently associated with the dysmorphology, dysfunction and underconnectivity of multiple fronto-striato-parietal and fronto-cerebellar networks that mediate 'cool' EF, such as the ventral fronto-ACC/SMA-striato-thalamic cognitive control system, the dorsal and ventral fronto-striato-thalamo-parieto-cerebellar attention systems and ventral fronto-striato-parieto-cerebellar timing networks. In addition, ADHD children have structural, functional and connectivity deficits in DMN systems that appear to be poorly deactivated during task performance and hence to intrude upon task-positive cognitive systems, leading to impaired EF. Furthermore, there is emerging evidence for OFC-limbic abnormalities in the context of reward processing, with particular implication for the ventral striatum, which, however, needs to be con-

firmed in larger-scale imaging studies, given that the findings are based on ROI analyses and confounded by comorbidity with CD. There is evidence for a delay in normal brain maturation from structural imaging studies, with indications that this may also apply to brain function and structural and functional connectivity. Studies on the clinically relevant question of disorder-specificity of imaging biomarkers for ADHD are only just emerging, but point towards potential disorder-specificity of ventral prefrontal under-recruitment relative to CD, OCD and BD [4,109] as well as of differential basal ganglia deficits relative to OCD and BD [18,109].

fMRI studies have shown that the most consistent mechanism of action of psychostimulants in ADHD is the upregulation and normalisation of the activation of the IFC and the basal ganglia. Studies on atomoxetine effects on brain function have only just emerged, but acute effects on prefrontal systems appear to be similar to those of psychostimulant medication. Some evidence from retrospective comparisons suggests long-term plastic medication effects on brain structure, function and neurochemistry, which, however, need to be confirmed in prospective randomized controlled designs. PET studies will need to investigate the underlying neurotransmitter abnormalities in ADHD other than the DA system, such as the serotonin, noradrenalin, glutamate and GABA systems with a particular emphasis on how they interact with the dysfunctional DA system.

The imaging literature of ADHD is confounded by relatively small numbers of often non-representative samples of convenience, by the use of ROI analyses and liberal uncorrected thresholds and by uncontrolled comorbidity with CD/ODD. Adult studies suffer from comorbidity confounds with secondary affective conditions and ascertainment bias from clinics, which do not capture remitted cases. The majority of imaging studies have been conducted in long-term medicated patients, which, as discussed, has an effect on structure and function, and is hence a crucial confound.

Given that cross-sectional imaging studies are confounded by cohort effects and ascertainment bias, there is an urgent need for longitudinal imaging studies, in particular following up children with ADHD into adulthood, to understand brain mechanisms of remittance and persistence. Furthermore, future studies will need to focus on multimodal imaging in representative populations to enhance our holistic understanding on the relationship between structural, functional, connectivity, blood flow as well as brain chemistry abnormalities within the

same patient groups. Also, future studies will need to focus on understanding the (differential) neurobiological basis of different ADHD subtypes as well as comorbid cases. Future studies therefore ideally should be longitudinal, multimodal, and tied to epidemiological samples.

Clinical translation of neuroimaging will be the challenge over the next decade. Several pioneering machine learning approaches using GPC have been promising, showing relatively high accuracy of up to 80% in classifying ADHD patients relative to controls and ASD patients based on structural or functional imaging scans. Multimodal multivariate approaches including several imaging as well as non-imaging modalities may provide higher classification accuracies than unimodal studies [110]. These multivariate classification methods, if successful and replicated across different representative patient groups, scanners and demographic populations, may be able to help with future imaging-based diagnosis or prognosis of individual patients and provide brain-based patient stratification and more personalised medicine.

Last, there is high potential for using neuroimaging as a neurotherapy. EEG-based neurofeedback has been shown to have similar effect sizes to stimulant medication in reducing ADHD behaviours [120]. fMRI or near infrared-spectroscopy (NIRS) have better spatial resolution and combined with neurofeedback could be used to upregulate the activation of ADHD-specific dysfunctional brain regions such as IFC, DLPFC and the basal ganglia (with fMRI only). Likewise, regional electrical stimulation via repetitive transcranial magnetic stimulation (rTMS) and transcranial direct current stimulation (tDCS) have been successful in other psychiatric disorders [121] and are promising for ADHD. So far, only one pilot study tested rTMS over right dorsolateral prefrontal cortex in one single session in 13 ADHD adults and found increases in behavioural attention scores [122].

In conclusion, we have acquired substantial knowledge on the underlying neurobiological mechanisms of ADHD. However, more studies are needed that integrate different imaging modalities to understand the interplay between the changes in neurochemistry, brain function and brain structure and to assess longitudinal trajectories of the disorder. The next decade will need to focus on using neuroimaging techniques in a more clinically applied fashion, either to aid with individual diagnosis, prognosis of disease progression and treatment success or as a neurotherapy to normalise abnormally functioning brain regions.

References

1. American Psychiatric Association. Diagnostic and statistical manual of mental disorders, fourth edition. Washington DC: APA; 2000.
2. Biederman J, Petty CR, Woodworth KY, Lomedico A, Hyder LL, Faraone SV. Adult outcome of attention-deficit/hyperactivity disorder: a controlled 16-year follow-up study. *J Clin Psychiatry* 2012; 73: 941-50.
3. Rubia K. Functional neuroimaging across development: a review. *Eur Child Adolesc Psychiatry* 2013; 22: 719-31.
4. Rubia K. 'Cool' inferior fronto-striatal dysfunction in attention deficit hyperactivity disorder (adhd) versus 'hot' ventromedial orbitofronto-limbic dysfunction in conduct disorder: a review. *Biol Psychiatry* 2011; 69: e69-87.
5. Willcutt EG, Sonuga-Barke EJS, Nigg JT, Sergeant GA. Recent developments in neuropsychological models of childhood psychiatric disorders. In Banaschewski T, Rohde LA, eds. *Biological child psychiatry. Recent trends and developments*. Basel: Karger; 2008. p. 195-226.
6. Rubia K, Halari R, Christakou A, Taylor E. Impulsiveness as a timing disturbance: neurocognitive abnormalities in attention-deficit hyperactivity disorder during temporal processes and normalization with methylphenidate. *Philos Trans R Soc Lond B Biol Sci* 2009; 364: 1919-31.
7. Noreika V, Falter C, Rubia K. Timing deficits in patients with ADHD. *Neuropsychologia* 2012; 51: 235-66.
8. Valera EM, Faraone SV, Murray KE, Seidman LJ. Meta-analysis of structural imaging findings in attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2007; 61: 1361-9.
9. Hoekzema E, Carmona S, Ramos-Quiroga JA, Richarte-Fernández V, Picado M, Bosch R, et al. Laminar thickness alterations in the fronto-parietal cortical mantle of patients with attention-deficit/hyperactivity disorder. *PLoS One* 2012; 7: e48286.
10. Shaw P, Malek M, Watson B, Sharp W, Evans A, Greenstein D. Development of Cortical surface area and gyrification in attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2012; 72: 191-7.
11. Shaw P, Malek M, Watson B, Greenstein D, De Rossi P, Sharp W. Trajectories of Cerebral cortical development in childhood and adolescence and adult attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2013; 74: 599-606.
12. Cubillo A, Halari R, Smith A, Giampietro V, Taylor E, Rubia K. Fronto-cortical and fronto-subcortical brain abnormalities in children and adults with ADHD: a review and evidence for fronto-striatal dysfunctions in adults with ADHD followed up from childhood during motivation and attention. *Cortex* 2012; 48: 194-215.
13. Lopez-Larson MP, King JB, Terry J, McGlade EC, Yurgelun-Todd D. Reduced insular volume in attention deficit hyperactivity disorder. *Psychiatr Res* 2012; 204: 32-9.
14. Plessen KJ, Bansal R, Zhu HT, Whiteman R, Amat J, Quackenbush GA, et al. Hippocampus and amygdala morphology in attention-deficit/hyperactivity disorder. *Arch Gen Psychiatry* 2006; 63: 795-807.
15. Ivanov I, Bansal R, Hao XJ, Zhu HT, Kellendonk C, Miller L, et al. Morphological abnormalities of the thalamus in youths with attention deficit hyperactivity disorder. *Am J Psychiatry* 2010; 167: 397-408.
16. Arnsten A, Rubia K. Neurobiological circuits regulating attention, movement and emotion and their disruptions in pediatric neuropsychiatric disorders. *J Am Acad Child Adolesc Psychiatry* 2012; 51: 356-67.
17. Ellison-Wright I, Ellison-Wright Z, Bullmore E. Structural brain change in attention deficit hyperactivity disorder identified by meta-analysis. *BMC Psychiatry* 2008; 8: 51.
18. Nakao T, Radua C, Rubia K, Mataix-Cols D. Gray matter volume abnormalities in ADHD and the effects of stimulant medication: voxel-based meta-analysis. *Am J Psychiatry* 2011; 168: 1154-63.
19. Frodl T, Skokauskas N. Meta-analysis of structural MRI studies in children and adults with attention deficit hyperactivity disorder indicates treatment effects. *Acta Psychiatr Scand* 2012; 125: 114-26.

20. Fusar-Poli P, Rubia K, Rossi G, Sartori G, Ballotín U. Dopamine transporter alterations in ADHD: pathophysiology or adaptation to psychostimulants? A meta-analysis. *Am J Psychiatry* 2012; 169: 264-72.
21. Christakou A, Murphy C, Chantiluke C, Cubillo A, Smith A, Giampietro V, et al. Disorder-specific functional abnormalities during sustained attention in youth with attention deficit hyperactivity disorder and with autism. *Mol Psychiatry* 2013; 18: 236-44.
22. Hart H, Radua J, Mataix D, Rubia K. Meta-analysis of fMRI studies of timing functions in ADHD. *Neurosci Biobehav Rev* 2012; 36: 2248-56.
23. Broyd SJ, Demanuele C, Debener S, Helps SK, James CJ, Sonuga-Barke EJS. Default-mode brain dysfunction in mental disorders: a systematic review. *Neurosci Biobehav Rev* 2009; 33: 279-96.
24. Shaw P, Eckstrand K, Sharp W, Blumenthal J, Lerch JP, Greenstein D, et al. Attention-deficit/hyperactivity disorder is characterized by a delay in cortical maturation. *Proc Natl Acad Sci U S A* 2007; 104: 19649-54.
25. Cortese S, Imperati D, Zhou J, Proal E, Klein RG, Mannuzza S, et al. White matter alterations at 33-year follow-up in adults with childhood attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2013; 74: 591-8.
26. Konrad A, Dielentheis TF, El Masri D, Bayerl M, Fehr C, Gesierich T, et al. Disturbed structural connectivity is related to inattention and impulsivity in adult attention deficit hyperactivity disorder. *Eur J Neurosci* 2010; 31: 912-9.
27. Konrad K, Eickhoff SB. Is the ADHD brain wired differently? A review on structural and functional connectivity in attention deficit hyperactivity disorder. *Hum Brain Mapp* 2010; 31: 904-16.
28. Dramsdaahl M, Westerhausen R, Haavik J, Hugdahl K, Plessen KJ. Adults with attention-deficit/hyperactivity disorder – a diffusion-tensor imaging study of the corpus callosum. *Psychiatry Res* 2012; 201: 168-73.
29. Konrad A, Dielentheis TF, El Masri D, Dellani PR, Stoeter P, Vucurevic G, et al. White matter abnormalities and their impact on attentional performance in adult attention-deficit/hyperactivity disorder. *Eur Arch Psychiatry Clin Neurosci* 2012; 262: 351-60.
30. Cao Q, Shu N, An L, Wang P, Sun L, Xia MR, et al. Probabilistic diffusion tractography and graph theory analysis reveal abnormal white matter structural connectivity networks in drug-naïve boys with attention deficit/hyperactivity disorder. *J Neurosci* 2013; 33: 10676-87.
31. Van Ewijk H, Heslenfeld DJ, Zwiers MP, Buitelaar JK, Oosterlaan J. Diffusion tensor imaging in attention deficit/hyperactivity disorder: a systematic review and meta-analysis. *Neurosci Biobehav Rev* 2012; 36: 1093-106.
32. Chuang TC, Wu MT, Huang SP, Weng MJ, Yang P. Diffusion tensor imaging study of white matter fiber tracts in adolescent attention-deficit/hyperactivity disorder. *Psychiatry Res* 2013; 211: 186-7.
33. Silk T, Vance A, Rinehart N, Bradshaw J, Cunnington R. Diffusion tensor imaging in attention deficit hyperactivity disorder. *J Neural Transm* 2007; 114: Lxxii-Lxxii.
34. Yap QJ, Teh I, Fusar-Poli P, Sum MY, Kuswanto C, Sim K. Tracking cerebral white matter changes across the lifespan: insights from diffusion tensor imaging studies. *J Neural Transm* 2013; 120: 1369-95.
35. Pavuluri MN, Yang S, Kamini K, Passarotti AM, Srinivasan G, Harral EM, et al. Diffusion tensor imaging study of white matter fiber tracts in pediatric bipolar disorder and attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2009; 65: 586-93.
36. Nagel BJ, Bathula D, Herting M, Schmitt C, Kroenke CD, Fair D, et al. Altered white matter microstructure in children with attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2011; 50: 283-92.
37. Cortese S, Kelly C, Chabernaud C, Proal E, Di Martino A, Milham MP, et al. Toward systems neuroscience of ADHD: a meta-analysis of 55 fMRI studies. *Am J Psychiatry* 2012; 169: 1038-55.
38. Hart H, Radua J, Mataix D, Rubia K. Meta-analysis of fMRI studies of inhibition and attention in ADHD: exploring task-specific, stimulant medication and age effects. *JAMA Psychiatry* 2013; 70: 185-98.
39. Wiener M, Turkeltaub P, Coslett HB. The image of time: a voxel-wise meta-analysis. *Neuroimage* 2010; 49: 1728-40.
40. Plichta MM, Scheres A. Ventral-striatal responsiveness during reward anticipation in ADHD and its relation to trait impulsivity in the healthy population: a meta-analytic review of the fMRI literature. *Neurosci Biobehav Rev* 2014; 38: 125-34.
41. Castellanos FX, Proal E. Large-scale brain systems in ADHD: beyond the prefrontal-striatal model. *Trends Cogn Sci* 2012; 16: 17-26.
42. Wang L, Zhu CZ, He Y, Zang YF, Cao QJ, Zhang H, et al. Altered small-world brain functional networks in children with attention-deficit/hyperactivity disorder. *Hum Brain Mapp* 2009; 30: 638-49.
43. Sun L, Cao Q, Long X, Sui M, Cao X, Zhu C, et al. Abnormal functional connectivity between the anterior cingulate and the default mode network in drug-naïve boys with attention deficit hyperactivity disorder. *Psychiatry Res* 2012; 201: 120-7.
44. Fair DA, Posner J, Nagel BJ, Bathula D, Dias TGC, Mills KL, et al. Atypical default network connectivity in youth with attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2010; 68: 1084-91.
45. Uddin LQ, Kelly AMC, Biswal BB, Margulies DS, Shehzad Z, Shaw D, et al. Network homogeneity reveals decreased integrity of default-mode network in ADHD. *J Neurosci Methods* 2008; 169: 249-54.
46. Cao XH, Cao QJ, Long XY, Sun L, Sui MQ, Zhu CZ, et al. Abnormal resting-state functional connectivity patterns of the putamen in medication-naïve children with attention deficit hyperactivity disorder. *Brain Res* 2009; 1303: 195-206.
47. Sato JR, Takahashi DY, Hoexter MQ, Massier KB, Fujita A. Measuring network's entropy in ADHD: a new approach to investigate neuropsychiatric disorders. *Neuroimage* 2013; 77: 44-51.
48. Cocchi L, Bramati IE, Zalesky A, Furukawa E, Fontenelle LF, Moll J, et al. Altered Functional brain connectivity in a non-clinical sample of young adults with attention-deficit/hyperactivity disorder. *J Neurosci* 2012; 32: 17753-61.
49. Fair DA, Nigg JT, Iyer S, Bathula D, Mills KL, Dosenbach NUF, et al. Distinct neural signatures detected for ADHD subtypes after controlling for micro-movements in resting state functional connectivity MRI data. *Front Syst Neurosci* 2012; 6: 80.
50. Posner J, Rauh V, Gruber A, Gat I, Wang Z, Peterson BS. Dissociable attentional and affective circuits in medication-naïve children with attention-deficit/hyperactivity disorder. *Psychiatry Res* 2013; 213: 24-30.
51. Rubia K, Halari R, Cubillo A, Mohammad M, Taylor E. Methylphenidate normalises activation and functional connectivity deficits in attention and motivation networks in medication-naïve children with ADHD during a rewarded Continuous Performance Task. *Neuropharmacology* 2009; 5: 640-52.
52. Vloet TD, Gilsbach S, Neufang S, Fink GR, Herpertz-Dahlmann B, Konrad K. Neural mechanisms of interference control and time discrimination in attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2010; 49: 356-67.
53. Cubillo A, Halari R, Ecker C, Giampietro V, Taylor E, Rubia K. Reduced activation and inter-regional functional connectivity of fronto-striatal networks in adults with childhood attention deficit hyperactivity disorder (ADHD) and persisting symptoms during tasks of motor inhibition and cognitive switching. *J Psychiatr Res* 2010; 44: 629-39.
54. Peterson BS, Potenza MN, Wang Z, Zhu H, Martin A, Marsh R, et al. An fMRI study of the effects of psychostimulants on default-mode processing during Stroop task performance in youths with ADHD. *Am J Psychiatry* 2009; 166: 1286-94.
55. Wolf RC, Plichta MM, Sambataro F, Fallgatter AJ, Jacob C, Lesch KP, et al. Regional brain activation changes and

- abnormal functional connectivity of the ventrolateral prefrontal cortex during working memory processing in adults with attention-deficit/hyperactivity disorder. *Hum Brain Mapp* 2009; 30: 2252-66.
56. Fassbender C, Zhang H, Buzy WM, Cortes CR, Mizuiri D, Beckett L, et al. A lack of default network suppression is linked to increased distractibility in ADHD. *Brain Res* 2009; 1273: 114-28.
 57. Clerkin SM, Schulz KP, Berwid OG, Fan J, Newcorn JH, Tang CY, et al. Thalamo-cortical activation and connectivity during response preparation in adults with persistent and remitted ADHD. *Am J Psychiatry* 2013; 170: 1011-9.
 58. Hoekzema E, Carmona S, Ramos-Quiroga JA, Richarte-Fernandez V, Bosch R, Soliva JC, et al. An independent components and functional connectivity analysis of resting state fMRI data points to neural network dysregulation in adult ADHD. *Hum Brain Mapp* 2013; Feb 18. [Epub ahead of print].
 59. Dosenbach NUF. Prediction of individual brain maturity using fMRI. *Science* 2010; 329: 1358.
 60. Sato JR, Hoexter MQ, Castellanos XF, Rohde LA. Abnormal brain connectivity patterns in adults with ADHD: a coherence study. *PLoS One* 2012; 7: e45671.
 61. Wilens TE. Effects of methylphenidate on the catechol-aminergic system in attention-deficit/hyperactivity disorder. *J Clin Psychopharmacol* 2008; 28 (Suppl 2): S46-53.
 62. Cubillo A, Smith A, Barrett N, Giampetro V, Brammer MJ, Simmons A, et al. Shared and drug-specific effects of atomoxetine and methylphenidate on inhibitory brain dysfunction in medication-naïve ADHD boys. *Cereb Cortex* 2014; 24: 174-85.
 63. Rubia K, Halari R, Cubillo A, Smith A, Mohammad M, Brammer M, et al. Methylphenidate normalises fronto-striatal underactivation during interference inhibition in medication-naïve boys. *Neuropsychopharmacology* 2011; 36: 1575-86.
 64. Rubia K, Halari R, Taylor E, Brammer M. Methylphenidate normalises fronto-cingulate underactivation during error processing in children with attention-deficit hyperactivity disorder. *Biol Psychiatry* 2011; 70: 255-62.
 65. Rubia K, Alzamora A, Cubillo A, Smith AB, Radua J, Brammer MJ. Effects of stimulants on brain function in ADHD: a systematic review and meta-analysis. *Biol Psychiatry* 2013; Oct 24. [Epub ahead of print].
 66. Epstein JN, Casey BJ, Tonev ST, Davidson MC, Reiss AL, Garrett A, et al. ADHD- and medication-related brain activation effects in concordantly affected parent-child dyads with ADHD. *J Child Psychol Psychiatry* 2007; 48: 899-913.
 67. Smith A, Cubillo A, Barrett N, Giampietro V, Simmons A, Brammer M, et al. Neurofunctional effects of methylphenidate and atomoxetine in boys with ADHD during time discrimination. *Biol Psychiatry* 2013; 74: 615-22.
 68. Cubillo A, Smith A, Barrat N, Giampietro V, Simmons A, Brammer M, et al. Drug-specific laterality effects on frontal lobe activation of atomoxetine and methylphenidate in ADHD boys during working memory. *Psychol Med* 2013; 19: 1-14.
 69. Kobel M, Bechtel N, Weber P, Specht K, Klarhofer M, Scheffler K, et al. Effects of methylphenidate on working memory functioning in children with attention deficit/hyperactivity disorder. *Eur J Paediatr Neurol* 2009; 13: 516-23.
 70. Vaidya CJ, Austin G, Kirkorian G, Ridlehuber HW, Desmond JE, Glover GH, et al. Selective effects of methylphenidate in attention deficit hyperactivity disorder: a functional magnetic resonance study. *Proc Natl Acad Sci U S A* 1998; 95: 14494-9.
 71. Posner J, Maia TV, Fair D, Peterson BS, Sonuga-Barke EJ, Nagel BJ. The attenuation of dysfunctional emotional processing with stimulant medication: an fMRI study of adolescents with ADHD. *Psychiatry Res* 2011; 193: 151-60.
 72. Konrad K, Neufang S, Fink GR, Herpertz-Dahlmann B. Long-term effects of methylphenidate on neural networks associated with executive attention in children with ADHD: results from a longitudinal functional MRI study. *J Am Acad Child Adolesc Psychiatry* 2007; 46: 1633-41.
 73. Mizuno K, Yoneda T, Komi M, Hirai T, Watanabe Y, Tomoda A. Osmotic release oral system-methylphenidate improves neural activity during low reward processing in children and adolescents with attention-deficit/hyperactivity disorder. *Neuroimage Clin* 2013; 2: 366-76.
 74. Bush G, Spencer TJ, Holmes J, Shin LM, Valera EM, Seidman LJ, et al. Functional magnetic resonance imaging of methylphenidate and placebo in attention-deficit/hyperactivity disorder during the multi-source interference task. *Arch Gen Psychiatry* 2008; 65: 102-14.
 75. Semrud-Clikeman M, Pliszka SR, Lancaster J, Liotti M. Volumetric MRI differences in treatment-naïve vs chronically treated children with ADHD. *Neurology* 2006; 67: 2091.
 76. Bledsoe J, Semrud-Clikeman M, Pliszka SR. A magnetic resonance imaging study of the cerebellar vermis in chronically treated and treatment-naïve children with attention-deficit/hyperactivity disorder combined type. *Biol Psychiatry* 2009; 65: 620-4.
 77. Ivanov I, Murrough JW, Bansal R, Hao X, Peterson BS. Cerebellar morphology and the effects of stimulant medications in youths with attention deficit-hyperactivity disorder. *Neuropsychopharmacology* 2014; 39: 718-26.
 78. Sobel LJ, Bansal R, Maia TV, Sanchez J, Mazzone L, Durkin K, et al. Basal ganglia surface morphology and the effects of stimulant medications in youth with attention deficit hyperactivity disorder. *Am J Psychiatry* 2010; 167: 977-86.
 79. Schnoebelen S, Semrud-Clikeman M, Pliszka SR. Corpus callosum anatomy in chronically treated and stimulant naïve ADHD. *J Atten Disord* 2010; 14: 256-66.
 80. Castellanos FX, Lee PP, Sharp W, Jeffries NO, Greenstein DK, Clasen LS, et al. Developmental trajectories of brain volume abnormalities in children and adolescents with attention-deficit/hyperactivity disorder. *JAMA* 2002; 288: 1740-8.
 81. Shaw P, Sharp WS, Morrison M, Eckstrand K, Greenstein DK, Clasen LS, et al. Psychostimulant treatment and the developing cortex in attention deficit hyperactivity disorder. *Am J Psychiatry* 2009; 166: 58-63.
 82. Bledsoe JC, Semrud-Clikeman M, Pliszka SR. Anterior cingulate cortex and symptom severity in attention-deficit/hyperactivity disorder. *J Abnorm Psychol* 2013; 122: 558-65.
 83. Volkow ND, Wang GJ, Tomasi D, Kollins SH, Wigal TL, Newcorn JH, et al. Methylphenidate-elicited dopamine increases in ventral striatum are associated with long-term symptom improvement in adults with attention deficit hyperactivity disorder. *J Neurosci* 2012; 32: 841-9.
 84. Hazell P. The challenges to demonstrating long-term effects of psychostimulant treatment for attention-deficit/hyperactivity disorder. *Curr Opin Psychiatry* 2011; 24: 286-90.
 85. Bush G, Holmes J, Shin LM, Surman C, Makris N, Mick E, et al. Atomoxetine increases fronto-parietal functional MRI activation in attention-deficit/hyperactivity disorder: a pilot study. *Psychiatry Res* 2013; 211: 88-91.
 86. Schulz KP, Fan J, Bedard ACV, Clerkin SM, Ivanov I, Tang CY, et al. Common and unique therapeutic mechanisms of stimulant and nonstimulant treatments for attention-deficit/hyperactivity disorder. *Arch Gen Psychiatry* 2012; 69: 952-61.
 87. Li A, Cao XH, Cao Q, Sun L, Yang L, Zou Q, et al. Methylphenidate normalizes resting-state brain dysfunction in boys with attention deficit hyperactivity disorder. *Neuropsychopharmacology* 2013; 38: 1287-95.
 88. Bussing R, Grudnik J, Mason D, Wasiak M, Leonard C. ADHD and conduct disorder: an MRI study in a community sample. *World J Biol Psychiatry* 2002; 3: 216-20.
 89. Stevens MC, Haney-Caron E. Comparison of brain volume abnormalities between ADHD and conduct disorder in adolescence. *J Psychiatry Neurosci* 2012; 37: 389-98.
 90. Rubia K, Smith A, Halari R, Matukura F, Mohammad M, Taylor E, et al. Disorder-specific dissociation of orbitofrontal dysfunction in boys with pure conduct disorder during reward and ventrolateral prefrontal dysfunction in boys with pure attention-deficit/hyperactivity disorder during sustained attention. *Am J Psychiatry* 2009; 166: 83-94.
 91. Rubia K, Halari R, Cubillo A, Mohammad A, Scott S, Brammer M. Disorder-specific inferior frontal dysfunction in boys with pure attention-deficit/hyperactivity disorder

- compared to boys with pure CD during cognitive flexibility. *Hum Brain Mapp* 2010; 31: 1823-33.
92. Rubia K, Halari R, Smith AB, Mohammad M, Scott S, Brammer MJ. Shared and disorder-specific prefrontal abnormalities in boys with pure attention-deficit/hyperactivity disorder compared to boys with pure CD during interference inhibition and attention allocation. *J Child Psychol Psychiatry* 2009; 50: 669-78.
 93. Rubia K, Halari R, Smith AB, Mohammed M, Scott S, Giampietro V, et al. Dissociated functional brain abnormalities of inhibition in boys with pure conduct disorder and in boys with pure attention deficit hyperactivity disorder. *Am J Psychiatry* 2008; 165: 889-97.
 94. Marsh AA, Finger EC, Michell DGV, Sims C, Kosson DS, Towbin KE, et al. Reduced amygdala response to fearful expressions in children and adolescents with callous-unemotional traits and disruptive behaviour disorders. *Am J Psychiatry* 2008; 165: 712-20.
 95. Finger EC, Marsh AA, Mitchell DG, Reid ME, Sims C, Budhani S, et al. Abnormal ventromedial prefrontal cortex function in children with psychopathic traits during reversal learning. *Arch Gen Psychiatry* 2008; 65: 586-94.
 96. Rubia K, Cubillo A, Smith AB, Woolley J, Heyman I, Brammer MJ. Disorder-specific dysfunction in right inferior prefrontal cortex during two inhibition tasks in boys with attention-deficit hyperactivity disorder compared to boys with obsessive-compulsive disorder. *Hum Brain Mapp* 2010; 31: 287-99.
 97. Rubia K, Cubillo A, Woolley J, Brammer MJ, Smith AB. Disorder-specific dysfunctions in patients with attention-deficit/hyperactivity disorder compared to patients with obsessive-compulsive disorder during interference inhibition and attention allocation. *Hum Brain Mapp* 2011; 32: 601-11.
 98. Radua J, Mataix-Cols D. Voxel-wise meta-analysis of grey matter changes in obsessive-compulsive disorder. *Br J Psychiatry* 2009; 195: 393-402.
 99. Brieber S, Neufang S, Bruning N, Kamp-Becker I, Remschmidt H, Herpertz-Dahlmann B, et al. Structural brain abnormalities in adolescents with autism spectrum disorder and patients with attention deficit/hyperactivity disorder. *J Child Psychol Psychiatry* 2007; 48: 1251-8.
 100. Lim L, Cubillo A, Smith A, Chantiluke K, Marquand A, Simmons A, et al. Disorder-specific predictive classification of adolescents with attention deficit hyperactivity disorder (ADHD) relative to autism using structural magnetic resonance imaging. *PLOS One* 2013; 8: e63660.
 101. Di Martino A, Zuo XN, Kelly C, Grzadzinski R, Mennes M, Schwarcz A, et al. Shared and distinct intrinsic functional network centrality in autism and attention-deficit/hyperactivity disorder. *Biol Psychiatry* 2013; 74: 623-32.
 102. Liu IY, Howe M, Garrett A, Karchemskiy A, Kelley R, Alegria D, et al. Striatal volumes in pediatric bipolar patients with and without comorbid ADHD. *Psychiatry Res* 2011; 194: 14-20.
 103. Lopez-Larson M, Michael ES, Terry JE, Breeze JL, Hodge SM, Tang L, et al. Subcortical differences among youths with attention-deficit/hyperactivity disorder compared to those with bipolar disorder with and without attention-deficit/hyperactivity disorder. *J Child Adolesc Psychopharmacol* 2009; 19: 31-9.
 104. Biederman J, Makris N, Valera EM, Monuteaux MC, Goldstein JM, Buka S, et al. Towards further understanding of the co-morbidity between attention deficit hyperactivity disorder and bipolar disorder: a MRI study of brain volumes. *Psychol Med* 2008; 38: 1045-56.
 105. Makris N, Seidman LJ, Brown A, Valera EM, Kaiser JR, Petty CR, et al. Further understanding of the comorbidity between attention-deficit/hyperactivity disorder and bipolar disorder in adults: an MRI study of cortical thickness. *Psychiatry Res* 2012; 202: 1-11.
 106. Hegarty CE, Foland-Ross LC, Narr KL, Sugar CA, McGough JJ, Thompson PM, et al. ADHD comorbidity can matter when assessing cortical thickness abnormalities in patients with bipolar disorder. *Bipolar Disord* 2012; 14: 843-55.
 107. Passarotti AM, Sweeney JA, Pavuluri MN. Neural correlates of response inhibition in pediatric bipolar disorder and attention deficit hyperactivity disorder. *Psychiatry Res* 2010; 181: 36-43.
 108. Passarotti AM, Sweeney JA, Pavuluri MN. Emotion processing influences working memory circuits in pediatric bipolar disorder and attention-deficit/hyperactivity disorder. *J Am Acad Child Adolesc Psychiatry* 2010; 49: 1064-80.
 109. Passarotti AM, Pavuluri MN. Brain functional domains inform therapeutic interventions in attention-deficit/hyperactivity disorder and pediatric bipolar disorder. *Exp Rev Neurother* 2011; 11: 897-914.
 110. Orru G, Pettersson-Yeo W, Marquand AF, Sartori G, Mechelli A. Using Support Vector Machine to identify imaging biomarkers of neurological and psychiatric disease: a critical review. *Neurosci Biobehav Rev* 2012; 36: 1140-52.
 111. Cheng W, Ji X, Zhang J, Feng J. Individual classification of ADHD patients by integrating multiscale neuroimaging markers and advanced pattern recognition techniques. *Front Syst Neurosci* 2012; 6: 58.
 112. Colby JB, Rudie JD, Brown JA, Douglas PK, Cohen MS, Shehzad Z. Insights into multimodal imaging classification of ADHD. *Front Syst Neurosci* 2012; 6: 59.
 113. Dai D, Wang J, Hua J, He H. Classification of ADHD children through multimodal magnetic resonance imaging. *Front Syst Neurosci* 2012; 6: 63.
 114. Eloyan A, Muschelli J, Nebel MB, Liu H, Han F, Zhao T, et al. Automated diagnoses of attention deficit hyperactive disorder using magnetic resonance imaging. *Front Syst Neurosci* 2012; 6: 61.
 115. Marquand A, De Simoni S, O'Daly O, Mourao-Miranda J, Mehta M. Quantifying the information content of brain voxels using target information, Gaussian processes and recursive feature elimination. *International Conference on Pattern Recognition. Istanbul, Turkey*; 2010.
 116. Hahn T, Marquand AF, Plichta MM, Ehlis AC, Schecklmann MW, Dresler T, et al. A novel approach to probabilistic biomarker-based classification using functional near-infrared spectroscopy. *Hum Brain Map* 2013; 34: 1102-14.
 117. Hart H, Chantiluke K, Cubillo A, Smith A, Simmons A, Marquand A, et al. Pattern classification of response inhibition in ADHD: toward the development of neurobiological markers for ADHD. *Hum Brain Mapp*; Oct 11. [Epub ahead of print].
 118. Hart H, Smith A, Cubillo A, Simmons A, Marquand A, Rubia K. Predictive neurofunctional markers of ADHD based on pattern classification of temporal processing. *J Am Acad Child Adolesc Psychiatry* 2014 [in press].
 119. Bansal R, Staib LH, Laine AF, Hao XJ, Xu DR, Liu J, et al. Anatomical brain images alone can accurately diagnose chronic neuropsychiatric illnesses. *PLoS One* 2012; 7: e50698.
 120. Arns M, De Ridder S, Strehl U, Breteler M, Coenen A. Efficacy of neurofeedback treatment in ADHD: the effects on inattention, impulsivity and hyperactivity: a meta-analysis. *Clin EEG Neurosci* 2009; 40: 180-9.
 121. Demirtas-Tatlidede A, Vahabzadeh-Hagh AM, Pascual-Leone A. Can noninvasive brain stimulation enhance cognition in neuropsychiatric disorders? *Neuropharmacology* 2013; 64: 566-78.
 122. Bloch Y, Harel EV, Aviram S, Govezensky J, Ratzoni G, Levkovitz Y. Positive effects of repetitive transcranial magnetic stimulation on attention in ADHD subjects: a randomized controlled pilot study. *World J Biol Psychiatry* 2010; 11: 755-8.

Anomalías cerebrales en el trastorno por déficit de atención/hiperactividad: una revisión

Objetivo. Revisar los hallazgos de los estudios con resonancia magnética en el trastorno por déficit de atención/hiperactividad (TDAH) infantil y adulto.

Desarrollo. Dichos estudios han demostrado que el TDAH se caracteriza por la presencia de múltiples anomalías de carácter estructural y funcional, primordialmente en los circuitos frontoestriatales, pero también en los circuitos frontoparieto-temporales, frontocerebelares e, incluso, frontolímbicos. Los datos aportados por los estudios longitudinales de resonancia magnética estructural demuestran que el TDAH se caracteriza por un retraso en la maduración estructural del cerebro. Esta conclusión se ve reforzada por los indicios indirectos ofrecidos por los estudios de cortes transversales, que indican la existencia de una inmadurez sustancial tanto en la función cerebral como en los patrones de conectividad estructural y funcional, indicios que, sin embargo, están pendientes de confirmar en estudios longitudinales. La alteración funcional de la corteza prefrontal ventrolateral parece estar más afectada en el TDAH que en otros trastornos pediátricos, y existen algunos indicios de anomalías distintivas en los ganglios basales. Un metaanálisis sobre los efectos de los estimulantes en la función cerebral demuestra que el mecanismo de acción agudo más congruente de los fármacos psicoestimulantes consiste en el aumento de la activación de la corteza prefrontal inferior y los ganglios basales. Los primeros intentos por utilizar los datos de los estudios de neuroimagen para elaborar clasificaciones diagnósticas individuales de los niños con TDAH a partir de técnicas de reconocimiento de patrones han cosechado resultados alentadores, pero todavía deben ser replicados por más centros y aparatos de resonancia magnética.

Conclusiones. Durante los últimos 20 años, las técnicas de neuroimagen han perfilado los biomarcadores del TDAH, pero es necesario que nuevos estudios descubran la utilidad clínica de esa información, como el uso de tales técnicas como instrumento de clasificación diagnóstica y pronóstica individualizada o como terapia para revertir las anomalías de la función cerebral que han sido confirmadas durante los dos decenios anteriores.

Palabras clave. Atomoxetina. Imágenes con tensor de difusión (DTI). Metilfenidato. Psicoestimulantes. Resonancia magnética funcional (RMf). Trastorno por déficit de atención/hiperactividad (TDAH).