



Ishikawa-Collocation Method for Nonlinear Fredholm Equations with Non-Separable Kernels

Chinedu Nwaigwe ^{a*} and Azubuike Weli ^a

^aDepartment of Mathematics, Rivers State University, Port Harcourt, Nigeria.

Author's contribution

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMCS/2023/v38i31747

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/96672>

Received: 25/11/2022

Accepted: 28/01/2023

Published: 10/02/2023

Original Research Article

Abstract

A fixed point method is developed on a mesh for the solution of nonlinear Fredholm equation. First, the problem is collocated at mesh points and a second order quadrature rule is used to approximate the nonlinear integral. Under the assumption of nonexpansivity of self-map, we construct an Ishikawa iteration to linearize the resulting system and approximate the solution at the mesh points. Four numerical examples are given to verify the accuracy and practicability of the method. The results show that indeed the method converges with second order of accuracy. One important lesson from this study is that the results support the claim, in previous studies, that fixed point iterations can provide reliable means of solving several nonlinear problems. It is recommended to extend this work to functional integral equations using higher order quadrature rules.

*Corresponding author: E-mail: nwaigwe.chinedu@ust.edu.ng;

Keywords: Integral equations; ishikawa iteration; discrete fixed-point algorithms; trapezoidal rule; experimental order of convergence.

2020 Mathematics Subject Classifications: 65R20, 45D05.

1 Introduction

An equation in which the unknown solution appears under an integral is known as an integral equation. This type of equations occur in several applications including physical sciences [1, 2, 3], economics and optimal control [4, 5, 6, 7], epidemiology [8, 9], heat transfer [10], antenna wire modelling [11, 12], neuron transport [13] and applied mathematical methods [14, 15]. More applications can be seen in [16, 17, 18, 19, 20].

For practical applications these equations are highly nonlinear and exact analytic solutions cannot be found. This necessitates the development of numerical methods for nonlinear integral equations. In this study we are interested in Fredholm equations which have applications in many areas including antenna wire modelling [11, 12]. Specifically, we are interested in approximating the solution of the following problem:

$$u(x) = g(x) + \int_{y=a}^b k(x, y, u(y)) dy, \quad x \in [a, b] \subset \mathbb{R}, \quad (1)$$

where $C[a, b] \ni g : [a, b] \rightarrow \mathbb{R}$, $k : [a, b] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and k is nonlinear in u and may or may not be separable in x and y . Further, we assume that k is Lipschitz continuous with respect to its third argument, namely there exists α_k such that

$$|k(x, y, u) - k(x, y, v)| \leq \alpha_k |u - v| \quad \forall \quad u, v.$$

In general, closed form analytical solutions do not exist hence numerical techniques [21, 22, 23, 24, 25, 26] are being developed to tackle the most general cases that may arise from practical applications. Linear Fredholm integral equations have been discussed in [27, 28, 29] whereas the nonlinear Fredholm integral equation poses more challenging problem. The use of Haar Wavelet method has been applied for nonlinear Fredholm equation equations [30], however a brief study of the papers cited in [31] have shown various complications and its usage not generally effective. In other studies [30, 32] investigated another approach for solving the nonlinear second kind of Fredholm equations, while He's homotopy perturbation method is used to find the solution of non-linear Fredholm equations in [33].

The existence and uniqueness of solution of functional Volterra equations is investigated in [34]. The solution was approximated by using nonlinear solver of Newton-Raphson type. The idea is extended to mixed Volterra-Fredholm equations by Nwaigwe and Benedict [21]. Nwaigwe [35] further extended the work to nonlinear functional mixed equations. Some remarkable contributions of the work of Nwaigwe [35] are the proof of existence and uniqueness of solution without any contraction assumption, and the propounding of new Grownwall inequalities which were used to prove the convergence of their proposed numerical scheme.

The challenge and downside of the above works are the use of nonlinear solvers which presents difficulties in computation, programming and mathematical theories. Hence, Nwaigwe [36] adopted the idea of fixed point iterations [15, 37] to propose a discrete Kranoselkij algorithm for functional Voltera equations without solving nonlinear systems. The results show that such algorithms produce very accurate solutions and are fast. A very recent study in using iterative methods to solve integral equations can be found in [38]. This has motivated us to look into approaching problem (1) through the use of fixed point methods. Obviously, fixed point methods have been widely investigated [39, 40, 41] and applied to variety of problems.

To the best of our knowledge, no study has investigated the combined use of collocation projection, trapezoid rule and Ishikawa iteration to solve Fredholm integral equations; this is the gap that the current work fills. Hence, the present study is to join the efforts of [15, 37, 36, 38], among others, to extend the application of fixed point methods to solve nonlinear Fredholm equations. The paper is organized as follows. In section 2 we recall

the Ishikawa iteration, formulate the numerical algorithm and give the pseudocode. The results are presented and discussed in section 3 and we give the concluding remarks in section 4.

2 The Numerical Method

We start this section by recalling the fixed point method of Ishikawa [42].

Lemma 2.1 (see Ishikawa [42]). *Let T be a contractive self-map, then the sequence, $\{u_n\}_{n \geq 0}$, generated by*

$$u_{n+1}(x) = (1 - a_n)u_n(x) + a_n T v_n(x) \quad (2)$$

$$v_n(x) = (1 - b_n)u_n(x) + b_n T u_n(x) \quad (3)$$

for a suitable u_0 , where (a_n) and (b_n) are in $(0, 1)$, converges to the fixed point of T .

In this work, we set $a_n = b_n = n^{-1/2}$ [42] and define the map T by

$$(Tu)(x) := g(x) + \int_{y=a}^b k(x, y, u(y)) dy. \quad (4)$$

Hence, the Fredholm equation (1) can be written in the operator form:

$$u(x) = (Tu)(x), \quad \text{for } x \in [a, b]. \quad (5)$$

We assume that the function k is Lipchitz continuous in $[a, b]$ with Lipchitz constant α_k and

$$\alpha_k(b - a) \leq 1. \quad (6)$$

Notice the equality sign included in the inequality above, it is because the Ishikawa iteration, like the Krasnoselskij iteration still converge for non-expansive maps.

Let $N \in \mathbb{Z}^+$ with $N > 1$. Define the $h = \frac{b-a}{N}$, $x_i = a + ih : i = 0, 1, \dots, N$, and let $u_{n,i}$ be the approximation of $u(x_i)$ after $n = 0, 1, 2, \dots$, Ishikawa iterations. By collocating at the grid points, we have

$$u(x_i) = (Tu)(x_i), \quad \text{for } i = 0, 1, \dots, N. \quad (7)$$

Applying the Ishikawa iteration, we obtain

$$u_{n+1,i} = (1 - a_n)u_{n,i} + a_n T v_{n,i}, \quad (8)$$

$$v_{n,i} = (1 - b_n)u_{n,i} + b_n T u_{n,i}, \quad n \geq 0, \quad (9)$$

where

$$T v_{n,i} = g(x_i) + I_{n,i}, \quad \text{where } I_{n,i} = \int_a^b k(x_i, y, u_n(y)) dy. \quad (10)$$

The integral is approximated by the use of the quadrature rule (see [43, 21]):

$$I_{u,n,i} = \frac{h}{2} \sum_{j=0}^{N-1} (k(x_i, x_j, u_{n,j}) + k(x_i, x_{j+1}, u_{n,j+1})). \quad (11)$$

The algorithm is initialize with

$$u_{0,i} = g(x_i) + \frac{h}{2} \sum_{j=0}^{N-1} (k(x_i, x_j, g(x_j)) + k(x_i, x_{j+1}, g(x_{j+1}))) \quad \forall i. \quad (12)$$

2.1 Pseudocode for the Algorithm

- (1) Define the grid, $\Omega_h = \{x_i, x_i = a + ih, h = \frac{b-a}{N}, \mathbb{Z} \ni N > 1\}$.
- (2) For each i compute $u_{0,i}$ using (12).
- (3) Set $n = 0$:
- (4) For each i , do the following:
 - (a) Evaluate the integral using (11).
 - (b) Compute $Tu_{n,i}$ using (10).
 - (c) Compute $u_{n+1,i}$ using (8).
 - (d) If $i < N$, set $i = i + 1$ and go to step 4a, otherwise go to the next step.
 - (e) If $\|u_{n+1} - u_n\| < TOL$ output u_{n+1} and stop. Otherwise, set $n = n + 1$ and go to step 4.

3 Numerical Experiments

We now present some examples to demonstrate the accuracy and practicality of the proposed method. The results are computed on 16 Gigabyte RAM quadcore machine running on a Linux operating system. These examples are derived via the method of Manufactured solutions [44, 45, 46, 47, 48, 49, 50].

Example 1

The first example is the problem:

$$u(x) = \frac{\frac{-x}{6} + (x^2 + 1)\cos^{-1}(x^2)}{1 + x^2} + \int_0^1 \frac{xy^3 \cos(u)}{1 + x^2} dy,$$

The exact solution of this problem is $u(x) = \cos^{-1}(x^2)$. The numerical solution of this problem is computed on a sequence of grids and their errors and experimental order of convergence (EOC) computed. The CPU time to compute all the EOCs is also recorded. The results are recorded in Table 1, it can be seen that proposed method is convergent and has second order of accuracy which is the theoretically expected order of accuracy. The CPU time taken to complete the EOC calculations of this problem 1 is 42.97808265686035.

Further, the numerical solution computed on different grids with 5, 10, 30 and 100 mesh points are plotted and compared with the exact solution in Fig. 1. We see that even with 30 mesh points the computed solution is already very accurate.

Table 1. Results for Problem 1, N = number of sub-intervals

N	Error	EOC
2	0.16333911635262954	-1
4	0.021646255808007164	2.915680921934721
8	0.003967379363779777	2.4478592401930115
16	0.0008612525713960623	2.203678065343175
32	0.00020136113788714327	2.0966511109303756
64	4.872781396353387e-05	2.046967867635893
128	1.1988354706990445e-05	2.0231118260012755
256	2.9733981802698706e-06	2.0114490968584344
512	7.404221460394211e-07	2.005692721549541
1024	1.8474197853446395e-07	2.002836322731808




Fig. 1. Plot of exact solution and numerical solution for problem 1

Table 2. Results for problem 2, N = number of sub-intervals

N	Error	EOC
2	0.05637482608295197	-1
4	0.0037625705430824885	3.905260541833301
8	0.00061178604074974	2.620619536587277
16	0.00012800706701432407	2.2568037282629883
32	2.9747578815797482e-05	2.1053793047664726
64	7.1905011685968745e-06	2.0486080176014885
128	1.7687111649733467e-06	2.0233938566428216
256	4.386726273342134e-07	2.0114818797252187
512	1.0923665838546981e-07	2.0056875996706625
1024	2.7255633971989823e-08	2.0028307037917417

Example 2

The second example is

$$u(x) = x^2 \left(x - \frac{\pi\sqrt{3}}{54} \right) + \int_0^1 \frac{x^2 y^2 u}{1 + u^2 + u^4} dy$$

The exact solution is $u(x) = x^3$

The solution of this problem is computed on different grids and the results are tabulated in Table 2. Just like in example 1, we can see that the proposed method converges to the exact solution at the expected order of convergence, 2. The CPU time taken to complete problem 2 is 22.412165880203247. Fig. 2. presents the plots of the exact solution and the numerical solution computed of different grids. We also see that the numerical solution converges very fast to the exact solution.




Fig. 2. Plot of exact solution and numerical solution for problem 2

Table 3. Results for problem 3, N = number of sub-intervals

N	Error	EOC
2	0.0033401995652793115	-1
4	0.0003850850798006178	3.1166851694545703
8	7.08155916803177e-05	2.4430382838362124
16	1.542482707894699e-05	2.1988127217211035
32	3.611565474237466e-06	2.0945580856224573
64	8.744631218027266e-07	2.046154876652523
128	2.151870582833837e-07	2.02280622917233
256	5.337573516506211e-08	2.011335373573074
512	1.329181165132809e-08	2.00564628078104
1024	3.316511398843147e-09	2.0027993653605405

Example 3

Our third problem is

$$u(x) = -\frac{x^2}{12} + \sin(x) - \frac{\sin(2)}{48} - \frac{1}{54} - \frac{\cos(3)}{432} + \frac{\cos(1)}{48} + \frac{1}{12} \int_0^1 \left(x^2 + y + \frac{u^3}{3} - u^2 \right) dy.$$

The exact solution is $u(x) = \sin(x)$. Table 3. tabulates the results computed for this problem using the proposed method on a sequence of grids. We also see that the numerical solution converges to the exact solution. This is also seen in Fig. 3. which displays the plots of the exact solution and numerical solution on several grids. The CPU time taken to compute the EOCs for problem 3 is 23.757.




Fig. 3. Plot of Exact solution and numerical solution for problem 3

Example 4

Our last example is

$$u(x) = e^{\frac{(x-1)}{2}} - \frac{1}{2}e^{x-e^2} + e^{-x} + \int_0^1 ue^{x-y-u^2} dy.$$

The exact solution is $u(x) = e^{-x}$.

The results for this problem are displayed in Table 4. and Fig. 4. One can also see that the numerical solution agrees with the exact solution as we refine the mesh. The CPU time taken to compute the EOCs for this problem is 41.409.

Table 4. Results for problem 4, N = number of sub-intervals

N	Error	EOC
2	0.03112031852815711	-1
4	0.005963255741055273	2.383684712094301
8	0.0011218506673921214	2.410219562070775
16	0.0002452206271312818	2.193728403766116
32	5.7458015915723415e-05	2.0935002576668724
64	1.3914595630604154e-05	2.045909192594972
128	3.4242276554241613e-06	2.0227484576080528
256	8.49363330335251e-07	2.0113248911773134
512	2.115096261467997e-07	2.005658500148919
1024	5.277262404845828e-08	2.0028616977181466




Fig. 4. Plot of exact solution and numerical solution for problem 4

4 Conclusion

A numerical method based on Ishikawa iteration has been proposed and implemented for solving nonlinear Fredholm integral equations with non-separable kernels. The algorithm is verified with four test problems with known exact solutions. The results show that

- (i) the scheme converges to the exact solution at the correct theoretical order of convergence,
- (ii) the scheme is fast and easy to program, and
- (iii) the quality of solution does not degrade as the mesh is refined.

We, therefore, conclude that provided the hypotheses of fixed point theorems are not violated, iterative schemes like the Ishikawa iteration can lead to a very reliable method for solving nonlinear integral equations. We recommended a further study to extend this work to functional integral equations using higher order quadrature rules.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Abdou, MA. On a asymptotic methods for Fredholm-Volterra integral equation of the second kind in contact problems. Journal of Computational and Applied Mathematics. 2003;154(2):431-446. Elsevier
- [2] Le, Tien-Dung, Moyne, Christian, Murad, MA, Lima, SA. A two-scale non-local model of swelling porous media incorporating ion size correlation effects. Journal of the Mechanics and Physics of Solids. 2013;61(12):2493-2521. Elsevier

- [3] Rocha, Aline C, Murad, Marcio A, Moyne, Christian, Oliveira, Saulo P, Le, Tien D. A new methodology for computing ionic profiles and disjoining pressure in swelling porous media. *Computational Geosciences*. 2016;20(5):975-996. Springer
- [4] Hu, Shouchuan, Khavanin, Mohammad, Zhuang, WAN. Integral equations arising in the kinetic theory of gases. *Applicable Analysis*. 1989;34(3-4):L261-266. Taylor & Francis
- [5] Jerri, Abdul. Introduction to integral equations with applications. John Wiley & Sons; 1999.
- [6] Oregan, Donal. Existence results for nonlinear integral equations. *Journal of Mathematical Analysis and Applications*. 1995;192(3):705-726. Elsevier
- [7] Prüss, Jan. Evolutionary integral equations and applications. Birkhäuser. 2013;87.
- [8] Maleknejad, K, Hadizadeh, M. A new computational method for Volterra-Fredholm integral equations. *Computers & Mathematics with Applications*. 1999;37(9):1-8. Elsevier
- [9] Wazwaz, Abdul-Majid. A reliable treatment for mixed Volterra-Fredholm integral equations. *Applied Mathematics and Computation*. 2002;127(2-3):405-414. Elsevier
- [10] Olmstead, WE, Handelsman, Richard A. Asymptotic solution to a class of nonlinear Volterra integral equations II. *SIAM Journal on Applied Mathematics*. SIAM. 1976;30(1):180-189.
- [11] Alipanah, Amjad. Solution of hallen's integral equation by using radial basis functions. *Mathematical Reports*. Editura Acad Romane Calea 13 Septembrie Nr 13, Sector 5, Bucuresti 050711. 2013;15(3):211-220.
- [12] Mei, K. On the integral equations of thin wire antennas. *IEEE Transactions on Antennas and Propagation*. 1965;13(3):374-378. IEEE
- [13] Argyros, Ioannis K. On a class of nonlinear integral equations arising in neutron transport. *Aequationes Mathematicae*. 1988;36:99-111. Springer
- [14] Wazwaz, Abdul-Majid. First course in integral equations, A. World Scientific Publishing Company; 2015.
- [15] Micula, Sanda. On some iterative numerical methods for mixed volterra-fredholm integral equations. *Symmetry*. Multidisciplinary Digital Publishing Institute. 2019;11(10):1200.
- [16] Honerkamp, Josef , Weese, Jürgen. Tikhonovs regularization method for ill-posed problems: A comparison of different methods for the determination of the regularization parameter. *Continuum Mechanics and Thermodynamics*. 1990;2:17-30. Springer
- [17] Ray, Saha S, Sahu PK. Numerical methods for solving Fredholm integral equations of second kind. *Book Abstract and Applied Analysis*; 2013. Hindawi
- [18] Schäfer H, Sternin E, Stannarius R, Arndt M, Kremer F. Novel approach to the analysis of broadband dielectric spectra. *Physical Review Letters*. 1996;76(12):2177. APS
- [19] Daddi-Moussa-Ider, Abdallah, Kaoui, Badr, Löwen, Hartmut. Axisymmetric flow due to a Stokeslet near a finite-sized elastic membrane. *Journal of the Physical Society of Japan*. The Physical Society of Japan. 2019;88(5):054401.
- [20] Asymmetric Stokes flow induced by a transverse point force acting near a finite-sized elastic membrane. Daddi-Moussa-Ider, Abdallah. *Journal of the Physical Society of Japan*. The Physical Society of Japan. 2020;89(12):124401.
- [21] Nwaigwe, Chinedu, Benedict, Deborah Ngochinma. Generalized Banach fixed-point theorem and numerical discretization for nonlinear Volterra-Fredholm equations. *Journal of Computational and Applied Mathematics*. 2022;115019. Elsevier
- [22] Manochehr, Kazemi. Triangular functions for numerical solution of the nonlinear Volterra Integral Equations. *Journal of Applied Mathematics and Computing*. 2021;1-24. Springer
- [23] Ordokhani, Yadollah and Razzaghi, Mohsen. Solution of nonlinear Volterra-Fredholm-Hammerstein integral equations via a collocation method and rationalized Haar functions. *Applied Mathematics Letters*. 2008;21(1):4-9. Elsevier

- [24] Brunner, Hermann. On the numerical solution of nonlinear Volterra-Fredholm integral equations by collocation methods. *SIAM Journal on Numerical Analysis*. 1990;27(4):987-1000. SIAM
- [25] Ezzati R, Najafalizadeh S. Numerical methods for solving linear and nonlinear Volterra-Fredholm integral equations by using CAS wavelets. *World Applied Sciences Journal*. 2012;18(12):1847-1854. Citeseer
- [26] Shiralashetti, SC, Lamani, Lata. CAS wavelets stochastic operational matrix of integration and its application for solving stochastic Itô-Volterra integral equations. *Jordan Journal of Mathematics and Statistics (JJMS)*. 2021;14(3):555-580.
- [27] Delves, LM, Abdel-Al, LF. The fast galerkin algorithm for the solution of linear fredholm equations; 1976. PRE-20036
- [28] Delves, LM. A fast method for the solution of Fredholm integral equations. *IMA Journal of Applied Mathematics*. Oxford University Press. 1977;20(2):173-182.
- [29] Delves, Leonard Michael, Mohamed, Julie L. *Computational methods for integral equations*; 1988. CUP Archive
- [30] Babolian, E, Shamsavaran, A. Numerical solution of nonlinear Fredholm integral equations of the second kind using Haar wavelets. *Journal of Computational and Applied Mathematics*. 2009;225(1):87-95. Elsevier
- [31] Lepik, Ülo, Tamme, Enn. Application of the Haar wavelets for solution of linear integral equations. *Proceedings*. 2004;494-507.
- [32] Borzabadi, Akbar Hashemi, Kamyad, Ali Vahidian, Mehne, Hamed Hashemi. A different approach for solving the nonlinear Fredholm integral equations of the second kind. *Applied Mathematics and Computation*. 2006;173(2):724-735. Elsevier
- [33] Biazar, Jafar, Ghazvini, H. Numerical solution for special non-linear Fredholm integral equation by HPM. *Applied Mathematics and Computation*. 2008;195(2):681-687. Elsevier
- [34] Analysis of the Euler and trapezoidal discretization methods for the numerical solution of nonlinear functional Volterra integral equations of Urysohn type. Bazm, Sohrab, Lima, Pedro, Nemati, Somayeh. *Journal of Computational and Applied Mathematics*. 2021;113628. Elsevier
- [35] Nwaigwe, Chinedu. Solvability and approximation of nonlinear functional mixed volterra-fredholm equation in banach space. *Journal of Integral Equations and Applications*. 2022;34(4):489-500.
- [36] Chinedu Nwaigwe, Sanda Micula. Fast and accurate numerical algorithm with performance assessment for nonlinear functional volterra equations. 2023.
Available: https://www.researchgate.net/publication/361722935_Fast_and_Accurate_Numerical_Algorithm-with_Performance_Assessment_for_Nonlinear_Functional_Volterra_Equations
- [37] Micula, Sanda. Numerical solution of two-dimensional fredholm-volterra integral equations of the second kind. *Symmetry. Multidisciplinary Digital Publishing Institute*. 2021;13(8)1326.
- [38] Micula, Sanda. Iterative numerical methods for a fredholm-hammerstein integral equation with modified argument. *Symmetry. Multidisciplinary Digital Publishing Institute*. 2023;15(1):66.
- [39] Okeke, Godwin Amechi. Convergence analysis of the Picard-Ishikawa hybrid iterative process with applications. *Afrika Matematika*. 2019;30(5):817-835. Springer
- [40] Berinde, Vasile. *Iterative approximation of fixed points*; 2007. Springer
- [41] Osilike, Micah, Chidume, Charles. *Nonlinear accretive and pseudo-contractive operator equations in Banach spaces*; 1996.
- [42] Shiro Ishikawa. Fixed points by a new iteration method. *Proceedings of the American Mathematical Society*. 1974;44(1):147.
- [43] Burden, Richard L. *Numerical analysis*. Boston, MA: Brooks/Cole, Cengage Learning; 2011.

- [44] Nwaigwe, Chinedu. An unconditionally stable scheme for two-dimensional convection-diffusion-reaction equations; 2022.
Available: https://www.researchgate.net/publication/357606287_An_Unconditionally_Stable_Scheme_for_Two-Dimensional_Convection-Diffusion-Reaction_Equations
- [45] Nwaigwe, Chinedu, Makinde, Oluwole Daniel. Finite difference investigation of a polluted non-isothermal non-newtonian porous media flow. Diffusion Foundations. 2019;26(4):145-156.
- [46] Nwaigwe, C, Ndu, RI, Weli, A. Wall motion effects on channel flow with temperature-dependent transport properties. Applied Mathematics. 2019;9(3)162-168.
- [47] Weli, Azubuike, Nwaigwe, Chinedu. Numerical analyses of channel flow with velocity-dependent suction and nonlinear heat source. Journal of Interdisciplinary Mathematics. Taylor and Francis. 2020;23(5):987-1008.
DOI: 10.1080/09720502.2020.1748278
- [48] Analysis and application of a convergent difference scheme to nonlinear transport in a Brinkman flow. International Journal of Numerical Methods for Heat & Fluid Flow. 2020; 30(10): 4453-4473.
DOI: <https://doi.org/10.1108/HFF-10-2019-0758>
- [49] Nwaigwe, Chinedu, Weli, Azubuike, Makinde, Oluwole Daniel. Computational analysis of porous channel flow with cross-diffusion. American Journal of Computational and Applied Mathematics. 2019;9(5):119-132.
DOI: 0.5923/j.ajcam.20190905.01
- [50] Analysis of two finite difference schemes for a channel flow problem. Nwaigwe, Chinedu, Weli, Azubuike. Asian Research Journal of Mathematics. 2019;15:1-14.

© 2023 Nwaigwe and Weli; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/96672>