Policy 533-07

| ANAPHYLAXIS / ALLERGIC REACTION | |
|---|--|
| ADULT | PEDIATRIC - (14 years and under) |
| BLS | Procedures |
| Assist with prescribed Epi Auto-Injector Administer oxygen as indicated Refer to Policy 533-02 Airway Management Epinephrine 1mg/mL - Auto-Injector IM - 0.3mg | Assist with prescribed Epi Auto-Injector Administer oxygen as indicated Refer to Policy 533-02 Airway Management Epinephrine 1mg/mL - Auto-Injector Patient weighs 15kg - 30kg IM - 0.15mg (Max total dose of 0.15mg) Patient weighs > 30kg IM - 0.3mg (Max total dose of 0.3mg) |
| Ехра | nded Scope |
| Epinephrine 1mg/mL − Auto-Injector or Syringe/Vial • IM − 0.3mg (0.3mL) ∘ May repeat x1 for ongoing distress | Epinephrine 1mg/mL – Auto-Injector or Syringe/Vial IM – 0.01mg/kg Patient weighs 15kg – 30kg IM – 0.15mg (Max total dose of 0.15mg) Patient weighs > 30kg IM – 0.3mg (Max total dose of 0.3mg) |
| ALS Prior to B | Base Hospital Contact |
| Vascular Access Cardiac monitor | Vascular Access Cardiac monitor |
| Anaphylaxis Without Shock Epinephrine 1mg/mL • IM − 0.3mg (0.3mL) ∘ May repeat every 5 mins x2 for ongoing distress | Anaphylaxis Without Shock Epinephrine 1mg/mL • IM – 0.01mg/kg • Patient weighs 15kg – 30kg • IM – 0.15mg (Max total dose of 0.15mg) • Patient weighs > 30kg • IM – 0.3mg (Max total dose of 0.3mg) • Contact BH for further |
| Anaphylaxis With Shock Same as above (anaphylaxis without shock) Initiate 2nd IV/IO Normal Saline IV/IO bolus – 1L | Anaphylaxis With Shock • Same as above (anaphylaxis without shock) • Initiate 2nd IV/IO • Normal Saline • IV/IO bolus – 20mL/kg |
| For Hypotension Refractory to Normal Saline Push-Dose Epinephrine Refer to Policy 533-10: Push-Dose Epinephrine | For Hypotension Refractory to Normal Saline Push-Dose Epinephrine Refer to Policy 533-10: Push-Dose Epinephrine |
| If Wheezing Is Present Albuterol • Nebulizer – 5mg (6mL) ○ Repeat as needed | If Wheezing Is Present Albuterol • Age < 2 years – 2.5mg (3mL) • Age ≥ 2 years – 5mg (6mL) • Repeat as needed |
| Allergic Reaction (Hives Only) Diphenhydramine • IM – 50mg | Allergic Reaction (Hives Only) Diphenhydramine • IM – 1mg/kg (max 50mg) |
| Base Hospital F | Physician Orders Only |
| Consult with ED Physician for further treatment measures | Consult with ED Physician for further treatment measures |
| Addition | nal Information |
| Anaphylaxis is a <u>true</u> medical emergency requiring immediate assessment, recognition, and intervention using Epinephrine It is pivotal to treat these patients with IM Epinephrine Once anaphylaxis is stabilized, continue to recognize and treat signs/symptoms throughout transport. | assessment, recognition, and intervention using Epinephrine It is pivotal to treat these patients with IM Epinephrine |

Last Reviewed/Revised: December 31, 2023 Next Review Date: December 31, 2025

signs/symptoms throughout transport.

Effective Date: April 1, 2024

Signature on File

signs/symptoms throughout transport.